

**Pure Mathematics**

**Differential & Integral Calculus**



# Question Bank & Practice Exams



By a group of supervisors

3<sup>rd</sup>  
Sec  
2022





# Question Bank & Practice Exams

3<sup>rd</sup>  
Sec

**AL TALABA BOOKSTORE**

For printing, publication &amp; distribution

El Faggala - Cairo - Egypt

Tel.: 02/ 259 340 12 - 259 377 91

E-mail: info@elmoasserbooks.com

www.elmoasserbooks.com

15014

## جميع حقوق الطبع والنشر محفوظة

لا يجوز، بأي صورة من الصور، التوزيع (النقل) المباشر أو غير المباشر لأي مما ورد في هذا الكتاب أو نسخه أو تصويره أو ترجمته أو تحويله أو الاقتباس منه أو تحويله رقميًا أو إنتاجه عبر شبكة الإنترنت إلا بإذن كتابي مسبق من الناشر. كما لا يجوز بأي صورة من الصور استخدام العلامة التجارية (EL-MOASSER) المسجلة باسم الناشر ومن يخالف ذلك يتعرض للمساءلة القانونية طبقاً لأحكام القانون ٨٢ لسنة ٢٠٠٢ الخاص بحماية الملكية الفكرية.



## Cataloging - in - Publishing

Prepared by Technical Affairs Department  
Egyptian National Library

El-Moasser Question Bank & Practice Exams /

By a group of supervisors. - 1<sup>st</sup> ed. - Cairo :

GPS for printing , publication and distribution , 2021-2022

2 vols ; 30 cm.

For 3<sup>rd</sup> Sec.

I.S.B.N. : 978 - 977- 475 - 86 - 70

1. Mathematics - study and teaching (secondary).

2. Education , secondary

510.7

Dep. No. 9508/2022



## *Preface*

Thanks to God who helped us to introduce one of our famous series "El Moasser" in mathematics.

We introduce this book to our colleagues.

We also introduce it to our students to help them study mathematics.

In fact, this book is the outcome of more than thirty years experience in the field of teaching mathematics.

This book will make students aware of all types of questions.

We would like to know your opinions about the book hoping that it will win your admiration.


We will be grateful if you send us your recommendations and your comments.

---

*The Authors*



# CONTENTS



- Summary for differential & integral calculus.
- Multiple choice question bank.
- Practice exams.
- School book examinations.
- Egypt exams (2017 : 2021 first and second sessions).
- Al-Azhar exams (2019 : 2021 first and second sessions).



# Summary

for

## Differential & Integral calculus







## Important notes related to (e)

The Nabier's constant  $e$  is an irrational number,  $2 < e < 3$ ,


$$e = 1 + \frac{1}{1} + \frac{1}{2} + \frac{1}{3} + \dots = \sum_{n=0}^{\infty} \frac{1}{n} \quad (\text{Taylor's series})$$

$$e = 2.71828$$

### • The natural exponential function :

$$f : \mathbb{R} \longrightarrow \mathbb{R}^+ \text{ where } f(x) = e^x$$

is exponential function whose base is  $e$  and it is one - to - one its domain  $= \mathbb{R}$ , its range  $= \mathbb{R}^+$ , its curve passes through  $(0, 1)$ ,  $(1, e)$



### • The natural logarithmic function :

$$f : \mathbb{R}^+ \longrightarrow \mathbb{R} \text{ where } f(x) = \ln x$$

is logarithmic function of base "e" ( $\log_e x = \ln x$ ) and it is one - to - one function its domain  $\mathbb{R}^+$ , its range  $= \mathbb{R}$ , its curve passes through  $(1, 0)$ ,  $(e, 1)$

## Remarks

From the previous we get :  $\lim_{x \rightarrow \infty} \ln x = \infty$ ,  $\lim_{x \rightarrow 0^+} \ln x = -\infty$

### ★ Some properties of natural logarithm :

if  $x, y \in \mathbb{R}^+$ ,  $n \in \mathbb{R}$  under condition the base of the logarithm  $\in \mathbb{R}^+ - \{1\}$ , then :

1.  $\ln e = 1$

2.  $\ln 1 = \text{zero}$

3.  $\ln x^n = n \ln x$

4.  $e^{\ln x} = x$

5.  $\ln xy = \ln x + \ln y$

6.  $\ln \frac{x}{y} = \ln x - \ln y$

7.  $\log_y x = \frac{\ln x}{\ln y}$

8.  $\ln x \times \log_x e = 1$



## Limits of functions related to the number e

$$1. \lim_{x \rightarrow \infty} \left(1 + \frac{1}{x}\right)^x = e \quad \text{and} \quad \lim_{x \rightarrow 0} \left(1 + x\right)^{\frac{1}{x}} = e$$

## Remarks

$$1. \lim_{x \rightarrow \infty} \left(1 + \frac{1}{x}\right)^{ax} = e^a$$

$$2. \lim_{x \rightarrow \infty} \left(1 + \frac{1}{x}\right)^{x+a} = e$$

$$3. \lim_{x \rightarrow \infty} \left(1 + \frac{a}{x}\right)^x = e^a$$

$$4. \lim_{x \rightarrow \infty} \left(1 - \frac{a}{x}\right)^{x+k} = e^{-a}$$

$$5. \lim_{x \rightarrow \infty} \left(1 + \frac{a}{x}\right)^{\frac{x}{a}} = e$$

$$2. \lim_{x \rightarrow 0} e^x = 1$$

$$3. \lim_{x \rightarrow 0} \frac{\log_a (1+x)}{x} = \log_a e \quad \text{and} \quad \lim_{x \rightarrow 0} \frac{\ln (1+x)}{x} = 1$$

$$4. \lim_{x \rightarrow 0} \frac{a^x - 1}{x} = \ln a \quad \text{and} \quad \lim_{x \rightarrow 0} \frac{e^x - 1}{x} = 1$$

## Derivative

## ★ Derivative of exponential and logarithmic functions :

$$1. \text{ If } y = e^x, \text{ then } \frac{dy}{dx} = e^x$$

$$2. \text{ If } y = a^x, \text{ then } \frac{dy}{dx} = a^x \ln a$$

$$3. \text{ If } y = \ln x, \text{ then } \frac{dy}{dx} = \frac{1}{x}$$

$$4. \text{ If } y = \log_a x, \text{ then } \frac{dy}{dx} = \frac{1}{x} \log_a e$$

## ★ In general :

$$1. \text{ If } y = e^{f(x)}, \text{ then } \frac{dy}{dx} = f'(x) e^{f(x)}$$

$$2. \text{ If } y = a^{f(x)}, \text{ then } \frac{dy}{dx} = f'(x) a^{f(x)} \cdot \ln a$$

$$3. \text{ If } y = \ln f(x), \text{ then } \frac{dy}{dx} = \frac{f'(x)}{f(x)}$$

$$4. \text{ If } y = \log_a f(x), \text{ then } \frac{dy}{dx} = \frac{f'(x)}{f(x)} \times \log_a e$$

**Notice that**

$$\odot \text{ If } y = \ln |f(x)|, \text{ then } \frac{dy}{dx} = \frac{f'(x)}{f(x)}$$

$$\odot \text{ If } y = \log_a |f(x)|, \text{ then } \frac{dy}{dx} = \frac{f'(x)}{f(x)} \log_a e$$

**★ Differentiation of trigonometric functions :**

$$1. \text{ If } y = \sin x, \text{ then } \frac{dy}{dx} = \cos x$$

$$2. \text{ If } y = \cos x, \text{ then } \frac{dy}{dx} = -\sin x$$

$$3. \text{ If } y = \tan x, \text{ then } \frac{dy}{dx} = \sec^2 x$$

$$4. \text{ If } y = \cot x, \text{ then } \frac{dy}{dx} = -\csc^2 x$$

$$5. \text{ If } y = \sec x, \text{ then } \frac{dy}{dx} = \sec x \tan x$$

$$6. \text{ If } y = \csc x, \text{ then } \frac{dy}{dx} = -\csc x \cot x$$

**★ In general :**

$$1. \text{ If } y = \sin(f(x)), \text{ then } \frac{dy}{dx} = f'(x) \cos(f(x))$$

$$2. \text{ If } y = \cos(f(x)), \text{ then } \frac{dy}{dx} = -f'(x) \sin(f(x))$$

$$3. \text{ If } y = \tan(f(x)), \text{ then } \frac{dy}{dx} = f'(x) \sec^2(f(x))$$

$$4. \text{ If } y = \cot(f(x)), \text{ then } \frac{dy}{dx} = -f'(x) \csc^2(f(x))$$

$$5. \text{ If } y = \csc(f(x)), \text{ then } \frac{dy}{dx} = -f'(x) \csc(f(x)) \cot(f(x))$$

$$6. \text{ If } y = \sec(f(x)), \text{ then } \frac{dy}{dx} = f'(x) \sec(f(x)) \tan(f(x))$$

**Implicit differentiation**

The derivative of a relation between two (or more) variables with respect to one of them without separating them.

**Notice**  $\frac{d x^n}{d x} = n x^{n-1}$  but  $\frac{d y^n}{d x} = n y^{n-1} \cdot \frac{dy}{dx}$

**Parametric differentiation**

If  $y = f(t)$ ,  $x = g(t)$  are the parametric equations of a curve, then :

$$\frac{dy}{dx} = \frac{dy}{dt} \times \frac{dt}{dx} = \frac{dy}{dt} \div \frac{dx}{dt}$$



### Higher derivatives of the function

The derivatives starting from the second derivative are called higher derivatives and the  $n^{\text{th}}$  derivative is denoted as  $y^{(n)} = \frac{d^n y}{d x^n} = f^{(n)}(x)$ , where  $n$  is a positive integer.

#### Notice that

$$\odot \frac{d y}{d x} \times \frac{d x}{d y} = 1 \quad \text{but} \quad \frac{d^2 y}{d x^2} \times \frac{d^2 x}{d y^2} \neq 1$$

$$\odot \frac{d y}{d z} \times \frac{d z}{d x} = \frac{d y}{d x} \quad \text{but} \quad \frac{d^2 y}{d z^2} \times \frac{d^2 z}{d x^2} \neq \frac{d^2 y}{d x^2}$$

"Chain rule can not be applied to find second derivative"

$$\odot \frac{d}{d x} \left[ y \frac{d y}{d x} \right] = 1^{\text{st}} \times \text{derivative of the } 2^{\text{nd}} + 2^{\text{nd}} \times \text{derivative of the } 1^{\text{st}}$$

$$= y \frac{d}{d x} \left( \frac{d y}{d x} \right) + \frac{d y}{d x} \times \frac{d y}{d x} = y \frac{d^2 y}{d x^2} + \left( \frac{d y}{d x} \right)^2$$

$$\odot \text{The rate of change of the slope of the tangent to the curve } y = f(x) \text{ equals } \frac{d}{d x} \left( \frac{d y}{d x} \right) = \frac{d^2 y}{d x^2}$$

$$\odot \text{If } y = (f \circ g)(x) = f(g(x)), \text{ then } \frac{d y}{d x} = \dot{f}(g(x)) \cdot \dot{g}(x)$$

$$\odot \text{If } y = \sin a x, \text{ then } y^{(n)} = a^n \sin \left( a x + \frac{n\pi}{2} \right)$$

$$\odot \text{If } y = \cos a x, \text{ then } y^{(n)} = a^n \cos \left( a x + \frac{n\pi}{2} \right)$$

$$\odot \text{If } y = \sin a x \text{ or } y = \cos a x, \text{ then } y^{(n)} = a^n y \text{ where } n \text{ is divisible by } 4$$

### Applications on first derivative (The equations of the tangent and the normal to a curve)

If  $A(x_1, y_1)$  is a point on the curve  $y = f(x)$ , then :

1. The slope of the tangent to the curve at A

$$= \left( \frac{d y}{d x} \right)_{(x_1, y_1)}$$

2. The slope of the normal to the curve at A

$$= \frac{-1}{\left( \frac{d y}{d x} \right)_{(x_1, y_1)}}$$

3. The tangent equation at A :  $y - y_1 = m(x - x_1)$


$$\text{and the normal equation : } y - y_1 = \frac{-1}{m}(x - x_1)$$

#### Remark

- ⊙ The slope of the curve at a point on it is the slope of the tangent to the curve at this point.
- ⊙ The normal to the curve is the straight line perpendicular to the tangent at the point of tangency.

**Related time rates**

- ⊙ If we have a relation between several variables  $X, y, z$ , then the derivative of this relation with respect to time ( $t$ ) gives the relation between the related rates of these variables :  $\frac{dX}{dt}, \frac{dy}{dt}, \frac{dz}{dt}$

**Remarks**

- ⊙ Let  $X_0$  be the initial value of variable  $X_{(t=0)}$ , and is the rate of change of  $X$  with respect to time is constant (**i.e.**  $\frac{dX}{dt} = \text{constant value}$ ), then after time ( $t$ ) the magnitude of the variable  $X$  is given by :  $X = X_0 + \frac{dX}{dt} \times t$
- ⊙ The distance between any two points :  $(X_1, y_1), (X_2, y_2)$  is  $\sqrt{(X_2 - X_1)^2 + (y_2 - y_1)^2}$
- ⊙ The volume of the bounded part between two concentric spheres, their radii lengths are  $r_1, r_2 = \frac{4}{3} \pi (r_2^3 - r_1^3)$
- ⊙ If  $k = X y z$  (Three variables), then  $\frac{dk}{dt} = \frac{dX}{dt} \times y z + \frac{dy}{dt} \times X z + \frac{dz}{dt} \times X y$
- ⊙ The distance between the point  $(X_1, y_1)$  and the straight line

$$aX + by + c = 0 \text{ is } \frac{|aX_1 + by_1 + c|}{\sqrt{a^2 + b^2}}$$

- ⊙ If the measure of angle  $X$  in radian, then :









$$1. \frac{d(\sin X)}{dt} = \cos X \cdot \frac{dX}{dt}$$

$$2. \frac{d(\cos X)}{dt} = (-\sin X) \times \frac{dX}{dt}$$

$$3. \frac{d(\tan X)}{dt} = \sec^2 X \cdot \frac{dX}{dt}$$



## Areas and perimeters of some geometrical figures

<b>Rectangle</b>		Perimeter = $2(x + y)$ Area = $x \times y$
<b>Square</b>		Perimeter = $4l$ Area = $l^2 = \frac{1}{2}d^2$
<b>Triangle</b>		Perimeter = sum of side lengths Area = $\frac{1}{2} \times b \times h$ $= \frac{1}{2}$ the product of any two sides $\times$ sine of included angle
<b>Parallelogram</b>		Perimeter = $2(b_1 + b_2)$ Area = $b_1 \times h_1 = b_2 \times h_2$
<b>Rhombus</b>		Perimeter = $4l$ Area = $l \times h$ $= \frac{1}{2}d_1 \times d_2$
<b>Trapezium</b>		Perimeter = sum of its side lengths Area = $\frac{1}{2}(b_1 + b_2) \times h$
<b>Circle</b>		Perimeter = $2\pi r$ Area = $\pi r^2$
<b>Sector</b>		Perimeter = $2r + l$ Area = $\frac{1}{2}lr = \frac{1}{2}\theta^{\text{rad}}r^2$ where $\theta^{\text{rad}} = \frac{l}{r}$ , $\frac{x^\circ}{180^\circ} = \frac{\theta^{\text{rad}}}{\pi}$





## Summary

Window in form of rectangle and equilateral triangle on its top		$\text{Perimeter} = 3x + 2y$ $\text{Area} = xy + \frac{1}{2} x^2 \sin 60^\circ$ $= xy + \frac{\sqrt{3}}{4} x^2$
Window in form of rectangle and semi-circle on its top		$r = \frac{1}{2} x$ $\text{Perimeter} = x + 2y + \pi r$ $= x + 2y + \frac{1}{2} \pi x$ $\text{Area} = xy + \frac{1}{2} \pi r^2$ $= xy + \frac{1}{8} \pi x^2$
Regular polygon	Where n is the number of sides and x is the side length.	$\text{Perimeter} = nx$ $\text{Area} = \frac{1}{4} n x^2 \cot \frac{\pi}{n}$

## Remember volume and lateral area and total surface area of some solids

Solid		Lateral area	Total area	Volume
Cube		$4l^2$	$6l^2$	$l^3$
Cuboid		$2(x+y) \times z$	$2(xy + yz + zx)$	$x \times y \times z$
Right circular cylinder		$2\pi r h$	$2\pi r h + 2\pi r^2$ $= 2\pi r(h+r)$	$\pi r^2 h$
Sphere		-----	$4\pi r^2$	$\frac{4}{3} \pi r^3$
Right cone		$\pi r l$	$\pi r l + \pi r^2$	$\frac{1}{3} \pi r^2 h$



<b>Prism</b>		Base perimeter $\times$ height	Lateral area + sum of areas of its two bases	Area of its base $\times$ height
<b>Regular pyramid</b>		$\frac{1}{2}$ Base perimeter $\times$ slant height	Lateral area + area of its base	$\frac{1}{3}$ base area $\times$ height

### The critical points

If the function  $f$  continuous in the interval  $]a, b[$ , then it has a critical point  $(c, f(c))$

where  $c \in ]a, b[$  if  $\boxed{\hat{f}(c) = 0}$  or  $\boxed{\hat{f}(c) \text{ is not exist}}$

The critical point at  $x = a$  must belong to the domain of the function

**i.e.**  $f(a)$  is defined

### Increasing and decreasing of the function

**1.** The function is increasing on an interval if the slope of the tangent to its curve at any point on it in this interval is positive.

**i.e.** If  $\hat{f}(x) > 0$  for all the values of  $x \in ]a, b[$ , then  $f$  is increasing on this interval.

**2.** The function is decreasing on an interval if the slope of the tangent to its curve at any point on it in this interval is negative.

**i.e.** If  $\hat{f}(x) < 0$  for all the values of  $x \in ]a, b[$ , then  $f$  is decreasing on this interval.

**So we use the first derivative in steps of investigation of increasing and decreasing functions as follow :**

**1.** Determine the domain of the function.

**2.** Find  $\hat{f}(x)$

**3.** Find the critical point. (the points at which  $\hat{f}(x) = 0$  or  $\hat{f}(x)$  is not exist)

**4.** Determine the intervals of the domain by which these points divide the domain.

**5.** Determine the sign of  $\hat{f}(x)$  in each of these intervals and so the increasing intervals where  $[\hat{f}(x) > 0]$  and the decreasing intervals where  $[\hat{f}(x) < 0]$



## Local maxima and minima of function

### ★ Using the first derivative to identify the local maxima and minima

If  $(c, f(c))$  is a critical point of the function  $f$  which is continuous at  $c$  and there is an open interval around  $c$  where :

1.  $f'(x) > 0$  at  $x < c$  ,  $f'(x) < 0$  at  $x > c$  , then  $f(c)$  is a local maximum value.
2.  $f'(x) < 0$  at  $x < c$  ,  $f'(x) > 0$  at  $x > c$  , then  $f(c)$  is a local minimum value.
3. If the sign of  $f'(x)$  on both sides of  $c$  does not change , then the function has no local maximum or local minimum value at  $c$

### ★ Using the second derivative :

If  $f$  is differentiable function twice on an open interval contains  $c$  where  $f'(c) = 0$  and

1.  $f''(c) < 0$  , then  $f(c)$  is local maximum value.
2.  $f''(c) > 0$  , then  $f(c)$  is local minimum value.
3.  $f''(c) = 0$  , then the 2<sup>nd</sup> derivative test failed to determine the kind of the point  $(c, f(c))$  if it is local maximum or local minimum value , in this case we use the first derivative test.

## Remarks

1. The local maximum points and local minimum points are critical points but the converse is not always true.
2. If the function  $f$  is only increasing (or decreasing) on an interval , so the function has no local maximum or local minimum in this interval.
3. The critical point at which the first derivative = 0  
**i.e.** The tangent is horizontal at this point sometimes is called stationary point.
4. The polynomial function of  $n^{\text{th}}$  degree has at most  $(n - 1)$  local maximum or minimum values.

## Steps to study the existence of the local maximum and local minimum values of continuous function not including the constant function :

1. Determine the domain of the function.
2. Calculate  $f'(x)$
3. Find out the critical points (**i.e.** The points at which  $f'(x) = 0$  or not exist) , let the  $x$ -coordinate of one of them be  $x_1$
4. Determine the type of the critical point if it is maximum or minimum by one of the next two methods.



## Applications of maxima and minima

To solve these questions, express the variable wanted to find its maximum or minimum value as a function in one variable, using the givens in the problem, then find the maximum or minimum of this function as explained before.

### Remarks

- ⊙ To find the greatest volume ( $v$ ), put  $\frac{dv}{dx} = 0$ , and make sure that  $\frac{d^2v}{dx^2} < 0$
- ⊙ To find the smallest cost ( $c$ ) put  $\frac{dc}{dx} = 0$ , and make sure that  $\frac{d^2c}{dx^2} > 0$  and so on.

## Absolute maxima and minima (Absolute extrema) on a closed interval

### Studying the absolute maximum and minimum values on a closed interval $[a, b]$

If  $f$  is a continuous function on the interval  $[a, b]$ :

1. Determine the critical points at which  $f'(x) = 0$  or not exist and belongs to the interval  $[a, b]$
2. Find the values of the function at the critical points and the endpoints  $f(a)$  and  $f(b)$
3. Compare among the previous values, then the greatest value is the absolute maximum value on  $[a, b]$  and the smallest value is the absolute minimum value on  $[a, b]$

### Remark

If the function  $f$  is defined on the interval  $[a, b]$  and if:

1.  $f'(x) > 0$  **i.e.** The function is increasing on the same interval, then:

- \* The absolute minimum value =  $f(a)$
- \* The absolute maximum value =  $f(b)$

2.  $f'(x) < 0$  **i.e.** The function is decreasing on the same interval, then:


- \* The absolute minimum value =  $f(b)$
- \* The absolute maximum value =  $f(a)$




## Convexity of curves and inflection points

★ A continuous part of a curve is said to be :

⊙ Convex upwards "concave downwards" If the curve lies below all its tangents.



⊙ Convex downwards "concave upwards" If the curve lies above all its tangents.



1. If  $f$  is a differentiable function on the interval  $]a, b[$ , then the curve of the function  $f$

- (a) Convex downwards If  $\hat{f}'$  is increasing on  $]a, b[$
- (b) Convex upwards If  $\hat{f}'$  is decreasing on  $]a, b[$

2. Let  $f$  be a differentiable function twice on the interval  $]a, b[$

- (a) If  $\hat{f}''(x) > 0$  for all the values of  $x \in ]a, b[$ , then the curve of  $f$  is convex downwards on the interval  $]a, b[$
- (b) If  $\hat{f}''(x) < 0$  for all the values of  $x \in ]a, b[$ , then the curve of  $f$  is convex upwards on the interval  $]a, b[$

★ The point of inflection :

The point  $C(c, f(c))$  is an inflection point of a curve of a function  $f$  if the following is satisfied.

1. The curve of the function  $f$  is continuous at  $C$

2. We can draw one tangent to the curve of the function at  $C$  Where :

- (a)  $\hat{f}(c) \in \mathbb{R}$  **i.e.** the tangent is inclined or horizontal.
- (b)  $\hat{f}(c) = \pm \infty$  (denominator of  $\hat{f}$  at the point  $C = \text{zero}$ ) **i.e.** the tangent is vertical.

3. The sign of  $\hat{f}''(x)$  changes before and after the point  $C$  **i.e.**  $[\hat{f}''(c) = 0 \text{ or not exist}]$

★ Steps to study convexity intervals and inflection points :

1. Find  $\hat{f}''(x)$ , then find the values of  $x$  which make  $\hat{f}'' = 0$  or not exist.
2. Determine the sign of  $\hat{f}''(x)$  to determine the intervals over which the function is convex upwards where  $[\hat{f}''(x) < 0]$  and the intervals over which the function is convex downwards where  $[\hat{f}''(x) > 0]$
3. Determine the inflection points from the obtained points at which the sign of  $\hat{f}''(x)$  changes around it. unless  $\hat{f}''(x)$  changes around any of these points, then it is not be considered an inflection point.



### Remarks

1. The inflection point at  $x = a$  must belong to the domain of the function  
i.e.  $f(a)$  is defined
2. The inflection points are the points which separated between convex upwards and convex downwards regions.
3. The tangent at the inflection point intersects the curve of the function.
4. The critical point of the function  $f$  is the point at which  $f'(x) = 0$  or not exist and if the sign of  $f'(x)$  changes around this point, then it is a local maximum or minimum.

Also we find that :

The critical point of the function  $f$  is the point at which  $f'(x) = 0$  or not exist and if the sign of  $f'(x)$  changes around this point, then it is an inflection point of the function  $f$

5. The inflection points of a function  $f$  which is differentiable twice is a local maximum or minimum of the function  $f''$
6. If the point  $(c, h) \in$  the curve of a function which is differentiable twice and
  - (a)  $(c, h)$  is an inflection point, then  $f''(c) = 0$ ,  $f(c) = h$
  - (b)  $(c, h)$  is a local maximum or a local minimum or critical point, then  $f'(c) = 0$ ,  $f(c) = h$

### Curve sketching of polynomial functions

Steps of drawing a curve of function  $f$  (where  $f$  is a polynomial of 3<sup>rd</sup> degree or less) :

1. Determine the domain of the function, then determine the symmetric of the function  $f$  if exist where :
  - (a)  $f(-x) = f(x)$  for every  $x \in$  the domain  
 $\therefore$  The function is even and so the curve is symmetric about y-axis.
  - (b)  $f(-x) = -f(x)$  for every  $x \in$  the domain  
 $\therefore$  The function is odd and so the function is symmetric about origin point.
2. Find out  $f'(x)$ ,  $f''(x)$
3. Use  $f'(x)$  to determine :
  - (a) The increasing interval where  $[f'(x) > 0]$ , decreasing interval where  $[f'(x) < 0]$
  - (b) The points of local maximum and local minimum (if exist) where  $f'(x) = 0$   
(notice that the function is differentiable) and the sign of  $f'(x)$  changes before and after this point.







4. Use  $f''(x)$  to determine :

- (a) The intervals where the curve convex upwards where  $[f''(x) < 0]$  , the intervals where the curve convex downwards where  $[f''(x) > 0]$
- (b) The inflection point (if exist) where  $f''(x) = 0$  (notice that the function is differentiable twice) and the sign of  $f''(x)$  changes before and after the point.

5. Determine some assisting points in sketching as :

- (a) The point / points of intersection with  $x$ -axis
- (b) The point / points of intersection with  $y$ -axis
- (c) Some extra points by substitution by some values of  $x$  and get the corresponding values of  $f(x)$

6. Arrange the points we get in a table and represent them graphically then join these points taking in consideration the following :

Sign of $f'(x)$ , $f''(x)$	Properties of the curve of function $f$	Shape of the curve
$f'(x) > 0$ , $f''(x) > 0$	Increasing , convex downwards	
$f'(x) > 0$ , $f''(x) < 0$	Increasing , convex upwards	
$f'(x) < 0$ , $f''(x) > 0$	Decreasing , convex downwards	
$f'(x) < 0$ , $f''(x) < 0$	Decreasing , convex upwards	

### Indefinite integration

#### • Some fundamental integrations (standard)

1.  $\int k \, dx = kx + c$  (where  $k$  is constant)

2.  $\int x^n \, dx = \frac{x^{n+1}}{n+1} + c$  (where  $n \neq -1$ )

3.  $\int (ax + b)^n \, dx = \frac{(ax + b)^{n+1}}{a(n+1)} + c$  (where  $n \neq -1$ )

4.  $\int e^x \, dx = e^x + c$

5.  $\int e^{kx+b} \, dx = \frac{e^{kx+b}}{k} + c$



$$6. \int a^x dx = \frac{a^x}{\ln a} + c$$

$$7. \int a^{kx+b} dx = \frac{e^{kx+b}}{k \ln a} + c$$

$$8. \int \frac{1}{x} dx = \ln |x| + c \quad (\text{where } x \neq 0)$$

• **Important rules of integration :**

$$1. \int (f(x))^n f'(x) dx = \frac{(f(x))^{n+1}}{n+1} + c$$

$$2. \int e^{f(x)} f'(x) dx = e^{f(x)} + c$$

$$3. \int a^{f(x)} f'(x) dx = \frac{a^{f(x)}}{\ln a} + c$$

$$4. \int \frac{f'(x)}{f(x)} dx = \ln |f(x)| + c$$

$$5. \int \frac{f'(x)}{\sqrt{f(x)}} dx = 2\sqrt{f(x)} + c$$

• **Some properties of the indefinite integration :**

$$1. \int a f(x) dx = a \int f(x) dx \quad (\text{where } a \text{ is a constant } \neq 0)$$

$$2. \int [f(x) \pm g(x)] dx = \int f(x) dx \pm \int g(x) dx$$

$$3. \frac{d}{dx} \int f(x) dx = f(x)$$

$$4. \int \frac{d}{dx} [f(x)] dx = f(x) + c$$

$$5. \int f(x) dx - \int f(x) dx = \text{constant (not necessary } = 0)$$

**Results**

$$1. \int \sin(ax+b) dx = -\frac{1}{a} \cos(ax+b) + c$$

$$2. \int \cos(ax+b) dx = \frac{1}{a} \sin(ax+b) + c$$

$$3. \int \sec^2(ax+b) dx = \frac{1}{a} \tan(ax+b) + c$$

$$4. \int \csc^2(ax+b) dx = -\frac{1}{a} \cot(ax+b) + c$$

$$5. \int \sec(ax+b) \tan(ax+b) dx = \frac{1}{a} \sec(ax+b) + c$$

$$6. \int \csc(ax+b) \cot(ax+b) dx = -\frac{1}{a} \csc(ax+b) + c$$

**Remember**

$$1. \int \tan x \, dx = \int \frac{\sin x}{\cos x} \, dx = - \int \frac{-\sin x}{\cos x} \, dx$$

$$= -\ln |\cos x| + c = \ln |\sec x| + c$$

**Notice that**

The numerator is the derivative of the denominator.

$$2. \int \cot x \, dx = \int \frac{\cos x}{\sin x} \, dx$$

$$= \ln |\sin x| + c = -\ln |\csc x| + c$$

**Notice that**

The numerator is the derivative of the denominator.

$$3. \int \sec x \, dx = \int \frac{\sec x (\sec x + \tan x)}{(\sec x + \tan x)} \, dx$$

(by multiplying numerator and denominator by  $(\sec x + \tan x)$ )

$$= \int \frac{\sec^2 x + \sec x \cdot \tan x}{\sec x + \tan x} \, dx$$

$$= \ln |\sec x + \tan x| + c$$

**Notice that**

The numerator is the derivative of the denominator.

$$4. \int \csc x \, dx = \int \frac{\csc x (\csc x + \cot x)}{(\csc x + \cot x)} \, dx$$

(by multiplying numerator and denominator by  $(\csc x + \cot x)$ )

$$= - \int \frac{-\csc^2 x - \csc x \cot x}{\csc x + \cot x} \, dx$$

$$= -\ln |\csc x + \cot x| + c$$

**Notice that**

The numerator is the derivative of the denominator.

**Generally**

$$1. \int \sin(f(x)) \times f'(x) \, dx = -\cos(f(x)) + c$$

$$2. \int \cos(f(x)) \times f'(x) \, dx = \sin(f(x)) + c$$

$$3. \int \sec^2(f(x)) \times f'(x) \, dx = \tan(f(x)) + c$$

$$4. \int \csc^2(f(x)) \times f'(x) \, dx = -\cot(f(x)) + c$$

$$5. \int \sec(f(x)) \times \tan(f(x)) \times f'(x) \, dx = \sec(f(x)) + c$$

$$6. \int \csc(f(x)) \times \cot(f(x)) \times f'(x) \, dx = -\csc(f(x)) + c$$



## ★ Methods of integration :

### 1. Integration by substitution :

⊙ If the given integration is given in the form  $\int f(g(x)) \cdot g'(x) \cdot dx$

use the substitution  $g(x) = z$


⊙ If the given integration contains the  $n^{\text{th}}$  root of a function as  $\sqrt[n]{g(x)}$ , use the

substitution  $g(x) = z^n$  or  $g(x) = z$

⊙ In some questions you can use an appropriate substitution in order to simplify the integration and rewrite it in standard form.

### 2. Integration by parts :

$$\int z \, dy = yz - \int y \, dz$$



## The definite integral

If the function  $f$  is continuous on the interval  $[a, b]$ , and  $F$  is any anti derivative to the function  $f$  on the same interval, then :  $\int_a^b f(x) \, dx = F(b) - F(a)$

## ★ Properties of definite integral :

1. If  $f$  is a continuous on  $[a, b]$ ,  $c \in ]a, b[$ , then :

$$1. \int_b^a f(x) \, dx = - \int_a^b f(x) \, dx$$

$$2. \int_a^a f(x) \, dx = \text{zero}$$

$$3. \int_a^b f(x) \, dx = \int_a^c f(x) \, dx + \int_c^b f(x) \, dx$$

2. If  $f$  is a continuous odd function on the interval  $[-a, a]$ , then  $\int_{-a}^a f(x) \, dx = \text{zero}$

3. If  $f$  is a continuous even function on the interval  $[-a, a]$

$$\text{, then } \int_{-a}^a f(x) \, dx = 2 \int_0^a f(x) \, dx$$

4. If  $f, g$  are two continuous function on the interval  $[a, b]$

$$1. \int_a^b [f(x) \pm g(x)] \, dx = \int_a^b f(x) \, dx \pm \int_a^b g(x) \, dx$$

$$2. \int_a^b k f(x) \, dx = k \int_a^b f(x) \, dx \text{ (where } k \in \mathbb{R})$$




## Definite integral and areas in the plane

**First** The area of a region bounded by the curve of the function  $f$  and  $x$ -axis in the interval  $[a, b]$

1.  $f(x) \geq 0$


i.e. "The region above  $x$ -axis"



, then  $A = \int_a^b f(x) dx$

2.  $f(x) \leq 0$

i.e. "The region below  $x$ -axis"



, then  $A = - \int_a^b f(x) dx = \left| \int_a^b f(x) dx \right|$


- ⊙ If the curve of the function  $f$  intersects the  $x$ -axis at  $x = c$  and  $x = d$  where  $c$  and  $d$  belong to  $[a, b]$  as in the opposite figure :

We find that :  $f(x) \geq 0$  for all  $x \in [c, d]$

,  $f(x) \leq 0$  for all  $x \in [a, c]$  and  $x \in [d, b]$

$\therefore$  The shaded area ( $A$ ) =  $A_1 + A_2 + A_3$

i.e.  $A = \left| \int_a^c f(x) dx \right| + \int_c^d f(x) dx + \left| \int_d^b f(x) dx \right|$



**Notice that**

The absolute value to the two areas  $A_1, A_3$  because they are below  $x$ -axis.



### Remarks

1. It is favorable to graph the curve of the function to identify the region above or below  $x$ -axis.
2. Sometimes it is difficult to sketch some curves then , its favorable to find zeroes of the function (even the limits of the integral are given) which divides the domain of the function  $[a, b]$  if exist into intervals and determine the sign of the function in each part and so you know if the region is above or below  $x$ -axis.
3. The value of the integration , could be positive or negative but the area is always positive.
4. In general , the area of the included region between any continuous function :  
 $y = f(x)$  and the  $x$ -axis and the two straight lines  $x = a$  ,  $x = b$

is  $A = \int_a^b |f(x)| dx$

### Second The area of plane region bounded by two curves

If  $f, g$  are two continuous functions on the interval  $[a, b]$  and  $f(x) \geq g(x)$  for every  $x \in [a, b]$  , then the area bounded by the region between the two curves  $y_1 = f(x)$  ,  $y_2 = g(x)$  and the two straight lines  $x = a$  ,  $x = b$  is given by the relation

$$A = \int_a^b [f(x) - g(x)] dx$$

and that's because the area between

the two curves  $y_1 = f(x)$  ,  $y_2 = g(x)$


= [The area under  $f(x)$  and above  $x$ -axis]

- [The area under  $g(x)$  and above  $x$ -axis]

$$= \int_a^b y_1 dx - \int_a^b y_2 dx$$

$$= \int_a^b f(x) dx - \int_a^b g(x) dx$$

$$= \int_a^b [f(x) - g(x)] dx$$



**Remarks**

1. To identify the higher function  $y_1 \geq y_2$  for every  $x \in [a, b]$  by using the graph or by choosing an arbitrary value of  $x \in [a, b]$  and substitution in the equations of the two curves, or by using the absolute value as follow :

The area (A) =  $\left| \int_a^b (\text{any of the two functions} - \text{the other function}) dx \right|$

2. When a region included between two curves, then the terms of integral with respect to  $x$  are the  $x$ -coordinates of their points of intersection and it will be found by solving the two equations algebraically.


3. If the two curves intersect at one point  $c \in ]a, b[$

and  $f(x) \geq g(x)$  for every  $x \in [a, c]$

and  $g(x) \geq f(x)$  for every  $x \in [c, b]$

, then  $A = A_1 + A_2$


$$= \int_a^c [f(x) - g(x)] dx + \int_c^b [g(x) - f(x)] dx$$

**Volumes of revolution solids****Revolution Solid**

- \* It is a solid generated by revolving a plane area a complete revolution about a fixed straight line in its plane called «axis of revolution»


**The volume of a solid produced by revolving a region about an axis.**

**The  $x$ -axis**



$$v = \pi \int_a^b y^2 dx = \pi \int [f(x)]^2 dx$$

**The  $y$ -axis**




$$v = \pi \int_c^d x^2 dy = \pi \int_c^d [f(y)]^2 dy$$



- \* Volume of the solid generated by revolving region bounded by two curves

$$v = \pi \int_a^b (y_1^2 - y_2^2) dx$$

$$= \pi \int_a^b [(f(x))^2 - (g(x))^2] dx$$




### Remarks

1. If a region bounded by two intersecting curves  $y_1 = f(x)$ ,  $y_2 = g(x)$  where  $y_1 \geq y_2$  for every  $x \in [a, b]$  revolve a complete revolution about  $x$ -axis, then the  $x$ -coordinates of the two intersecting points of the two curves are the terms of integration  $a, b$ , where  $a < b$  and,

$$v = \pi \int_a^b (y_1^2 - y_2^2) dx$$


**i.e.**  $v = \pi \int_a^b y_1^2 dx - \pi \int_a^b y_2^2 dx$



2. If a region bounded by two intersecting curves  $x_1 = f(y)$ ,  $x_2 = g(y)$  where  $x_1 \geq x_2$  for every  $y \in [c, d]$  revolve a complete revolution about  $y$ -axis, then the  $y$ -coordinates of the two intersecting points of the two curves are the terms of integration  $c, d$  where  $c < d$  and

$$v = \pi \int_c^d (x_1^2 - x_2^2) dy$$


**i.e.**  $v = \pi \int_c^d x_1^2 dy - \pi \int_c^d x_2^2 dy$



# Multiple Choice Question Bank

in

## Differential & Integral calculus







### First

### Questions on limits

Choose the correct answer from the given ones :

1  $\lim_{x \rightarrow \infty} \left(1 + \frac{1}{x}\right)^{2x} = \dots\dots\dots$

- (a) 1                                      (b) 2                                      (c) e                                      (d)  $e^2$

2  $\lim_{x \rightarrow 0} (1 + x)^{\frac{1}{3x}} = \dots\dots\dots$

- (a)  $\frac{1}{3}$                                       (b)  $e^3$                                       (c)  $e^{\frac{1}{3}}$                                       (d)  $\frac{e}{3}$

3  $\lim_{x \rightarrow \infty} \left(1 + \frac{1}{1+x}\right)^x = \dots\dots\dots$

- (a) e                                      (b)  $\frac{1}{1+e}$                                       (c)  $e+1$                                       (d)  $e^{-1}$

4  $\lim_{x \rightarrow \infty} \left(\frac{x+5}{x+3}\right)^x = \dots\dots\dots$

- (a) e                                      (b)  $e^2$                                       (c)  $\frac{1}{e}$                                       (d)  $\frac{2}{e}$

5  $\lim_{x \rightarrow \infty} \left(\frac{x+7}{x+3}\right)^{x+4} = \dots\dots\dots$

- (a)  $e^4$                                       (b)  $e^3$                                       (c)  $e^2$                                       (d)  $e^5$

6  $\lim_{x \rightarrow 0} \frac{e^{4x} - 1}{5x} = \dots\dots\dots$

- (a)  $\frac{4}{5}$                                       (b)  $e^{\frac{4}{5}}$                                       (c)  $e^{\frac{5}{4}}$                                       (d)  $\ln 5$

7  $\lim_{x \rightarrow 0} \frac{e^x - \sin x - 1}{x} = \dots\dots\dots$

- (a) 2                                      (b) 1                                      (c)  $e-1$                                       (d) zero

8  $\lim_{x \rightarrow 0} \frac{5^x - 1}{3^x - 1} = \dots\dots\dots$

- (a)  $\ln \frac{5}{3}$                                       (b)  $\frac{\log 5}{\log 3}$                                       (c)  $\log \frac{5}{3}$                                       (d)  $\frac{5}{3}$



9  $\lim_{x \rightarrow 0} \frac{e^{-x} - 1}{x} = \dots\dots\dots$

(a)  $e$

(b)  $-1$

(c)  $-e$

(d)  $e^{-1}$

10  $\lim_{x \rightarrow 0} \frac{2^x - 1}{3x} = \dots\dots\dots$

(a)  $3 \ln 2$

(b)  $\frac{1}{3} \ln 2$

(c)  $\ln \frac{2}{3}$

(d)  $2 \ln 3$

11  $\lim_{x \rightarrow 0} \frac{e^x - \cos x}{x} = \dots\dots\dots$

(a) zero

(b)  $-1$

(c)  $1$

(d)  $e$

12  $\lim_{x \rightarrow 0} \frac{e^{\tan x} - 1}{\tan 2x} = \dots\dots\dots$

(a)  $e$

(b)  $1$

(c)  $\frac{1}{2}$

(d)  $2$

13  $\lim_{x \rightarrow 0} \sqrt[x]{1+x} = \dots\dots\dots$

(a) zero

(b)  $1$

(c)  $\frac{1}{e}$

(d)  $e$

14  $\lim_{x \rightarrow 0} (e^x - 4) = \dots\dots\dots$

(a)  $-4$

(b)  $-3$

(c)  $e$

(d)  $e^4$

15  $\lim_{x \rightarrow 1} \left( \frac{\ln x}{x-1} \right) = \dots\dots\dots$

(a) zero

(b)  $1$

(c)  $e$

(d)  $e^{-1}$

16  $\lim_{x \rightarrow 0} \frac{1 - e^{2x}}{1 - e^x} = \dots\dots\dots$

(a)  $1$

(b)  $-1$

(c)  $2$

(d)  $-2$

17  $\lim_{x \rightarrow 0} \left( 1 + \frac{x}{a} \right)^{\frac{a}{x}} = \dots\dots\dots$

(a)  $e^{-1}$

(b)  $-e$

(c)  $-e^{-1}$

(d)  $e$

18  $\lim_{x \rightarrow 0} \frac{\ln(1+x^2)}{x^2} = \dots\dots\dots$

(a)  $1$

(b)  $\log_a e$

(c)  $\ln a$

(d)  $2$



19  $\lim_{x \rightarrow 0} \frac{\log_3(1+x)}{\sin 2x} = \dots\dots\dots$

- (a)  $\log_3 e$  (b)  $2 \log_3 e$  (c)  $\log_3 \sqrt{e}$  (d)  $\frac{1}{2}$

20  $\lim_{x \rightarrow 0} \frac{\log(1+2x)}{5^x - 1} = \dots\dots\dots$

- (a)  $2 \ln 10 \ln 5$  (b)  $\frac{2}{5}$  (c)  $\log_5 2e$  (d)  $2 \log e \log_5 e$

21  $\lim_{x \rightarrow 0} \frac{2^x - 1}{\sqrt{x+1} - 1} = \dots\dots\dots$

- (a)  $\ln 2$  (b)  $\ln 4$  (c)  $\log_2 e$  (d)  $\log_4 e$

22  $\lim_{x \rightarrow 0} \frac{e^x - e^{\sin x}}{x - \sin x} = \dots\dots\dots$

- (a)  $e^{\sin x}$  (b)  $e$  (c)  $1$  (d) zero

23  $\lim_{x \rightarrow 6} \frac{e^x - e^6}{x - 6} = \dots\dots\dots$

- (a)  $e$  (b)  $e^6$  (c)  $\ln 6$  (d)  $e^{12}$

24  $\lim_{x \rightarrow \frac{\pi}{2}} (1 + \cos x)^{\frac{\sec x}{4}} = \dots\dots\dots$

- (a)  $4e$  (b)  $e^4$  (c)  $\frac{1}{4}e$  (d)  $e^{\frac{1}{4}}$

25  $\lim_{x \rightarrow 0} (1 + 3 \tan^2 x)^{\cot^2 x} = \dots\dots\dots$

- (a)  $e^3$  (b)  $e^{-3}$  (c)  $3e$  (d)  $-3e$

26  $\lim_{x \rightarrow 0} \left(4 \sin^3 x + \sin \frac{\pi}{2}\right)^{\csc^3 x} = \dots\dots\dots$

- (a)  $e^{-4}$  (b)  $4e$  (c)  $e^2$  (d)  $e^4$

27  $\lim_{x \rightarrow \infty} \left(\frac{x^2 - 7x + 12}{x^2 - 8x + 16}\right)^{2x-4} = \dots\dots\dots$

- (a)  $1$  (b)  $e$  (c)  $e^2$  (d)  $e^4$



# Multiple choice question bank

28  $\lim_{x \rightarrow 0} (1 + 2x + x^2)^{\frac{1}{2x}} = \dots\dots\dots$

(a)  $e^2$

(b)  $\frac{1}{2}e$

(c)  $e$

(d)  $e^{\frac{1}{2}}$

29  $\lim_{x \rightarrow 0} \frac{a^x + b^x + c^x - 3}{x} = \dots\dots\dots$

(a)  $\ln(a b c)$

(b)  $\log a + \log b + \log c$

(c)  $\ln a \cdot \ln b \cdot \ln c$

(d) 1

30  $\lim_{n \rightarrow \infty} n (\ln(n+1) - \ln n) = \dots\dots\dots$

(a) zero

(b)  $e$

(c)  $\frac{1}{e}$

(d) 1

31 If  $\lim_{x \rightarrow a} \left(1 + \frac{1}{x}\right)^{2x} = e^2$ , then  $a$  is  $\dots\dots\dots$

(a) zero

(b)  $e$

(c) 1

(d)  $\infty$

32 If  $\lim_{x \rightarrow 0} (1 + kx)^{\frac{4}{x}} = e^{-12}$ , then  $k = \dots\dots\dots$

(a)  $-\frac{4}{3}$

(b) -3

(c) -12

(d) -48

33 If  $\lim_{x \rightarrow \infty} \left(1 + \frac{a+2}{x}\right)^x = e^4$ , then  $a = \dots\dots\dots$

(a) -2

(b) 2

(c) 3

(d) 4

34 If  $\lim_{x \rightarrow \infty} \left(\frac{x}{x+k}\right)^x = e$ , then  $k = \dots\dots\dots$

(a) 1

(b) -1

(c) -2

(d) 2

35  $\lim_{x \rightarrow 0} \frac{(10)^{\sin x} - 1}{\tan x} = \dots\dots\dots$

(a)  $\ln 10$

(b)  $\log x$

(c) zero

(d) 1

36  $\lim_{x \rightarrow 0} \frac{\ln(1-x)}{x} = \dots\dots\dots$

(a)  $e$

(b)  $-e$

(c) -1

(d) 1

37  $\lim_{x \rightarrow 3} \frac{\ln(x-2)}{x-3} = \dots\dots\dots$

(a)  $e$

(b) 1

(c)  $e^2$

(d) -1



- 38  $\lim_{x \rightarrow 0} \frac{\ln(x^2 + 3x + 1)}{\ln(x^2 + 5x + 1)} = \dots\dots\dots$   
 (a) 1 (b)  $\frac{3}{5}$  (c)  $\frac{5}{3}$  (d)  $\ln \frac{3}{5}$
- 
- 39 If  $\lim_{x \rightarrow 0} \frac{e^{a^2 x} - e^{5ax}}{3x} = 2$ , then  $a = \dots\dots\dots$  where  $a \in \mathbb{R}^+$   
 (a) 2 (b) 3 (c) 5 (d) 6
- 
- 40 If  $\lim_{x \rightarrow 0} \frac{\ln(1 + ax)}{bx} = -1$ , then  $a + b = \dots\dots\dots$   
 (a) -1 (b) zero (c) 1 (d) 2
- 
- 41 (Trial 2021) If  $\lim_{x \rightarrow 0} \frac{\ln(x+1)^{\sqrt{k}}}{x} = 4$ , then  $k = \dots\dots\dots$   
 (a) 2 (b) 4 (c) 8 (d) 16
- 
- 42 (1<sup>st</sup> session 2021) If  $\lim_{x \rightarrow 0} \frac{e^{x+a} - e^a}{x} = \frac{1}{e}$ , then the value of  $a = \dots\dots\dots$   
 (a) zero (b) 1 (c)  $e$  (d) -1
- 
- 43 If  $\lim_{x \rightarrow k} (2 + x)^{\frac{1}{x+1}} = e$ , then  $k = \dots\dots\dots$   
 (a) zero (b) 1 (c) -1 (d)  $e$
- 
- 44 (2<sup>nd</sup> session 2021) If  $\lim_{x \rightarrow 0} \frac{\ln(1+x)^{4a}}{x} = a^2 + 4$ , then the value of constant  $a = \dots\dots\dots$   
 (a) -2 (b) -4 (c) 2 (d) 4
- 
- 45 If  $\lim_{x \rightarrow 0} \frac{e^{ax} - 1}{\ln(1 + 3x)} = 2$ , then  $a = \dots\dots\dots$   
 (a) 1 (b) 2 (c) 3 (d) 6
- 
- 46  $\lim_{x \rightarrow 0} \frac{xe^x - x}{1 - \cos 2x} = \dots\dots\dots$   
 (a) 1 (b)  $\frac{1}{2}$  (c)  $e^2$  (d)  $e^{\frac{1}{2}}$
- 
- 47  $\lim_{x \rightarrow 0} \frac{2^{2x} - 2^{3x}}{x} = \dots\dots\dots$   
 (a)  $\ln 2$  (b)  $\ln \frac{1}{2}$  (c)  $\ln 4$  (d)  $\ln \frac{1}{4}$
- 
- 48  $\lim_{x \rightarrow 0} \frac{(10)^x - 2^x - 5^x + 1}{x \sin x} = \dots\dots\dots$   
 (a)  $\ln 10$  (b)  $\ln \frac{5}{2}$  (c)  $\ln 5 \times \ln 2$  (d)  $\log_2 5$



49 If  $\lim_{x \rightarrow 0} \frac{2^x + 3^x + 4^x - 3}{x} = \ln a + \ln b$  then  $a b = \dots\dots\dots$

(a) 12

(b) 16

(c) 24

(d) 44

50 If  $f(x) = e^{\tan x}$ , then  $\lim_{x \rightarrow \frac{\pi}{4}} \frac{f(x) - f(\frac{\pi}{4})}{x - \frac{\pi}{4}} = \dots\dots\dots$

(a) e

(b)  $2e$ (c)  $e^2$ (d)  $2e^2$ 

51  $\lim_{h \rightarrow 0} \frac{\cot(x+h) - \cot(x)}{h} = \dots\dots\dots$

(a)  $-\csc^2 x$ (b)  $\sec^2 x$ (c)  $\cot^2 x$ (d)  $\tan^2 x$ 

52  $\lim_{h \rightarrow 0} \frac{\sec(\frac{\pi}{4} + h) - \sec(\frac{\pi}{4})}{h} = \dots\dots\dots$

(a)  $\sqrt{2}$ 

(b) zero

(c)  $\frac{1}{\sqrt{2}}$ 

(d) undefined.

53 If  $f(x)$  is a rule of a polynomial function, then  $\lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \dots\dots\dots$

(a)  $\dot{f}(x)$ (b)  $\dot{f}(h)$ (c)  $\ddot{f}(x)$ (d)  $\ddot{f}(h)$ 

54 If  $\dot{f}(x) = \cos^3 x$  and  $f(0) = 0$ , then  $\lim_{x \rightarrow 0} \frac{f(x)}{x} = \dots\dots\dots$

(a) -1

(b) zero

(c) 1

(d) does not exist

55 If  $\lim_{x \rightarrow 2} f(x) = \text{zero}$ , then  $\lim_{x \rightarrow 2} (1 + f(x))^{\frac{1}{f(x)}} = \dots\dots\dots$

(a) zero

(b) b

(c) e

(d)  $e^{f(2)}$ 

56 If  $\lim_{x \rightarrow \infty} \left(\frac{x+a}{x+b}\right)^x = e^c$ , then  $c = \dots\dots\dots$

(a)  $a+b$ (b)  $a-b$ (c)  $b-a$ (d)  $\frac{a+b}{2}$ 

57 If  $e^x = \sum_{n=0}^{\infty} \frac{x^n}{n!}$ , then  $\lim_{x \rightarrow 0} \frac{e^x - 1 - x - \frac{x^2}{2}}{x^3} = \dots\dots\dots$

(a)  $\frac{1}{2}$ (b)  $\frac{1}{3}$ (c)  $\frac{1}{6}$ (d)  $e^3 - 1$



## Second

## Questions on derivatives - Implicit and parametric differentiation - Higher derivatives

Choose the correct answer from the given ones :

- 1 If  $f(x) = \cot(5x - \pi)$ , then  $f'\left(\frac{\pi}{4}\right) = \dots\dots\dots$   
 (a)  $5\sqrt{2}$  (b)  $-5\sqrt{2}$  (c) 10 (d) -10
- 
- 2 If  $y = \csc 2x$ , then  $\frac{dy}{dx} = \dots\dots\dots$  at  $x = \frac{\pi}{6}$   
 (a)  $\frac{3}{4}$  (b)  $-\frac{4}{3}$  (c)  $\frac{1}{2}$  (d)  $\sqrt{3}$
- 
- 3 If  $y = \sec\left(\frac{\pi}{4}x\right)$ , then  $\frac{dy}{dx} = \dots\dots\dots$   
 (a)  $\frac{\pi}{4} \sec\left(\frac{\pi}{4}x\right) \tan\left(\frac{\pi}{4}x\right)$  (b)  $\frac{\pi}{4}x$   
 (c)  $\frac{\pi}{4}$  (d)  $\sqrt{2}$
- 
- 4 If  $y = \left(\sec \frac{\pi}{4}\right)x$ , then  $\frac{dy}{dx} = \dots\dots\dots$   
 (a)  $\frac{\pi}{4} \sec\left(\frac{\pi}{4}x\right) \tan\left(\frac{\pi}{4}x\right)$  (b)  $\frac{\pi}{4}x$   
 (c)  $\frac{\pi}{4}$  (d)  $\sqrt{2}$
- 
- 5 If  $y = \csc \sqrt{x}$ , then  $\frac{dy}{dx} = \dots\dots\dots$   
 (a)  $-\frac{1}{2\sqrt{x}} \csc \sqrt{x} \cot \sqrt{x}$  (b)  $-\csc \sqrt{x} \times \cot \sqrt{x}$   
 (c)  $-\csc^2 \sqrt{x}$  (d)  $-\frac{1}{2\sqrt{2}} \csc^2 \sqrt{x}$
- 
- 6 If  $y = \cot(x^2 + 3)$ , then  $\frac{dy}{dx} = \dots\dots\dots$   
 (a)  $-2x \csc^2(x^2 + 3)$  (b)  $2x \csc^2(x^3 + 3)$   
 (c)  $-2x \cot(x^2 + 3) \csc(x^2 + 3)$  (d)  $\csc^2(x^2 + 3)$
- 
- 7 If  $f(x) = (5 - 2 \cot x)^3$ , then  $f'\left(\frac{\pi}{4}\right) = \dots\dots\dots$   
 (a) -108 (b) 27 (c) 54 (d) 108



- 8 If  $f(x) = \frac{x-4}{2\sqrt{x}}$ , then  $f'(0) = \dots\dots\dots$   
 (a) zero (b) 1 (c) not exist (d) -1
- 
- 9 If  $g(x) = |x|$ , then  $g'(-5) = \dots\dots\dots$   
 (a) 5 (b) -5 (c) 1 (d) -1
- 
- 10 If  $y = \sqrt{f(x)}$ , and  $f'(2) = 4$ ,  $f(2) = 9$ , then  $\frac{dy}{dx}$  when  $x = 2$  equals  $\dots\dots\dots$   
 (a)  $\frac{4}{3}$  (b)  $\frac{2}{9}$  (c)  $\frac{1}{6}$  (d)  $\frac{2}{3}$
- 
- 11 If  $x^3 y^2 = 1$ , then  $\left[\frac{dy}{dx}\right]_{y=1} = \dots\dots\dots$   
 (a)  $-\frac{2}{3}$  (b)  $-\frac{3}{2}$  (c)  $\frac{2}{3}$  (d)  $\frac{3}{2}$
- 
- 12 If  $f(5x) = x^2 + x$ , then  $f'(2) = \dots\dots\dots$   
 (a) 5 (b) 1 (c)  $\frac{3}{25}$  (d)  $\frac{9}{25}$
- 
- 13 If  $xy = 1$ , then  $\frac{dy}{dx}$  equals each of the following except  $\dots\dots\dots$   
 (a)  $x^{-2}$  (b)  $\frac{-1}{x^2}$  (c)  $\frac{-y}{x}$  (d)  $-y^2$
- 
- 14 If  $\frac{x}{y} + \frac{y}{x} = a$  where  $a$  is constant, then  $\frac{dy}{dx} = \dots\dots\dots$   
 (a)  $\frac{x}{y}$  (b)  $\frac{y}{x}$  (c)  $\frac{ax}{y}$  (d)  $\frac{ay}{x}$
- 
- 15  $\frac{d}{dx} \left(2 \cot \frac{\pi}{4}\right) = \dots\dots\dots$   
 (a)  $-2 \csc^2 \frac{\pi}{4}$  (b)  $-2 \cot \frac{\pi}{4} \csc \frac{\pi}{4}$  (c) 2 (d) zero
- 
- 16 If  $\frac{2}{\sqrt{x} + \sqrt{y}} = 9$ , then  $\left(\frac{dy}{dx}\right)^2 = \dots\dots\dots$   
 (a)  $\frac{x}{y}$  (b)  $\frac{y}{x}$  (c)  $\frac{2y}{x}$  (d)  $\frac{1}{2\sqrt{xy}}$
- 
- 17 If  $y = 8x^3$ , then  $dy = \dots\dots\dots$   
 (a)  $24x^2 dx$  (b)  $24x^2 + c$  (c)  $2x^4 + c$  (d)  $24x^2$



18 If  $m = 4\pi r^2$ , which of the following is equal to differential  $m$ ?

- (a)  $4\pi$  (b)  $2r dr$  (c)  $8\pi r dr$  (d)  $8\pi dr$

19  $\frac{d}{dx}(2\cos^2 x - 1) = \dots\dots\dots$

- (a)  $\sin 2x$  (b)  $\cos 2x$  (c)  $-2\sin 2x$  (d)  $-2\cos 2x$

20 If  $f(x) = \sin x \sec 2x$ , then  $\dot{f}(0) = \dots\dots\dots$

- (a)  $-1$  (b) zero (c)  $1$  (d)  $2$

21  $\frac{d}{dx}\left[x^2 + \frac{d}{dx}(x + \sec x)\right] = \dots\dots\dots$

- (a)  $2x + 1 + \sec x \tan x$  (b)  $2x + 2\sec^3 x - \sec x$   
(c)  $2 + \sec^3 x - \sec x$  (d)  $2x + \sec^2 x \tan^2 x$

22 If  $f(x) = 8x \sin x \cos x \cos 2x$ , then  $\dot{f}\left(\frac{\pi}{8}\right) = \dots\dots\dots$

- (a)  $-2$  (b) zero (c)  $1$  (d)  $2$

23  $\frac{d}{dx}(\cos x \csc x) = \dots\dots\dots$

- (a) zero (b)  $\csc^2 x$  (c)  $\sec^2 x$  (d)  $-\csc^2 x$

24 If  $y = \sin x \sec\left(\frac{\pi}{2} - x\right)$  where  $x$  is an acute angle, then  $\frac{dy}{dx} = \dots\dots\dots$

- (a) zero (b)  $1$   
(c)  $\cos x \csc x + \sin x \sec x$  (d)  $\sin x \cos x + \sec x \cot x$

25 If  $y + \cot x = 0$ , then  $\frac{dy}{dx} = \dots\dots\dots$

- (a)  $1 + y$  (b)  $1 + y^2$  (c)  $y^2 - 1$  (d)  $y^2$

26 (2<sup>nd</sup> session 2021) If  $y = \cot 5x$ , then  $\frac{dy}{dx} + 5y^2 = \dots\dots\dots$

- (a)  $5$  (b)  $-5y$  (c)  $5y$  (d)  $-5$

27 If  $y = \csc x$ , then  $\frac{dy}{dx} = \dots\dots\dots$  where  $x$  is acute angle.

- (a)  $-y\sqrt{y^2 - 1}$  (b)  $y\sqrt{y^2 + 1}$  (c)  $\sqrt{y^2 - 1}$  (d)  $\sqrt{y^2 + 1}$



28 If  $y = \cos\left(\frac{\pi}{2} - \frac{1}{x}\right)$ , then  $\frac{dy}{dx} = \dots\dots\dots$

(a)  $-\cos \frac{1}{x}$

(b)  $\frac{1}{x^2} \cos \frac{1}{x}$

(c)  $\frac{-1}{x^2} \sin\left(\frac{\pi}{2} - \frac{1}{x}\right)$

(d)  $-\frac{1}{x^2} \cos\left(\frac{\pi}{2} - \frac{1}{x}\right)$

29 (1<sup>st</sup> session 2021) If  $y = \sqrt{2(\sec x + \tan x)}$  where  $x \in ]0, \frac{\pi}{2}[$ , then  $\frac{1}{y} \left(\frac{dy}{dx}\right) = \dots\dots\dots$

(a)  $2 \sec x$

(b)  $-\frac{1}{2} \sec x$

(c)  $\frac{1}{2} \sec x$

(d)  $-2 \sec x$

30 If  $y = \frac{1}{4} \sec^4 x$ , then  $\frac{dy}{dx} = \dots\dots\dots$

(a)  $4 y \tan x$

(b)  $4 y$

(c)  $\sec^3 x \tan x$

(d)  $y \tan x$

31 If  $y = (\csc x - \cot x)(\csc x + \cot x)$ , then  $\frac{dy}{dx} = \dots\dots\dots$

(a) zero

(b) 1

(c)  $-y \csc x$

(d)  $-y \cot x$

32 If  $f(x) = \tan(x - \theta) \cdot \cot(x + \theta)$ , then  $f'(x) = \dots\dots\dots$  at  $x = \theta$

(a)  $\cot \theta$

(b)  $\tan 2\theta$

(c)  $\cot 2\theta$

(d)  $\tan \theta + \cot \theta$

33 If  $f(x) = x^3 + 5x - 3$ , then  $\frac{d}{dx} [\hat{f}(4)] = \dots\dots\dots$

(a) zero

(b) 4

(c) 24

(d) 53

34 If  $f(x) = \sec x \cos x + \csc x \sin x$ , then  $f\left(\frac{13\pi}{4}\right) + f'\left(\frac{13\pi}{4}\right) = \dots\dots\dots$

(a)  $-\frac{13\pi}{4}$

(b)  $\frac{13\pi}{4}$

(c) 2

(d) undefined.

35 If  $y = \sec x (\sin x + \cos x)$ , then  $\frac{dy}{dx} = \dots\dots\dots$

(a)  $1 - \tan^2 x$

(b)  $\cot^2 x$

(c)  $\sec^2 x$

(d)  $\csc^2 x$

36 If  $x^2 + 4xy - 3y^2 - 2x = 3$ , then  $\frac{dy}{dx}$  could be equal to  $\dots\dots\dots$  at  $x = 3$

(a) 3

(b)  $\frac{1}{3}$

(c)  $-\frac{1}{3}$

(d) -3



- 37 The number of solutions that satisfy :  $x^2 + y^2 = 25$  and  $\frac{dy}{dx} = 2$  is .....
- (a) zero (b) 1  
(c) 2 (d) infinite number
- 
- 38 If :  $y^2 = 2 + xy$ , then  $\frac{dy}{dx}$  ..... at  $x = 1$
- (a) has one value (b) has two values  
(c) has an infinite number of values (d) undefined
- 
- 39 If  $f(x) = x^3 - 5x^2 + 9x - 3$ , then  $f''(0) = \dots\dots\dots$
- (a) -20 (b) -10 (c) 0 (d) 10
- 
- 40 If  $f(n) = x^4$  where  $x$  is constant, then  $f''(n) = \dots\dots\dots$
- (a)  $12x^2$  (b)  $4x^3$  (c) zero (d)  $12n^2$
- 
- 41 If  $f(x) = ax^3 + 3x^2 + 4x + 1$  and  $f'(1) = 6$ , then  $a = \dots\dots\dots$
- (a) zero (b)  $-\frac{4}{3}$  (c) -1 (d)  $-\frac{2}{3}$
- 
- 42 If  $f(x) = \sin 2x$ , then  $f''\left(\frac{\pi}{4}\right) = \dots\dots\dots$
- (a) zero (b) -2 (c) -4 (d) -6
- 
- 43 If  $f(x) = \sin^2 x + \cos^2 x$ , then  $f''(-1) = \dots\dots\dots$
- (a) -1 (b) zero (c) 1 (d) 2
- 
- 44 If  $f(x) = \cot x$ , then  $f''\left(\frac{\pi}{4}\right) = \dots\dots\dots$
- (a)  $-\frac{4}{9}$  (b)  $\frac{4}{9}$  (c) 4 (d)  $\frac{9}{2}$
- 
- 45 If  $f(x) = \sin 2x \cos 2x$ , then  $f''\left(\frac{\pi}{3}\right) = \dots\dots\dots$
- (a) -4 (b) zero (c)  $4\sqrt{3}$  (d) 8
- 
- 46  $\frac{d^2}{dx^2} \left( \cos^2 \frac{x}{2} - \sin^2 \frac{x}{2} \right) = \dots\dots\dots$
- (a)  $-\cos x$  (b)  $\cos x$  (c)  $-\sin x$  (d)  $\cos x - \sin x$



- 47  $\frac{d}{dX} \left( \frac{dy}{dz} \right) = \frac{d^2 y}{dz^2} \times \dots\dots\dots$   
(a)  $\frac{dz}{dX}$  (b)  $\frac{dy}{dX}$  (c)  $\frac{dX}{dz}$  (d)  $\frac{dy}{dz}$
- 
- 48 If  $y = -\sin X$ , then  $\frac{d^2 y}{dX^2} + y = \dots\dots\dots$   
(a) -4 (b) 2 (c) 4 (d) zero
- 
- 49 If  $y = X^2 + 2X$ , and  $\frac{d^2 y}{dX^2} - y - 3 = 0$ , then  $X = \dots\dots\dots$   
(a) 1 (b) 2 (c) -1 (d) -2
- 
- 50 If  $f(X) = (X - 3)^{-1}$ , then  $f'''(3) = \dots\dots\dots$   
(a) zero (b) -6 (c) undefined (d) 2
- 
- 51 If  $f$  is a polynomial function of fifth degree, then the fifth derivative of the function  $f$  equals  $\dots\dots\dots$   
(a) zero (b) non zero constant (c)  $X$  (d)  $5X$
- 
- 52 The third derivative of the quadratic function is a  $\dots\dots\dots$  function.  
(a) linear (b) quadratic (c) constant (d) zero
- 
- 53 The second derivative of the cubic function is a  $\dots\dots\dots$  function.  
(a) linear (b) quadratic (c) constant (d) zero
- 
- 54 If  $f(X) = \frac{X}{X-2}$ , then  $f'''(3) = \dots\dots\dots$   
(a) 12 (b) -8 (c) -12 (d) -16
- 
- 55  $\frac{d}{dX} \left[ y \frac{dy}{dX} \right] = y \frac{d^2 y}{dX^2} + \dots\dots\dots$   
(a)  $\frac{d^2 y}{dX^2}$  (b)  $\left( \frac{dy}{dX} \right)^2$  (c)  $\frac{dy}{dX}$  (d)  $y$
- 
- 56 If  $y = a \sin(mX) + b \cos(mX)$ ,  $\frac{d^2 y}{dX^2} = \dots\dots\dots$   
(a)  $m^2 y$  (b)  $-m^2 y$  (c)  $my$  (d)  $-my$



- 57 If  $y = \sin 3x + \cos 3x$ , then  $\frac{d^4 y}{dx^4} = \dots\dots\dots$   
 (a)  $9y$  (b)  $81y$   
 (c)  $3 \cos 3x - 3 \sin 3x$  (d)  $81x$
- 
- 58 If  $f(x) = \cos 3x \cos x - \sin 3x \sin x$ , then  $f''\left(\frac{\pi}{4}\right) = \dots\dots\dots$   
 (a) 2 (b) 4 (c) 8 (d) 16
- 
- 59  $\frac{d^3}{dx^3}(\sin x \cot x) = \dots\dots\dots$   
 (a)  $-\sin x$  (b)  $-\cos x$  (c)  $\sin x$  (d)  $\cos x$
- 
- 60 If  $x = (1-y)(1+y)(1+y^2)(1+y^4)$ , then  $\frac{d^2 y}{dx^2} = \dots\dots\dots$   
 (a)  $-\frac{1}{8}y^{-7}$  (b)  $-56y^6$  (c)  $-\frac{7}{64}y^{-15}$  (d)  $\frac{7}{8}y^6$
- 
- 61 If  $(x+y)^5 = 3$ , then  $\frac{d^2 y}{dx^2} + \frac{dy}{dx} = \dots\dots\dots$   
 (a) -1 (b) zero (c)  $20(x+y)^4$  (d) 1
- 
- 62 If  $y = f(x)$  satisfies the relation  $\frac{d^4 y}{dx^4} = y$ , then  $y$  can be equal to  $\dots\dots\dots$   
 (a)  $(x+1)^4$  (b)  $\sin x$   
 (c)  $\tan x$  (d) constant value  $\neq 0$
- 
- 63 If  $f(x+1) = x^2 + 2x + 1$ , then  $f''(3) = \dots\dots\dots$   
 (a) 1 (b) 2 (c) 3 (d) 4
- 
- 64 If  $3f(x) + f(4-x) = 6x^2 + 1$ , then  $f''(2) + f'''(2) = \dots\dots\dots$   
 (a) 3 (b) 6 (c) 12 (d) 15
- 
- 65 If  $x^2 - y^2 = 9$  and  $\frac{d^2 y}{dx^2} = \frac{a}{y^3}$ , then  $a = \dots\dots\dots$   
 (a) 1 (b) -1 (c) 9 (d) -9
- 
- 66 If  $f(x) = (x-3)(x-4)(x-k)$  and  $f''(3) = 2$  where  $k \in \mathbb{R}$ , then  $k = \dots\dots\dots$   
 (a) zero (b) 1 (c) 2 (d) 3



- 67 If  $y = x^5 - 10x^4 + 60x^2 + 120$ , then  $\frac{d^2 y}{dx^2} = \dots\dots\dots$  at  $\frac{d^4 y}{dx^4} = 0$
- (a) 300 (b) 200 (c) -200 (d) -300
- 
- 68 If  $f(x) = 2 \sin \frac{x}{2} \cos \frac{x}{2}$ , then the 1000<sup>th</sup> derivative of this function =  $\dots\dots\dots$
- (a)  $[\sin x]^{1000}$  (b)  $\sin x$  (c)  $-\sin x$  (d)  $-\cos x$
- 
- 69 If  $f(x) = e^{3x}$ , then  $\ddot{f}(x) = \dots\dots\dots$
- (a)  $e^{2x}$  (b)  $3e^{3x}$  (c)  $9e^{3x}$  (d)  $3e^{2x}$
- 
- 70 If  $y = e^{ax}$ , then  $\frac{d^4 y}{dx^4} = \dots\dots\dots$
- (a)  $a^4$  (b)  $a^4 y$  (c)  $a e^{ax}$  (d)  $a^2 e^{ax}$
- 
- 71 If  $f(x) = x^2 - 3 \ln 5x$ , then  $\dot{f}(2) = \dots\dots\dots$
- (a) -1 (b) 1 (c)  $\frac{5}{2}$  (d) 6
- 
- 72 If  $f(x) = a e^x$ , then  $\dot{f}(-2)$  equals  $\dots\dots\dots$
- (a)  $-f(2)$  (b)  $-\dot{f}(2)$  (c)  $-f(-2)$  (d)  $f(-2)$
- 
- 73 If  $y = \ln(\sec x + \tan x)$ , then  $\frac{dy}{dx} = \dots\dots\dots$
- (a)  $\tan x$  (b)  $\sec x$  (c)  $\tan^2 x$  (d)  $\csc x$
- 
- 74 If  $y = \ln(\csc x - \cot x)$ , then  $\frac{dy}{dx} = \dots\dots\dots$
- (a)  $\csc x$  (b)  $\cot x$  (c)  $\frac{1}{\csc x - \cot x}$  (d)  $\sin x - \tan x$
- 
- 75 If  $y = (e^{-x} \ln x)$ , then  $\frac{dy}{dx} = \dots\dots\dots$
- (a)  $e^{-x} \left( \frac{1}{x} - \ln x \right)$  (b)  $e^x \left( \frac{1}{x} - \ln x \right)$   
(c)  $\frac{e^{-x}}{x} - \ln x$  (d)  $e^{-x} \left( \frac{1}{x} + \ln x \right)$



76 If  $y = \frac{e^x}{1+x}$ , then  $\dot{y} = \dots\dots\dots$

- (a)  $\frac{xy}{1+x}$  (b)  $\frac{xy}{(1+x)^2}$  (c)  $\frac{ye^x}{(1+x)^2}$  (d)  $\frac{xe^x}{1+x}$

77 If  $f: \mathbb{R} \rightarrow \mathbb{R}$  where  $f(x) = e^x + e^{-x}$ , then  $f(1) + \dot{f}(1) = \dots\dots\dots$

- (a)  $-2e$  (b)  $-e$  (c)  $e$  (d)  $2e$

78 If  $y = x^6 + 6^x$ , then  $\frac{dy}{dx} = \dots\dots\dots$

- (a)  $12x$  (b)  $x+6$   
(c)  $6x^5 + 6^x \ln 6$  (d)  $6x^5 + x \times 6^{x-1}$

79 If  $y = \ln |x^2 - 1|$ , then  $\frac{dy}{dx} = \dots\dots\dots$

- (a)  $2x|x^2 - 1|$  (b)  $\frac{2x}{x^2 - 1}$  (c)  $\ln\left(\frac{2x}{x^2 - 1}\right)$  (d)  $\frac{1}{x^2 - 1}$

80 If  $y = \ln(\tan x)$ , then  $\frac{dy}{dx} = \dots\dots\dots$

- (a)  $2 \sec x \tan x$  (b)  $2 \csc 2x$   
(c)  $2 \cot 2x$  (d)  $-2 \csc x \cot x$

81 If  $y = e^\pi$ , then  $\frac{dy}{dx} = \dots\dots\dots$

- (a) zero (b)  $e^\pi$  (c)  $\pi e^\pi$  (d)  $\frac{1}{\pi} e^\pi$

82 (Trial 2021) If  $y = a e^{bx}$ ,  $\frac{d^2y}{dx^2} = y$ , then  $b^2 = \dots\dots\dots$

- (a) zero (b) 1 (c)  $-1$  (d) 2

83 (Trial 2021) If:  $f(x) = \ln(x^2 + 1)^2 + e^{\sin x}$ , then  $f(0) \times \dot{f}(0) = \dots\dots\dots$

- (a) zero (b) 1 (c)  $1+e$  (d)  $e$

84 If  $y = \ln \frac{e^{x^2}}{x^2}$ , then  $\frac{dy}{dx} = \text{zero at } x = \dots\dots\dots$

- (a) zero (b) 1 (c)  $-1$  (d)  $\pm 1$



- 85 (Trial 2021) If  $y \times \log_{x^3} e = 1$  where  $x > 1$ , then  $\frac{dy}{dx} = \dots\dots\dots$  at  $x = 3$
- (a) zero (b) 1 (c)  $3 \ln 3$  (d)  $\ln 3$
- 
- 86 If  $f : f(x) = \log_x e$ , then  $\dot{f}(x) = \dots\dots\dots$
- (a)  $\frac{1}{x}$  (b)  $-\frac{1}{x}$  (c)  $-\frac{(\log_x e)^2}{x}$  (d)  $\frac{(\ln x)^2}{x}$
- 
- 87 If  $y = \frac{\log_x 2e}{\log_x 3e}$ , then  $\frac{dy}{dx} = \dots\dots\dots$
- (a) zero (b) 1 (c)  $e$  (d)  $\frac{1}{e}$
- 
- 88 Derivative of  $\frac{ax+b}{cx+d}$  with respect to  $\frac{ax+b}{cx+d}$  equals  $\dots\dots\dots$
- (a) zero (b) 1 (c)  $\frac{a}{(cx+d)^2}$  (d)  $\frac{b}{(ax+b)^2}$
- 
- 89 If  $f(x) = \dot{f}(x) = e^{g(x)}$  then  $g(x)$  could be  $\dots\dots\dots$
- (a)  $x+2$  (b)  $x^2$  (c)  $\ln x$  (d)  $2x$
- 
- 90 Which of the following function does satisfy the equation :  $\frac{dy}{dx} - y^2 = 1$  ?
- (a)  $y = \sin x$  (b)  $y = \tan x$  (c)  $y = \sec x$  (d)  $y = \cot x$
- 
- 91 If  $f(x)$  is an even function,  $\dot{f}(0)$  is exist, then  $\dot{f}(0) = \dots\dots\dots$
- (a) zero (b)  $-1$  (c) 1 (d) otherwise
- 
- 92 If  $f(x)$  is an odd function and is differentiable in the interval  $]-\infty, \infty[$ ,  $\dot{f}(3) = 2$ , then  $\dot{f}(-3) = \dots\dots\dots$
- (a) zero (b) 1 (c) 2 (d) 4
- 
- 93  $\frac{d}{dx} (1 + \tan^2 x)^3 = \dots\dots\dots$
- (a)  $6 \sec^5 x \tan x$  (b)  $3 \sec^2 x \tan x$   
(c)  $6 \sec^6 x \tan x$  (d)  $3 \sec^4 x \tan x$



94  $\frac{d}{dX} [(\sec X - 1)(\sec X + 1)] = \dots\dots\dots$

- (a)  $\sec^2 X \tan^2 X$  (b)  $2 \sec^2 X \tan X$  (c)  $\sec^2 X \tan X$  (d)  $\sec^4 X$

95 If  $y = \sec 3X + \tan 3X$ , then  $\frac{dy}{dX} = \dots\dots\dots$

- (a)  $3y \sec 3X$  (b)  $3y \tan 3X$  (c)  $3y \sec^2 3X$  (d)  $3y \tan^2 3X$

96 If  $y = \cot aX$  and  $\frac{dy}{dX} + 4(1 + y^2) = 0$ , then  $a = \dots\dots\dots$

- (a) 1 (b) -2 (c) 4 (d) -4

97 If  $y = \sin^2 3X + \cos^2 3X + \cot^2 3X$  and  $\frac{dy}{dX} = a \csc^2 3X \cot 3X$ , then  $a = \dots\dots\dots$

- (a) -1 (b) -2 (c) -3 (d) -6

98 (Trial 2021) If  $\sin X \cos y = \frac{1}{2}$  where  $X, y$  are measure of acute angles

, then  $\frac{dy}{dX} = \dots\dots\dots$  at  $X = \frac{\pi}{4}$

- (a) 1 (b) -1 (c) zero (d)  $\frac{1}{2}$

99 If  $y = X \sin y$ , then  $\frac{dX}{dy} = \dots\dots\dots$

- (a)  $\frac{1 - \sin y}{X \cos y}$  (b)  $\frac{1 - X \sin y}{X \cos y}$  (c)  $\frac{1 - X \cos y}{\sin y}$  (d)  $\frac{1 - X \cos y}{X \sin y}$

100 If  $y \in ]0, \frac{\pi}{4}[$ ,  $X = \frac{2 \tan y}{1 - \tan^2 y}$ , then  $\frac{dy}{dX} = \dots\dots\dots$

- (a)  $\frac{1}{2} \cos^2 2y$  (b)  $2 \sec^2 2y$  (c)  $\sin^2 2y$  (d)  $\cot 2y$

101 If  $y = \frac{2 \cot X}{\cot^2 X - 1}$ , then  $\frac{dy}{dX} = \dots\dots\dots$  where  $X \in ]0, \frac{\pi}{6}[$

- (a)  $2 \sec^2 2X$  (b)  $2 \cot^2 2X$  (c)  $4 \csc^2 2X$  (d)  $\tan 2X$

102 If  $y = \frac{\sec X \csc X}{\csc^2 X - \sec^2 X}$ , then  $\frac{dy}{dX} = \dots\dots\dots$

- (a)  $\sec^2 (2X)$  (b)  $\frac{1}{2} \tan (2X)$  (c)  $\sec X \csc X$  (d)  $2 \sin^2 (2X)$



## Multiple choice question bank

- 103 If  $y = x \sec x$ , then  $\frac{dy}{dx} = \dots\dots\dots$   
 (a)  $y \tan x + y x^{-1}$  (b)  $\sec x \tan x$  (c)  $x \sec x \tan x$  (d)  $y (\tan x + 1)$
- 
- 104 If  $y = \csc x + \cot x$ , then  $\frac{dy}{dx} = \dots\dots\dots$   
 (a)  $y \csc x$  (b)  $-y \csc x$  (c)  $y \cot x$  (d)  $-y \cot x$
- 
- 105 If  $y = \sec^n(x)$ , then  $\frac{dy}{dx} = \dots\dots\dots$   
 (a)  $n y \sec x$  (b)  $n y \tan x$  (c)  $n y \sec^2 x$  (d)  $n y \tan^2 x$
- 
- 106 If  $y = x \sin x$ , then  $\ddot{y} + y = \dots\dots\dots$   
 (a)  $-\sin x$  (b)  $x$   
 (c)  $2 \cos x$  (d)  $-x \sin x + 2 \cos x$
- 
- 107 If  $x = \sin y$ ,  $y$  is an acute angle, then  $\frac{dy}{dx} = \dots\dots\dots$   
 (a)  $\sqrt{1-x^2}$  (b)  $\frac{1}{\sqrt{1-x^2}}$  (c)  $\sqrt{x^2-1}$  (d)  $\frac{1}{\sqrt{x^2-1}}$
- 
- 108 (Trial 2021) If  $x = \sec y$  where  $y \in ]\frac{\pi}{2}, \pi[$ , then  $\frac{dx}{dy} = \dots\dots\dots$   
 (a)  $x\sqrt{x^2+1}$  (b)  $-x\sqrt{x^2+1}$  (c)  $-x\sqrt{x^2-1}$  (d)  $x\sqrt{x^2-1}$
- 
- 109 If  $x = \tan y$ , then  $\frac{dy}{dx} = \dots\dots\dots$   
 (a)  $x^2 - 1$  (b)  $\frac{1}{x^2 - 1}$  (c)  $\frac{1}{x^2 + 1}$  (d)  $\frac{-1}{x^2 + 1}$
- 
- 110 (2<sup>nd</sup> session 2021) If  $x = \tan(1+y)$ , then  $\frac{dy}{dx} = \dots\dots\dots$  at  $x = 2$   
 (a) 0.5 (b) 0.2 (c) 1 (d) 5
- 
- 111 (2<sup>nd</sup> session 2021) If  $x^2 = e^{2y}$ , then  $x \times \frac{dy}{dx} = \dots\dots\dots$   
 (a) -1 (b) 1 (c)  $x$  (d)  $e^{2x}$
- 
- 112 If  $x = \tan y \csc y$ , then  $\frac{dy}{dx} = \dots\dots\dots$   
 (a)  $\csc y \cot y$  (b)  $\csc y - \cos y$  (c)  $\csc y - \sin y$  (d)  $\sin y + \sec y$



113  $\frac{d^3}{dX^3} (\sin^2 X) = \dots\dots\dots$

(a)  $-4 \sin X$

(b)  $-4 \sin 2X$

(c)  $2 \cos 2X$

(d)  $-\sin 2X$

114  $\frac{d^2}{dX^2} (\cos^4 X + \sin^4 X) = \dots\dots\dots$

(a) zero

(b)  $-2 \sin 2X$

(c)  $-4 \cos 4X$

(d) 1

115 If  $y = \tan X + \frac{1}{3} \tan^3 X$ , then  $\frac{dy}{dX} = (\dots\dots\dots)^4$

(a)  $\csc X$

(b)  $\sec X$

(c)  $\tan X$

(d)  $\sec^4 X$

116 If  $y = \tan X$ , then  $\frac{d^2 y}{dX^2} = \dots\dots\dots$

(a)  $y + y^3$

(b)  $\sec^2 X$

(c)  $2y(1 + y^2)$

(d)  $\sec X \tan X$

117 If  $y = X \tan \frac{X}{2}$ , then  $(1 + \cos X) \frac{dy}{dX} - \sin X = \dots\dots\dots$

(a)  $Xy$

(b)  $y$

(c) zero

(d)  $X$

118 If  $y = e^X \sin X$ , then  $2 \frac{dy}{dX} - \frac{d^2 y}{dX^2} = \dots\dots\dots$

(a)  $2y$

(b)  $4y$

(c)  $5y$

(d)  $8y$

119 If  $y = \ln \left( \frac{e^{4X}}{1 + e^{4X}} \right)$ , then  $\frac{dy}{dX} = \dots\dots\dots$

(a)  $\frac{-1}{1 + e^{4X}}$

(b)  $\frac{2}{1 + e^{4X}}$

(c)  $\frac{-3}{1 + e^{4X}}$

(d)  $\frac{4}{1 + e^{4X}}$

120 If  $f(X) = \ln(\ln X)$ , then  $\ddot{f}(e) = \dots\dots\dots$

(a)  $e^2$

(b)  $e$

(c)  $\frac{1}{e}$

(d)  $\frac{1}{e^2}$

121 If  $y = \frac{z+1}{z-1}$ ,  $X = \frac{z-1}{z+1}$ , then  $\frac{dy}{dX} = \dots\dots\dots$  at  $X = 2$

(a)  $-\frac{1}{8}$

(b)  $-\frac{1}{4}$

(c)  $\frac{1}{4}$

(d)  $-4$

122 If  $y = 3t^2 + 1$ ,  $z = 2t - 5$ , then  $\frac{dy}{dz} = \dots\dots\dots$

(a)  $\frac{t}{3}$

(b)  $3t$

(c) 3

(d)  $\frac{3}{t}$



### Multiple choice question bank

- 123 If  $X = 2t^2 + 3$ ,  $y = \sqrt{t^3}$ , then  $\frac{dy}{dX}$  at  $t = 1$  equals .....
- (a)  $\frac{1}{8}$  (b)  $\frac{1}{4}$  (c)  $\frac{3}{8}$  (d)  $\frac{8}{3}$
- 
- 124 If  $X = at^2$ ,  $y = 2at$ , then  $\frac{dy}{dX} = \dots\dots\dots$
- (a)  $2at$  (b)  $2a$  (c)  $t$  (d)  $\frac{1}{t}$
- 
- 125 If  $y = \cot\left(\frac{\pi}{6}z\right)$ ,  $z = 3\sqrt{X}$ , then  $\frac{dy}{dX} = \dots\dots\dots$  at  $X = 1$
- (a)  $-\frac{\pi}{3}$  (b)  $\frac{\pi}{36}$  (c)  $-\frac{\pi}{6}$  (d)  $-\frac{\pi}{4}$
- 
- 126 If  $y = e^X$ ,  $z = \sin X$ , then  $\frac{dy}{dz} = \dots\dots\dots$
- (a)  $\frac{e^X}{\sin X}$  (b)  $e^X \tan X$  (c)  $e^X \cos X$  (d)  $\frac{e^X}{\cos X}$
- 
- 127 If  $X = \sin 2\pi\theta$ ,  $y = \cos 2\pi\theta$ , then  $\frac{dy}{dX} = \dots\dots\dots$  at  $\theta = \frac{1}{6}$
- (a)  $\sqrt{2}$  (b)  $\sqrt{3}$  (c)  $\frac{1}{\sqrt{2}}$  (d)  $-\sqrt{3}$
- 
- 128 If  $X = a(\theta - \sin \theta)$ ,  $y = a(1 - \cos \theta)$ , then all the following are true except .....
- (a)  $\frac{dX}{d\theta} = y$  (b)  $\frac{dy}{dX} = \cot \frac{\theta}{2}$   
(c)  $\frac{dy}{d\theta} + \frac{dX}{d\theta} = 1$  (d)  $y \frac{dy}{dX} = a \sin \theta$
- 
- 129 If  $X = a \sec \theta$ ,  $y = b \tan \theta$ , then  $\frac{dy}{dX} = \frac{2b}{a}$  at  $\theta = \dots\dots\dots$  where  $\theta$  is positive acute angle.
- (a)  $\frac{\pi}{12}$  (b)  $\frac{\pi}{6}$  (c)  $\frac{\pi}{4}$  (d)  $\frac{\pi}{3}$
- 
- 130 If  $X = e^{\sin \theta}$ ,  $\frac{dy}{d\theta} = 2^{\log_2 \cos \theta}$ , then  $\frac{dy}{dX} = \dots\dots\dots$  at  $\theta = \text{zero}$
- (a)  $\frac{2}{e}$  (b)  $1$  (c)  $e$  (d)  $\frac{1}{e}$
- 
- 131 If  $X = a\left(t - \frac{1}{t}\right)$ ,  $y = a\left(t + \frac{1}{t}\right)$ , then  $\frac{dy}{dX} = \dots\dots\dots$
- (a)  $-\frac{X}{y}$  (b)  $\frac{X}{y}$  (c)  $-\frac{y}{X}$  (d)  $\frac{y}{X}$



132 If  $y = t^3 - t$ ,  $t = \frac{1}{z^2} + z$ ,  $z = 2x - 1$ , then  $\left(\frac{dy}{dx}\right)_{x=1} = \dots\dots\dots$

(a) -22                      (b) -20                      (c) -18                      (d) -15

133 If  $y = (5x - 4)(x + 3)$ ,  $z = 3x^2 - 4x + 17$ , then  $\frac{d^2y}{dx^2} + \frac{d^2z}{dx^2} = \dots\dots\dots$

(a) 12                      (b) 14                      (c) 16                      (d) 18

134 If  $x = \frac{t}{t+1}$ ,  $y = \frac{t+1}{t}$ , then  $\frac{d^2y}{dx^2} = \frac{\dots\dots\dots}{x^2}$

(a)  $\frac{2}{x}$                       (b)  $2x^{-3}$                       (c)  $-x^{-2}$                       (d) zero

135 If  $x = \sec z$ ,  $\sqrt{y} = \tan z$ , then  $\frac{d^2y}{dx^2} = \dots\dots\dots$

(a)  $2 \tan z \sec z$                       (b)  $\sec^2 z \tan^2 z$                       (c) 3                      (d) 2

136 If  $\frac{dz}{dx} = 2x - 3$ ,  $\frac{dy}{dx} = x^2 + 1$ , then  $\frac{d^2z}{dy^2}$  at  $x = 1$  equals  $\dots\dots\dots$

(a) 1                      (b)  $\frac{3}{4}$                       (c)  $\frac{3}{2}$                       (d)  $\frac{4}{3}$

137 If  $y = x^2 + 3x + 2$ ,  $z = 3x^2 - 5x + 4$ , then  $\frac{d^2y}{dz^2}$  at  $x = 2$  equals  $\dots\dots\dots$

(a)  $-\frac{4}{7}$                       (b)  $-\frac{4}{49}$                       (c) 8                      (d) 56

138 If  $x = 2t^3 + 3$ ,  $y = t^4$ , then  $\frac{d^2y}{dx^2} = \dots\dots\dots$  at  $t = 1$

(a)  $\frac{2}{3}$                       (b)  $\frac{1}{9}$                       (c) 4                      (d)  $\frac{1}{3}$

139 If  $\frac{dz}{d\theta} = \cos 2\theta$ ,  $\frac{dy}{d\theta} = \sin 2\theta$ , then  $\frac{d^2y}{dz^2} = \dots\dots\dots$  at  $\theta = \frac{\pi}{8}$

(a)  $4\sqrt{2}$                       (b) 4                      (c)  $2\sqrt{2}$                       (d) 2

140 If  $y = \tan x$ ,  $z = \cot x$ , then  $\frac{d^2y}{dz^2} = \dots\dots\dots$

(a)  $2 \tan^2 x$                       (b)  $-2 \tan^3 x$                       (c)  $2 \tan^3 x$                       (d)  $-2 \cot^3 x$

141 If  $x = \ln t$ ,  $y = t^2 - 1$ , then  $\frac{d^2y}{dx^2} = \dots\dots\dots$  at  $t = 1$

(a) 2                      (b) 3                      (c) 4                      (d) 6



142 If  $x = \cos \theta$ ,  $y = \cos 2\theta$ , then  $\frac{d^2 y}{d x^2} = \dots\dots\dots$

(a) 4 (b)  $4 \sin \theta$  (c)  $4 \cos \theta$  (d)  $-4 \sin \theta$

143 If  $x^2 + y^2 = t + \frac{1}{t}$ ,  $x^4 + y^4 = t^2 + \frac{1}{t^2}$  where  $xy > 0$ , then  $\frac{d^2 y}{d x^2} = \dots\dots\dots$

(a)  $\frac{1}{x^3}$  (b)  $\frac{2}{x^3}$  (c)  $\frac{2\sqrt{2}}{x^3}$  (d)  $\frac{2\sqrt{2}}{x^4}$

144 If  $y = x^9 - 14x^7 - x^2 + 3$ , then  $\frac{d^{10} y}{d x^{10}} = \dots\dots\dots$

(a) 9 (b) 10 (c) zero (d) 9

145 If  $y = x^7$ , then  $\frac{d^7 y}{d x^7} = \dots\dots\dots$

(a)  $7x^6$  (b)  $42x^5$  (c) 49 (d) 7

146 If  $y = x^n$  where  $x \neq 0$  and  $n \in \mathbb{Z}^+$ , then the smallest value of  $n$  to make  $\frac{d^3 y}{d x^3} \neq \text{zero}$  is  $\dots\dots\dots$

(a) 1 (b) 2 (c) 3 (d) 4

147 If  $y = x^n$  where  $n$  is a natural number and  $\frac{d^4 y}{d x^4} = 360x^{n-4}$ , then the value of  $n = \dots\dots\dots$

(a) 7 (b) 13 (c) 5 (d) 6

148 If  $f(x) = 20x^{n-1}$  and  $f'(x) = c$  where  $c \in \mathbb{R}$ ,  $n \in \mathbb{Z}^+$ , then  $n + c = \dots\dots\dots$

(a) 104 (b) 123 (c) 124 (d) 125

149 If  $f(x) = \frac{x^{25}}{25}$ , then  $f^{(25)}(x) = \dots\dots\dots$

(a) 25 (b) 25 (c) 1 (d) zero

150 If  $f(x) = x \ln x$  and  $\frac{d^8 y}{d x^8} = \frac{k}{x^7}$ , then  $k = \dots\dots\dots$

(a) 5 (b) 6 (c) 7 (d) 8

151 If  $y$  is a function in  $x$ , then  $\frac{d}{d x} [y^{(4)}] = \dots\dots\dots$

(a)  $4y^{(3)}$  (b)  $4y^{(3)} \frac{d y}{d x}$  (c)  $y^{(5)}$  (d)  $\frac{1}{5} y^{(5)}$



- 152 If  $y = x^{2021}$ , then ..... = zero  
 (a)  $y^{(2019)}$  (b)  $y^{(2020)}$  (c)  $y^{(2021)}$  (d)  $y^{(2022)}$
- 
- 153 If  $y = x^{2021}$ , then ..... = 2021  
 (a)  $y^{(2019)}$  (b)  $y^{(2020)}$  (c)  $y^{(2021)}$  (d)  $y^{(2022)}$
- 
- 154 If  $y = x^{n+1} + n x^{n-1} + 1$ , then  $\frac{d^n y}{d x^n} = \dots\dots\dots$   
 (a)  $\underline{n+1}$  (b)  $x \underline{n+1}$  (c)  $x \underline{n}$  (d)  $x^{-1} \underline{n}$
- 
- 155 If  $y = \ln x$ , then  $\frac{d^{10} y}{d x^{10}} = \dots\dots\dots$   
 (a)  $\frac{9}{-x^{10}}$  (b)  $\frac{10}{-x^9}$  (c)  $\frac{9}{x^{10}}$  (d)  $\frac{10}{x^9}$
- 
- 156 If  $f(x) = 3x^4 - 2x^2 + 1$ , then .....  
 (a)  $f'''(x) = 36x^2 - 4$  (b)  $f^{(4)}(x) = \text{zero}$   
 (c)  $f'''(0) = -4$  (d)  $f^{(n)}(x) = \text{zero for every } n \geq 5$
- 
- 157 If  $f(x) = x^{\frac{4}{3}}$ , then .....  
 (a)  $f^{(n)}(\text{zero})$  is undefined for every  $n \geq 2$  (b)  $f''(\text{zero}) = \frac{4}{3}$   
 (c)  $f''(\text{zero}) = \frac{4}{9}$  (d)  $f^{(3)}(1) = \frac{8}{27}$
- 
- 158 If  $f(x) = x^n$  where  $n$  is natural number,  $x \neq 0$  and  $\frac{d^m(y)}{d x^m} \neq \text{zero}$ , then .....  
 (a)  $m \geq n$  (b)  $m \leq n$  (c)  $m = n + 1$  (d)  $m > n$
- 
- 159 If  $f$  is a polynomial function and  $f^{(4)}(x)$  is of 7<sup>th</sup> degree, then  $f^{(7)}(x)$  is from ..... degree.  
 (a) 10<sup>th</sup> (b) 7<sup>th</sup> (c) 4<sup>th</sup> (d) 3<sup>rd</sup>
- 
- 160 If  $f(x) = a^x$ , then  $f^{(n)}(\text{zero}) = \dots\dots\dots$   
 (a) 1 (b)  $\ln a$  (c)  $(\ln a)^n$  (d)  $n \ln a$



161 If  $y = e^a x$ ,  $\frac{d^n y}{d x^n} = b e^a x$ , then .....

(a)  $a = b$

(b)  $a = b^n$

(c)  $b = a^n$

(d)  $\frac{a}{b} = n$

162 If  $y = \ln x$ ,  $n$  is a positive integer, then  $\frac{d^n (y)}{d x^n} = \dots$

(a)  $\left(\frac{-e}{x}\right)^n$

(b)  $(n-1) x^{-n}$

(c)  $(n+1) x^{-n-1}$

(d)  $(-1)^{n-1} \frac{n-1}{x^n}$

163 If  $xy = a$  (where  $a$  is a positive real number) and  $\frac{d^2 y}{d x^2} \times \frac{d^2 x}{d y^2} > \frac{d y}{d x} \times \frac{d x}{d y}$ , then  $a \in \dots$

(a)  $]2, \infty[$

(b)  $]0, 4[$

(c)  $]4, \infty[$

(d)  $]0, 2[$

164 If  $y = \sin(ax)$ , then  $y^{(2017)} = \dots$

(a)  $a^{2017} y$

(b)  $-a^{2017} y$

(c)  $a^{2017} \cos(ax)$

(d)  $-a^{2017} \cos(ax)$

165 If  $y = \cos(ax)$ , then  $y^{(2020)} = \dots$

(a)  $a^{2020} y$

(b)  $-a^{2020} y$

(c)  $a^{2020} \sin(ax)$

(d)  $-a^{2020} \sin(ax)$

166 If  $y = \sin 5x$  and  $\frac{d^{20} y}{d x^{20}} = a \sin 5x$ , then  $a = \dots$

(a) 20

(b)  $5^{20}$

(c)  $-5^{20}$

(d) 20

167 If  $(x, y)$  is any point on the unit circle, then .....

(a)  $xy - 2(y)^2 + 1 = 0$

(b)  $xy + (y)^2 + 1 = 0$

(c)  $xy + (y)^2 - 1 = 0$

(d)  $xy + 2(y)^2 + 1 = 0$

168 If  $f(2x+1) = x$ ,  $h(2x-5)$  and  $h(1) = 2$ ,  $\hat{h}(1) = 4$ , then  $\hat{f}(7) = \dots$

(a) 4

(b) 7

(c) 11

(d) 13

169 If  $f(x) + \hat{f}(x) = x^3 + 5x^2 + x + 2$ , then  $f$  is a polynomial function, then  $f(x) = \dots$

(a)  $x^3 + 2x^2 - 3x$

(b)  $3x^2 + 4x - 3$

(c)  $x^3 + 2x^2 + 5$

(d)  $x^3 + 2x^2 - 3x + 5$



170 If  $f(x) = e^x g(x)$ ,  $g(0) = 2$ ,  $g'(0) = 1$ , then  $f'(0) = \dots\dots\dots$

- (a) 1 (b) 3 (c) 2 (d) zero

171 If  $f(x) = \ln(x \cdot g(x))$  and  $g(4) = g'(4) = 4$ , then  $f'(4) = \dots\dots\dots$

- (a) 1 (b)  $\frac{5}{4}$  (c)  $\frac{3}{2}$  (d) 2

172 If  $y = f(x)$  and  $f(x+h) - f(x) = 5x^2h + h^2$ , then  $\frac{d^3y}{dx^3} = \dots\dots\dots$

- (a) 10 (b)  $10x$  (c)  $5x^2$  (d) zero

173 The rate of change of the volume of a sphere with respect to its surface area when  $r = 2$  cm. is  $\dots\dots\dots$  cm.

- (a) 1 (b) 2 (c) 3 (d) 4

174 A cube of side length  $l$  length unit and its volume =  $V$  cubic unit and its total surface area =  $A$  square unit, then  $\frac{d^2V}{dA^2} = \dots\dots\dots$  when  $l = \frac{1}{4}$

- (a)  $\frac{1}{4}$  (b)  $\frac{1}{12}$  (c)  $\frac{1}{16}$  (d)  $\frac{1}{48}$

175 If  $y = \sin 2x \cos 2x$  where  $\frac{\ddot{y}}{y} + a^2 \leq 0$ , then  $a \in \dots\dots\dots$

- (a)  $[-4, 4]$  (b)  $] -4, 4[$  (c)  $\mathbb{R} - ] -4, 4[$  (d)  $\mathbb{R} - [-4, 4]$

176 If  $f(x) = \begin{cases} x^3 & \text{when } x < 2 \\ 4x & \text{when } x \geq 2 \end{cases}$ , then  $f'(2) = \dots\dots\dots$

- (a) 2 (b) 4 (c) 1 (d) does not exist

177 If  $f$  is an even function and  $f'$  exists, then  $f'(h) + f'(-h) \dots\dots\dots$  zero

- (a) = (b) < (c) > (d)  $\neq$

178 If  $f : f(x) = \begin{cases} ax^2 + bx + c & x < 0 \\ 2x + 5 & x \geq 0 \end{cases}$  is differentiable twice at  $x = 0$ , then  $a = \dots\dots\dots$

- (a) zero (b) 2 (c) 5 (d) 1



179 (Trial 2021) If  $\frac{dy}{dx} = \sqrt{ay}$ ,  $\frac{d^2y}{dx^2} = 3$ , then the value of  $a = \dots\dots\dots$

- (a) 6 (b) 5 (c) 3 (d) 4

180 If  $y = \frac{x-3}{x+4}$  and  $(y-1)\ddot{y} = k(\dot{y})^2$ , then  $k = \dots\dots\dots$

- (a) 1 (b) 2 (c) 3 (d) 4

181 If  $y = \sqrt{e^{ax}}$  and  $\ddot{y} + 4\dot{y} + 4y = 0$ , then  $a = \dots\dots\dots$

- (a) 4 (b) -4 (c) 16 (d) -16

182 If  $x = ae^\theta(\sin\theta - \cos\theta)$ ,  $y = ae^\theta(\sin\theta + \cos\theta)$ , then  $\left(\frac{dy}{dx}\right)_{\theta=\frac{\pi}{4}} = \dots\dots\dots$

- (a) 1 (b) 2 (c)  $\frac{2}{\pi}$  (d)  $\frac{1}{\pi}$

183 If  $f(x)$ ,  $g(x)$  are two differentiable functions at  $x=2$  and  $h(x) = f(x) \cdot g(x)$  where  $f(2) = 2$ ,  $\dot{f}(2) = 3$ ,  $\ddot{f}(2) = 12$ ,  $g(2) = 3$ ,  $\dot{g}(2) = 4$ ,  $\ddot{g}(2) = 6$ , then  $\ddot{h}(2) = \dots\dots\dots$

- (a) 66 (b) 144 (c) zero (d) 60

184 If  $f, g$  are two functions and  $\dot{f}(x) = -f(x)$ ,  $\dot{g}(x) = g(x)$ , then  $\frac{d}{dx}[(f(x))^2 + (g(x))^2] = \dots\dots\dots$

- (a)  $\dot{f}(x) + \dot{g}(x)$  (b) 1 (c)  $-\dot{f}(x)$  (d) zero

185 If  $\dot{f}(x) = xf(x)$ ,  $f(3) = -5$ , then  $\ddot{f}(3) = \dots\dots\dots$

- (a) -50 (b) -40 (c) 15 (d) 27

186 If  $0 < x < \frac{\pi}{2}$ ,  $f(\sin x) = \sin^2 x$ , then  $\ddot{f}(1) = \dots\dots\dots$

- (a) 1 (b) 2 (c)  $\pi$  (d)  $\frac{\pi}{2}$

187 (Trial 2021) If  $f\left(\frac{1}{2}x\right) = |x|^3$ , then  $\ddot{f}(-1) = \dots\dots\dots$

- (a) -48 (b) 48 (c) 14 (d) 1

188 If  $f(x^2 - 1) = mx^2 + 4x + 1$  where  $x > 0$ ,  $\dot{f}(3) = 4$ , then value of  $m = \dots\dots\dots$

- (a) 3 (b) 4 (c) 6 (d) 12



- 189 If  $y = 1 + \frac{x}{1} + \frac{x^2}{2} + \frac{x^3}{3} + \dots$  to  $\infty$ , then  $\dot{y} + \ddot{y} = \dots$
- (a)  $x$  (b) 1 (c)  $y$  (d)  $2y$
- 
- 190 If  $y = 1 + x + x^2 + x^3 + \dots$  to  $\infty$  where  $|x| < 1$ , then  $\frac{d^2 y}{dx^2} = \dots$  at  $x = \frac{1}{2}$
- (a) 2 (b) 4 (c) 8 (d) 16
- 
- 191 If  $y = e^x \times e^{2x} \times e^{3x} \times \dots \times e^{10x}$  and  $\frac{d^2 y}{dx^2} = a e^{bx}$ , then  $\frac{a}{b} = \dots$
- (a) 1 (b) 10 (c) 55 (d)  $(55)^2$
- 
- 192 (1<sup>st</sup> session 2021) If  $y = \frac{1}{(x+1)^2}$ , then the rate of change of  $\frac{1}{y}$  with respect to  $x^2$  at  $x = 1$  is  $\dots$
- (a)  $\frac{1}{2}$  (b) 4 (c) 2 (d)  $\frac{1}{4}$
- 
- 193 The rate of change of  $e^{x^3}$  with respect to  $\ln x$  equals  $\dots$
- (a)  $3x^2 e^{x^3} + 3x^2$  (b)  $e^{x^3}$  (c)  $3x^3 e^{x^3}$  (d)  $3x^2 e^{x^3}$
- 
- 194 The rate of change of  $\sin x^3$  respect to  $\cos x^3$  equals  $\dots$
- (a)  $-\cot x^3$  (b)  $\cot x^3$  (c)  $\tan x^3$  (d)  $-\tan x^3$
- 
- 195 If  $x = \sin^3 \theta$ ,  $y = \cos^3 \theta$ , then  $\frac{d^2 y}{dx^2} = \dots$  at  $\theta = \frac{\pi}{4}$
- (a)  $\frac{2}{3}$  (b)  $\frac{8}{3}$  (c)  $\frac{4\sqrt{3}}{3}$  (d)  $\frac{4\sqrt{2}}{3}$
- 
- 196 The rate of change of  $(x - \sin x)$  with respect to  $(1 - \cos x)$  at  $x = \frac{\pi}{3}$  equals  $\dots$
- (a)  $\frac{\sqrt{3}}{3}$  (b)  $2\sqrt{3}$  (c)  $\sqrt{3}$  (d)  $\frac{2}{3}$
- 
- 197  $\frac{d}{dx} \left( \sum_{n=0}^{\infty} \frac{1}{n} \right) = \dots$
- (a) zero (b) 1 (c)  $\ln x$  (d)  $\ln + 1$
- 
- 198 If  $y = \frac{e^x + 1}{e^x - 1}$ , then  $\frac{y^2}{2} + \frac{dy}{dx} = \dots$
- (a) 1 (b) -1 (c)  $-\frac{1}{2}$  (d)  $\frac{1}{2}$



- 199 If  $f(x) = \ln(x + \sqrt{x^2 + 1})$ , then  $f'(x) = \dots\dots\dots$
- (a)  $\sqrt{x^2 + 1}$  (b)  $\frac{x}{\sqrt{x^2 + 1}}$  (c)  $1 + \frac{x}{\sqrt{x^2 + 1}}$  (d)  $\frac{1}{\sqrt{x^2 + 1}}$
- 
- 200 If  $y = \ln \sqrt{\tan x}$ , then  $\frac{dy}{dx} = \dots\dots\dots$  when  $x = \frac{\pi}{4}$
- (a) 1 (b) zero (c)  $\frac{1}{2}$  (d)  $\infty$
- 
- 201 If  $f(x) = e^x \sin x$ ,  $g(x) = e^x \cos x$ , which of the following statements is not true?
- (a)  $f'(x) = f(x) + g(x)$  (b)  $g'(x) = g(x) - f(x)$   
(c)  $f''(x) = 2g(x)$  (d)  $g''(x) = 2f(x)$
- 
- 202 If  $y = \frac{1 + \tan x}{1 - \tan x}$ , then  $\frac{dy}{dx} = \dots\dots\dots$
- (a)  $\cos^2(x + \frac{\pi}{4})$  (b)  $\sec^2(x + \frac{\pi}{4})$   
(c)  $\sin^2(x + \frac{\pi}{4})$  (d)  $\csc^2(x + \frac{\pi}{4})$
- 
- 203 If  $y = \ln(\sin x)$ , then  $\frac{d^2y}{dx^2} = \dots\dots\dots$
- (a)  $-\csc^2 x$  (b)  $\sec x$  (c)  $-\csc x \cot x$  (d)  $\sec x \tan x$
- 
- 204 If  $f(x) = e^{\ln x}$ , then  $f'(x) = \dots\dots\dots$
- (a)  $[\ln x] e^{\log x}$  (b)  $e^{\ln x}$  (c)  $(\ln x) e^{x - \ln x}$  (d) 1
- 
- 205 If  $f(x) = e^{\ln(x^3 - 2x + 1)}$ , then  $f'(0) = \dots\dots\dots$
- (a) -4 (b) -2 (c) zero (d) 2
- 
- 206 If  $y = x^2 \ln e^x$ , then  $\frac{dy}{dx} = \dots\dots\dots$
- (a)  $3x^2$  (b)  $x^3$  (c)  $\ln x^3$  (d) zero
- 
- 207 If  $y = e^{(1 + \ln x)}$ , then  $\frac{dy}{dx} = \dots\dots\dots$
- (a)  $x$  (b)  $e x$  (c)  $e$  (d) 1
- 
- 208 If  $f(x) = \ln(\sin x) - \ln(\cos x)$ , then  $f'(\frac{\pi}{4}) = \dots\dots\dots$
- (a) 2 (b) -2 (c) 1 (d) -1



- 209 If  $y = x^{\sin x}$ , then  $\frac{dy}{dx} = \dots\dots\dots$  at  $x = \frac{\pi}{2}$   
 (a) 1 (b) 2 (c)  $\frac{\pi}{2}$  (d)  $\frac{2}{\pi}$
- 
- 210 If  $f(x) = (\cos x)^{\cos x}$ , then  $\dot{f}(\text{zero}) = \dots\dots\dots$   
 (a) -3 (b) -2 (c) -1 (d) zero
- 
- 211 If  $y = \sqrt{x + \sqrt{x + \sqrt{x + \sqrt{x + \dots}}}}$ , then  $\frac{dy}{dx} = \dots\dots\dots$   
 (a)  $2y - 1$  (b)  $\frac{1}{2y - 1}$  (c)  $x^2 + y$  (d)  $1 + x + x^2 + \dots$
- 
- 212 If  $a^y = b^x$  where  $a, b \in \mathbb{R}^+$ , then  $\frac{dy}{dx} = \dots\dots\dots$   
 (a)  $\log \frac{a}{b}$  (b)  $\log_a b$  (c)  $\log_b a$  (d)  $\log \frac{b}{a}$
- 
- 213 If  $y = x^x$ ,  $x > 0$ , then  $\frac{dy}{dx} = \dots\dots\dots$   
 (a)  $\ln x$  (b)  $2 + \ln x$  (c)  $x^x \ln x$  (d)  $x^x (1 + \ln x)$
- 
- 214 If  $y^x = x^y$ , then  $\frac{dy}{dx} = \dots\dots\dots$   
 (a)  $\frac{y}{x}$  (b)  $\frac{y(x \ln y - y)}{x(y \ln x - x)}$   
 (c)  $\frac{x}{y}$  (d)  $\frac{x \ln y}{y \ln x}$
- 
- 215 If  $y = x e^{xy}$ , then  $\frac{dy}{dx} = \dots\dots\dots$   
 (a)  $e^{xy} + x$  (b)  $\frac{y + xy}{x}$   
 (c)  $x e^{xy} + y e^{xy}$  (d)  $\frac{y(1 + xy)}{x(1 - xy)}$
- 
- 216 If  $f: \mathbb{R}^+ \rightarrow \mathbb{R}^+$  where  $f(x) = x^{2x}$ , then  $\dot{f}(e) = \dots\dots\dots$   
 (a)  $4e^{2e}$  (b)  $2e^{2e}$  (c)  $2e^e$  (d)  $4e^e$
- 
- 217 If  $y = \left(\frac{x^a}{x^b}\right)^{a+b} \times \left(\frac{x^b}{x^c}\right)^{b+c} \times \left(\frac{x^c}{x^a}\right)^{c+a}$ , then  $\frac{dy}{dx} = \dots\dots\dots$   
 (a) 1 (b) zero (c)  $a + b + c$  (d)  $2(a + b + c)$



- 218 If  $f(x) = 2x^3 + 1$ ,  $g(x) = x^2$ , then  $\hat{f}(g(3)) = \dots\dots\dots$   
(a) zero (b) 9 (c) 486 (d) 2916
- 219 If  $f(x) = \frac{2}{x+1}$ ,  $g(x) = 3x$ , then  $\frac{d}{dx}[(f \circ g)(x)] = \dots\dots\dots$  at  $x = -2$   
(a)  $-\frac{3}{25}$  (b) 6 (c)  $\frac{1}{25}$  (d)  $-\frac{6}{25}$
- 220 If  $f(x) = 3x^2 - 2$ , then  $(f \circ f)(-1) = \dots\dots\dots$   
(a) -36 (b) -18 (c) zero (d) 18
- 221 (Trial 2021) If  $\hat{f}(x) = \frac{1}{1+x^2}$ ,  $g(x) = \tan x$ , then  $(f \circ g)(x) = \dots\dots\dots$   
(a)  $\sec^2 x$  (b)  $\sec^2 x \tan^2 x$  (c)  $\cos^2 x$  (d) 1
- 222 If  $f(x) = e^x$ ,  $g(x) = \ln x$ , then  $(g \circ f)(x) = \dots\dots\dots$   
(a) 1 (b)  $x$  (c)  $e^x$  (d)  $\ln x$
- 223 If  $f, g$  are two functions where  $f(x) = x^2$ ,  $g(2) = 3$ ,  $\hat{g}(2) = -2$ ,  $\hat{\hat{g}}(2) = 5$ , then  $(f \circ \hat{\hat{g}})(2) = \dots\dots\dots$   
(a) 16 (b) 24 (c) 32 (d) 38
- 224 If  $y = x^2 + 5x + 3$ , then  $\frac{dy}{d(x^2)} = \dots\dots\dots$  at  $x = 1$   
(a)  $\frac{7}{2}$  (b)  $\frac{5}{7}$  (c) 7 (d)  $\frac{7}{5}$
- 225 If  $y = f(f(f(x)))$  and  $f(1) = 1$ ,  $\hat{f}(1) = 3$  then  $[\hat{y}]_{x=1} = \dots\dots\dots$   
(a) 3 (b) 9 (c) 27 (d) 81
- 226 If  $\lim_{h \rightarrow 0} \frac{\hat{f}(1) - \hat{f}(1+h)}{2h} = 21$  where  $f(x) = 2x^4 - ax^3$ , then  $a = \dots\dots\dots$   
(a) 11 (b) 12 (c) 13 (d) 14
- 227 If  $\sin x = e^y$  where  $0 < x < \pi$ , then  $\frac{dy}{dx} = \dots\dots\dots$   
(a)  $\tan x$  (b)  $\cot x$  (c)  $-\tan x$  (d)  $-\cot x$



228 If  $y = 4^{\log_2 \sin x} + 9^{\log_3 \cos x}$ , then  $\frac{dy}{dx} = \dots\dots\dots$

- (a) zero (b) 1 (c)  $\sin x + \cos x$  (d) -1

229 If  $k, m \in \mathbb{R}$ , then  $f(x) = x e^x$  and  $f^{(10)}(x) = k e^x + m x e^x$ , then  $k + m = \dots\dots\dots$

- (a) 9 (b) 10 (c) 11 (d) 12

230 If  $f(x) = \begin{vmatrix} 2x & 7 & 6 \\ 0 & 3x^2 & 4 \\ 0 & 0 & x^3 \end{vmatrix}$ , then  $\dot{f}(1) = \dots\dots\dots$

- (a) 1 (b) 6 (c) 36 (d) 72

231 If  $f(x) = \begin{vmatrix} g(x) & h(x) \\ I(x) & J(x) \end{vmatrix}$ , then  $\dot{f}(x) = \dots\dots\dots$

- (a)  $\begin{vmatrix} g(x) & h(x) \\ I(x) & J(x) \end{vmatrix} + \begin{vmatrix} g(x) & h(x) \\ \dot{I}(x) & \dot{J}(x) \end{vmatrix}$  (b)  $\begin{vmatrix} \dot{g}(x) & \dot{h}(x) \\ I(x) & J(x) \end{vmatrix} + \begin{vmatrix} g(x) & h(x) \\ \dot{I}(x) & \dot{J}(x) \end{vmatrix}$   
 (c)  $\begin{vmatrix} g(x) & \dot{h}(x) \\ I(x) & \dot{J}(x) \end{vmatrix} + \begin{vmatrix} \dot{g}(x) & h(x) \\ I(x) & \dot{J}(x) \end{vmatrix}$  (d)  $\begin{vmatrix} g(x) & h(x) \\ \dot{I}(x) & \dot{J}(x) \end{vmatrix} + \begin{vmatrix} \dot{g}(x) & h(x) \\ I(x) & J(x) \end{vmatrix}$

232 If  $f: \mathbb{R} - \{\frac{\pi}{2} + \pi n, n \in \mathbb{Z}\} \rightarrow \mathbb{R}$  where  $f(x) = e^{\tan x}$ , then for all values of  $x$  we find that  $\dots\dots\dots$

- (a)  $\dot{f}(x) \geq f(x)$  (b)  $\dot{f}(x) \leq f(x)$   
 (c)  $\dot{f}(x) > f(x)$  (d)  $\dot{f}(x) < f(x)$

233 If  $0 < x < \frac{\pi}{2}$  and  $f(\sin x) = a \cos x$  where  $a$  is a constant and  $\dot{f}(\frac{3}{5}) = -6$ , then  $a = \dots\dots\dots$

- (a) 2 (b) 4 (c) 6 (d) 8

234 If  $f(x) = g(x^2)$ , then  $\frac{\dot{f}(2)}{\dot{g}(4)} = \dots\dots\dots$

- (a) 1 (b) 2 (c) 3 (d) 4



235 If the function  $g$  is the inverse of the function  $f$  where  $f$  and  $g$  are differentiable functions on  $\mathbb{R}$  and  $\hat{f}(a) = 2$ ,  $f(a) = b$ , then  $\hat{g}(b) = \dots\dots\dots$

- (a)  $\frac{1}{2}$  (b) 2 (c)  $\frac{2}{3}$  (d) 1

236 If  $f: [1, \infty[ \rightarrow \mathbb{R}$ ,  $f(x) = 2x^2 - 4x + 5$  and  $g$  is the inverse function of the function  $f$ , then  $\hat{g}(5) = \dots\dots\dots$

- (a) 4 (b) 2 (c)  $\frac{1}{2}$  (d)  $\frac{1}{4}$

237 If  $y = \cot x$  where  $x$  is in degree, then  $\frac{dy}{dx} = \dots\dots\dots$

- (a)  $-\csc^2 x$  (b)  $-\csc x \cot x$   
(c)  $-\frac{\pi}{180} \csc^2 x$  (d)  $-\frac{180}{\pi} \csc^2 x$

238 If  $y = \sin 2x$  and  $\frac{d^n y}{dx^n} = 2^n \sin 2x$ , then  $\dots\dots\dots$

- (a)  $n$  is an even number. (b)  $n$  is an odd number.  
(c)  $n$  is divisible by 3 (d)  $n$  is divisible by 4


239 If  $f(x) = \frac{e^{5x} + e^{4x} + e^{3x}}{e^{2x} + e^x + 1}$ , then  $\hat{f}(0) = \dots\dots\dots$

- (a) 1 (b) 2 (c) 3 (d) 4

240 In the opposite figure :

If  $\overline{AB}$  is a tangent to the circle  
, then  $\frac{dy}{dx} = \dots\dots\dots$  at  $x = 4$


- (a) 0.4 (b) 0.6 (c) 0.8 (d) 0.9



241 In the opposite figure :

If  $\overline{AD}$  bisects  $\angle BAC$   
, then  $\frac{dy}{dx} = \dots\dots\dots$


- (a)  $-\csc 2x \cot 2x$  (b)  $-2 \csc x \cot x$   
(c)  $-2 \csc 2x$  (d)  $-2 \csc 2x \cot 2x$






242 If the two opposite figures show the relations :

$y = f(t)$  ,  $x = g(t)$  , then  $\frac{dy}{dx}$  could be equals .....




- (a) zero
- (b) a positive number.
- (c) a negative number.
- (d) a non-negative number.

243 The opposite graph represents each of the two functions  $f(x) = a^x$  and  $f'(x)$  , then .....




- (a)  $0 < a < 1$
- (b)  $1 < a < e$
- (c)  $a = e$
- (d)  $a > e$

244 If  $y = a^x$  where  $0 < a < 1$  , then which of the following graphs does represent  $\dot{y}$  ?



245 If  $y = a x^n - b x^{n+1} + 5$  is a polynomial and  $a, b \in \mathbb{R}^+$  , then  $\frac{d^n y}{d x^n}$  may be represented by the figure .....





246 If  $f$ ,  $g$ ,  $h$  are differentiable functions with respect to  $x$ , by using the given variables in the following table :

**First :**  $h(x) = 3f(x) - 2g(x)$ , then  $h'(1) = \dots\dots\dots$

(a)  $-5$

(b)  $2$

(c)  $1$

(d)  $-1$

$x$	1	2
$f(x)$	$-1$	$4$
$g(x)$	$2$	$1$
$f'(x)$	$1$	$5$
$g'(x)$	$2$	$-3$

**Second :**  $h(x) = f(x) [5 + g(x)]$ , then  $h'(2) = \dots\dots\dots$

(a)  $8$

(b) Zero

(c)  $18$

(d)  $-18$

**Third :**  $h(x) = f(x) \div [g(x) + 2]$ , then  $h'(1) = \dots\dots\dots$

(a)  $\frac{-1}{4}$

(b)  $\frac{1}{4}$

(c)  $\frac{3}{8}$

(d)  $\frac{-3}{8}$

**Fourth :**  $h(x) = f[g(x)]$ , then  $h'(1) = \dots\dots\dots$

(a)  $2$

(b) Zero

(c)  $10$

(d)  $-6$

**Fifth :**  $h(x) = g[3x - f(x)]$ , then  $h'(2) = \dots\dots\dots$

(a)  $6$

(b)  $-8$

(c)  $-1$

(d)  $8$

**Sixth :**  $h(x) = [x^3 + g(x)]^{-2}$ , then  $h'(1) = \dots\dots\dots$

(a)  $\frac{-2}{27}$

(b)  $\frac{-10}{27}$

(c) Zero

(d)  $-30$



## Third

## Questions on the geometric applications (the two equations of the tangent and the normal to a curve)

Choose the correct answer from the given ones :

- 1 If  $f(x) = 3x + g(x)$ , and  $g'(2) = -5$ , then the slope of the tangent to the curve of the function  $f$  at  $x = 2$  equals .....  
 (a)  $-2$                       (b)  $3$                       (c)  $-15$                       (d)  $\frac{1}{2}$

---

- 2 If the tangent to the curve  $y = f(x)$  at the point  $(3, 4)$  makes an angle of measure  $\frac{3\pi}{4}$  with the positive direction of  $x$ -axis, then  $f'(3) = \dots\dots\dots$   
 (a)  $-1$                       (b)  $-\frac{3}{4}$                       (c)  $\frac{3}{4}$                       (d)  $1$

---

- 3 The tangent to the curve :  $y = 3x^2 - 5$  at the point  $(1, -2)$  passes through the point .....  
 (a)  $(5, -2)$                       (b)  $(3, 1)$                       (c)  $(2, -4)$                       (d)  $(0, -8)$

---

- 4 If the curves of the two functions  $f(x)$  and  $g(x)$  are touching at the point  $(2, 4)$ , and  $f'(2) = 3$ , then  $g'(2) = \dots\dots\dots$   
 (a)  $2$                       (b)  $3$                       (c)  $4$                       (d)  $5$

---

- 5 The slope of the normal to the curve of the function  $y = |x^3|$  at the point  $(-2, 8)$  is .....  
 (a)  $-12$                       (b)  $12$                       (c)  $\frac{1}{12}$                       (d)  $-\frac{1}{12}$

---

- 6 The slope of the tangent to the curve  $y = e^x$  at the point  $(1, e)$  lies on it equals .....  
 (a)  $1$                       (b)  $-1$                       (c)  $e$                       (d)  $e^2$

---

- 7 The slope of the tangent to the curve  $y = \ln x$  at the point  $(e^2, 2)$  lies on it equals .....  
 (a)  $2$                       (b)  $-2$                       (c)  $e^2$                       (d)  $e^{-2}$

---

- 8 If  $f : f(x) = x - x \ln x$ , then the slope of the tangent to the curve at  $x = e$  equals .....  
 (a) zero                      (b)  $-1$                       (c)  $1$                       (d)  $e$



- 9 If the tangent drawn from the point  $(2, 4)$  to the curve  $y = \frac{1-2x}{x-2}$  touches the curve at the point B, then B = .....
- (a)  $(1, 1)$       (b)  $(2, 2)$       (c)  $(3, -5)$       (d)  $(-1, -1)$
- 
- 10 The measure of the angle which the tangent to the curve  $\sin 2x = \tan y$  makes with the positive direction of  $x$ -axis at the point  $(\frac{3\pi}{4}, \frac{3\pi}{4})$  equals .....
- (a) zero      (b)  $135^\circ$       (c)  $45^\circ$       (d)  $26^\circ 34'$
- 
- 11 The tangent to the curve :  $x^2 - xy + y^2 = 27$  which drawn at the point  $(6, 3)$ , makes an angle of measure ..... with the positive direction of  $x$ -axis.
- (a)  $90^\circ$       (b) zero      (c)  $45^\circ$       (d)  $180^\circ$
- 
- 12 The slope of the tangent to the curve  $y = t^2 + 2t$ ,  $x = t^2 - 2t$  equals ..... at  $t = 2$
- (a)  $\frac{1}{3}$       (b)  $-\frac{1}{2}$       (c) 2      (d) 3
- 
- 13 Which of the following curves has a tangent with constant slope ?
- (a)  $x = \sin t$ ,  $y = \cos t$       (b)  $x = t^2$ ,  $y = 3t^2$   
(c)  $x = 2t - 1$ ,  $y = t^2 - 4$       (d)  $xy = 7$
- 
- 14 The slope of the tangent to the curve of the circle  $x^2 + y^2 = 1$  at  $x = \frac{3}{5}$  equals .....
- (a)  $\pm \frac{4}{3}$       (b)  $\pm \frac{3}{4}$       (c)  $\frac{4}{5}$       (d)  $\pm \frac{4}{5}$
- 
- 15 If the tangent to the curve of the function  $y = x^2 + a$  at the point  $(1, b)$  intersects the  $x$ -axis at  $x = -1$ , then  $a \times b = \dots\dots\dots$
- (a) 3      (b) 4      (c) 12      (d) -12
- 
- 16 The equation of the tangent to the curve  $x = y^2$  at the origin point is .....
- (a)  $y = 0$       (b)  $x = 0$       (c)  $x = y$       (d)  $x + 4y = 0$
- 
- 17 If the equation of the normal to the curve  $y = f(x)$  at the point  $(1, 1)$  is  $x + 4y = 5$ , then  $f'(1) = \dots\dots\dots$
- (a) -3      (b)  $-\frac{1}{4}$       (c) 4      (d) -4



18 The equation of the normal to the curve of the function :  $y = x|x|$  at the point  $(-2, -4)$  is .....

(a)  $y + 4x + 12 = 0$

(b)  $4y + x + 18 = 0$

(c)  $4y + x + 14 = 0$

(d)  $y + 4x - 4 = 0$

19 The equation of the normal to the curve :  $y = \sin x$  at the point  $(0, 0)$  is .....

(a)  $x = 0$

(b)  $y = 0$

(c)  $x + y = 0$

(d)  $x - y = 0$

20 The equation of the tangent to the curve of the function  $f$  where  $f(x) = e^{2x+1}$  at the point  $(-\frac{1}{2}, 1)$  is .....

(a)  $2y = x + 1$

(b)  $y = 2x + 2$

(c)  $y = 2x - 3$

(d)  $2y = 3x + 1$

21 (Trial 2021) The slope of the curve  $x^y - y^x = 0$  at the point  $(1, 1)$  which lies on it equals .....

(a) 1

(b) zero

(c) 2

(d) -1

22 (Trial 2021) Slope of tangent to the curve  $y = 5^x \log_5(x+1)$  at  $x = 0$  equals .....

(a)  $\ln 5$

(b)  $\log_5 e$

(c) zero

(d)  $5 \log_5 e$

23 Equation of the tangent to the curve  $y + \ln(xy) = x$  at the point  $(1, 1)$  is .....

(a)  $x = 1$

(b)  $x = -1$

(c)  $y = 1$

(d)  $y = -1$

24 The equation of the normal to the curve  $y = 3e^x$  at point lies on it and its  $x$ -coordinate is  $-1$  is .....

(a)  $e^2 x = 0$

(b)  $y - \frac{3}{e} = \frac{3}{e}(x+1)$

(c)  $y - 3 = \frac{e}{3}(x+1)$

(d)  $e^2 x + 3ey + e^2 - 9 = 0$

25 The equation of the normal to the curve  $y = e^{2x} \cos x$  at  $x = 0$  is .....

(a)  $y + 2x = 1$

(b)  $2y + x = 2$

(c)  $x + y = 2$

(d)  $y - 2x = 1$



- 26 If  $y = X + c$  is a tangent to the curve  $9X^2 + 16Y^2 = 144$ , then  $c = \dots\dots\dots$   
(a)  $\pm 2$  (b)  $\pm 3$  (c)  $\pm 5$  (d)  $\pm 6$
- 27 The normal to the circle  $X^2 + Y^2 = 12$  at any point on it, passes through the point  $\dots\dots\dots$   
(a)  $(2, 2)$  (b)  $(1, 1)$  (c)  $(0, 0)$  (d)  $(-2, -2)$
- 28 If  $X = a(1 + \cos \theta)$ ,  $Y = a \sin \theta$ , then the perpendicular to the curve passes through a fixed point which is  $\dots\dots\dots$   
(a)  $(a, 0)$  (b)  $(0, a)$  (c)  $(1, 0)$  (d)  $(a, a)$
- 29 (Trial 2021) If equation of tangent to the curve  $Y = -\ln X$  at a point lies on it  $(a, b)$  is  $Y = mX$ , then  $a = \dots\dots\dots$   
(a)  $e$  (b)  $\frac{1}{2}e$  (c)  $\frac{3}{2}e$  (d)  $2\sqrt{e}$
- 30 (2<sup>nd</sup> Session 2021) If the equation of the tangent to the curve :  $\log_y (4X) = 2X$  at the point  $(1, 2)$  which lies on it is  $Y = mX + c$ , then  $m = \dots\dots\dots$   
(a)  $1 - 2 \ln 2$  (b)  $2 \ln 2 - 1$  (c)  $2 \ln 2 + 1$  (d)  $2 + \ln 2$
- 31 The ratio between the slope of the tangent to the curve :  $Y = \ln (3\sqrt{X+1})$  and the slope of the tangent to the curve :  $Y = \ln (5\sqrt{X+1})$  at  $X = a$  equals the ratio  $\dots\dots\dots$   
(a)  $3 : 5$  (b)  $5 : 3$  (c)  $1 : 1$  (d)  $\ln 3 : \ln 5$
- 32 The rate of change of slope of the tangent of the function  $f(X) = 2X^3$  at  $X = 3$  equals  $\dots\dots\dots$   
(a) 36 (b) 54 (c) 6 (d) 12
- 33 If the normal to the curve  $Y = X \ln X$  parallel to the straight line  $2X - 2Y + 3 = 0$ , then the equation of this normal is  $\dots\dots\dots$   
(a)  $X - Y = 3e^{-2}$  (b)  $X - Y = 6e^{-2}$   
(c)  $X - Y = 3e^2$  (d)  $X - Y = 6e^2$



34 (1<sup>st</sup> session 2021) The equation of the normal to the curve  $y = \ln (\tan x)$  at the point which lies on the curve and its  $x$ -coordinate equals  $\frac{\pi}{4}$  is .....

- (a)  $4x - 8y = \pi$  (b)  $8y + 4x = \pi$   
(c)  $4x + 2y = \pi$  (d)  $4x - 2y = \pi$

35 If the tangent to the curve of the function  $f : f(x) = ax^3$  at  $x = 1$  is perpendicular to the tangent to the curve of the function  $h : h(x) = b \sin^2 x$  at  $x = \frac{\pi}{4}$ , then  $a \times b =$  .....

- (a)  $\frac{4}{3}$  (b)  $\frac{2}{3}$  (c)  $\frac{1}{3}$  (d)  $-\frac{1}{3}$

36 (2<sup>nd</sup> session 2021) The tangent of the curve  $2x + 1 = \sin y$  is parallel to  $y$ -axis at the point .....

- (a)  $(\frac{\pi}{2}, 0)$  (b)  $(-\frac{1}{2}, \pi)$  (c)  $(0, \frac{\pi}{2})$  (d)  $(-\frac{1}{2}, 0)$

37 If the tangent to the curve :  $y^2 = 4ax$  is perpendicular to  $x$ -axis, then .....

- (a)  $\frac{dy}{dx} = 0$  (b)  $\frac{dy}{dx} = 1$  (c)  $\frac{dx}{dy} = 1$  (d)  $\frac{dx}{dy} = 0$

38 The tangent to the curve  $x = t^2 - 1$ ,  $y = t^2 - t$  parallel to  $x$ -axis at  $t =$  .....

- (a) zero (b)  $\frac{1}{\sqrt{3}}$  (c)  $\frac{1}{2}$  (d)  $-\frac{1}{\sqrt{3}}$

39 The tangent to the curve :  $x = 3 \cos \theta$ ,  $y = 3 \sin \theta$  where  $(0 \leq \theta \leq \pi)$ , parallel to  $x$ -axis if  $\theta =$  .....

- (a) zero (b)  $\frac{\pi}{3}$  (c)  $\frac{\pi}{2}$  (d)  $\pi$

40 (Trial 2021) If  $y = f(x)$  where  $y = \sqrt[3]{n^2 + 7}$ ,  $6n^2x + n = 1$ , then the perpendicular to the curve of the function at a point lies on it and its  $x$ -coordinate equals zero is parallel to .....

- (a)  $x$ -axis (b)  $y$ -axis  
(c) the straight line  $y = x$  (d) the straight line  $y = -x$



- 41 (2<sup>nd</sup> session 2021) The slope of the normal to the curve  $x = e^t$ ,  $y = e^{2t+2}$  at the point which lies on the curve and its  $x$ -coordinate equals one is .....
- (a)  $-2e^2$       (b)  $-\frac{1}{2e^2}$       (c)  $2e^2$       (d)  $\frac{e^2}{2}$
- 
- 42 (Trial 2021) If  $y^2 = 8x$  where  $y > 0$ , then the point lies on this curve at which  $\frac{dy}{dx} = \frac{dx}{dy}$  is .....
- (a)  $(0, 0)$       (b)  $(1, 2\sqrt{2})$       (c)  $(\frac{1}{2}, 2)$       (d)  $(2, 4)$
- 
- 43 The curve  $y = x^{\frac{1}{5}}$  at  $(0, 0)$  has .....
- (a) a vertical tangent.      (b) a horizontal tangent.  
(c) an inclined tangent.      (d) no tangent.
- 
- 44 If the curve  $x = 2z^3 - 5z^2 - 4z + 12$ ,  $y = 2z^2 + z - 4$  has a horizontal tangent, then  $z =$  .....
- (a)  $-\frac{1}{4}$       (b)  $-\frac{1}{3}$       (c)  $2$       (d)  $-\frac{2}{3}$
- 
- 45 The curve  $y - e^{xy} + x = 0$  has a vertical tangent at the point .....
- (a)  $(1, 1)$       (b)  $(0, 0)$       (c)  $(1, 0)$       (d)  $(2, e^2)$
- 
- 46 The tangent to the curve  $x = e^\theta \cos \theta$ ,  $y = e^\theta \sin \theta$  at the point which at  $\theta = \frac{\pi}{4}$  makes with the positive direction of  $x$ -axis an angle of measure .....
- (a) zero      (b)  $\frac{\pi}{4}$       (c)  $\frac{\pi}{3}$       (d)  $\frac{\pi}{2}$
- 
- 47 The slope of the tangent to the curve  $x = y^{100} + \log(100)^y$  at the point  $(3, 1)$  equals .....
- (a) 102      (b) 100      (c)  $100 \times 3^{99}$       (d)  $\frac{1}{102}$
- 
- 48 If the tangent to the curve :  $y = x^3 - 3x^2$  makes obtuse angle with the positive direction of  $x$ -axis, then  $x \in$  .....
- (a)  $[0, 2]$       (b)  $]0, 2[$       (c)  $\mathbb{R} - [0, 2]$       (d)  $\mathbb{R} - ]0, 2[$



- 49 If the straight line :  $y + x - 1 = 0$  touches the curve of the function  $f : f(x) = x^2 - 3x + a$ , then  $a = \dots\dots\dots$
- (a) 1                      (b) 2                      (c) 3                      (d) 4
- 
- 50 If the straight line :  $y = x + c$  where  $c > 0$  touches the curve  $y = \frac{x}{x+1}$  at the point  $(a, b)$ , then  $a, b = \dots\dots\dots$
- (a) -5                      (b) -4                      (c) -1                      (d) zero
- 
- 51 The tangent to the curve of the function  $y = \sqrt[3]{x}$  at  $x = 0$  is parallel to  $\dots\dots\dots$
- (a)  $x$ -axis                      (b)  $y$ -axis  
(c) the straight line  $y = x$                       (d) the straight line  $x + y = 0$
- 
- 52 If the curve :  $y = x^2 - ax + a - 1$  touches  $x$ -axis where  $a \in \mathbb{R}$ , then  $a = \dots\dots\dots$
- (a) 2                      (b) zero                      (c)  $-\frac{1}{2}$                       (d) -1
- 
- 53 The area of the triangle bounded by two coordinate axes and the tangent to the curve  $xy = a^2$  at the point  $(x_1, y_1)$  lies on it equals  $\dots\dots\dots$
- (a)  $\frac{a^2 x_1}{y_1}$                       (b)  $\frac{a^2 y_1}{x_1}$                       (c)  $2a^2$                       (d)  $4a^2$
- 
- 54 Area of triangle included between perpendicular to the curve  $x = e^{\sin y}$  at the point  $(1, 0)$  and the two coordinate-axes equals  $\dots\dots\dots$  area unit.
- (a)  $\frac{1}{2}$                       (b)  $\frac{1}{4}$                       (c)  $\frac{3}{4}$                       (d) 1
- 
- 55 Tangent to the curve  $x^2 y^2 + 2x - 3y = 0$  at the point  $(1, 1)$  does not pass through the point  $\dots\dots\dots$
- (a)  $(0, -3)$                       (b)  $(-1, -7)$                       (c)  $(2, 5)$                       (d)  $(3, 0)$
- 
- 56 The curve  $\left(\frac{x}{a}\right)^n + \left(\frac{y}{b}\right)^n = 2$  touches the straight line  $\frac{x}{a} + \frac{y}{b} = 2$  at the point  $(a, b)$  when  $\dots\dots\dots$
- (a)  $n = 3$                       (b)  $n = 2$                       (c) for all values of  $n$                       (d) false for all values of  $n$



# Multiple choice question bank

- 57 The equation of the tangent to the curve  $y = be^{-\frac{x}{a}}$  at the point of intersection with y-axis is .....
- (a)  $\frac{x}{a} - \frac{y}{b} = 1$       (b)  $ax + by = 1$       (c)  $ax - by = 1$       (d)  $\frac{x}{a} + \frac{y}{b} = 1$
- 
- 58 The straight line :  $ax + by + c = 0$  is a normal to the curve  $xy = 1$  , then .....
- (a)  $a > 0, b > 0$       (b)  $a < 0, b < 0$   
 (c)  $a = 0, b \neq 0$       (d)  $a > 0, b < 0$  or  $a < 0, b > 0$
- 
- 59 Length of the intercepted part from y-axis by the tangent to the curve  $y = x \sin x$  at  $x = \pi$  equals ..... length unit.
- (a)  $-\pi$       (b)  $\pi$       (c)  $-\pi^2$       (d)  $\pi^2$
- 
- 60 Length of the intercepted part from the y-axis by the tangent to the curve  $y = \ln(\ln x)$  at  $x = e$  equals .....
- (a)  $\frac{1}{e}$       (b)  $e$       (c)  $1$       (d) zero
- 
- 61 If the function  $f$  is even and  $2x + y + 5 = 0$  is the equation of the tangent to the curve at  $x = a$  , then equation of tangent to the curve at  $x = -a$  is .....
- (a)  $2x + y + 5 = 0$       (b)  $-2x + y + 5 = 0$   
 (c)  $\frac{1}{2}x + y + 5$       (d)  $2x + y - 5 = 0$
- 
- 62 If the normal to the curve :  $9y^2 = x^3$  at the point  $(a, b)$  which lies on the curve cuts equal parts of the coordinate axes , then  $a =$  .....
- (a)  $2$       (b)  $-4$       (c)  $4$       (d)  $-2$
- 
- 63 If the tangent to the curve :  $2y^3 = ax^2 + x^3$  at the point  $(a, a)$  which lies on the curve cuts from the coordinate axes two parts of lengths  $L, M$  where  $L^2 + M^2 = 61$  , then  $a =$  .....
- (a)  $\pm 20$       (b)  $\pm 30$       (c)  $\pm 40$       (d)  $\pm 50$
- 
- 64 Slope of the tangent to the curve  $y = \ln \tan t + \ln \cos t, x = \ln \sin t + \ln \cot t$  at  $t = \frac{\pi}{4}$  is .....
- (a)  $-1$       (b) zero      (c)  $1$       (d)  $2$



- 65 Measure of the angle that the tangent to the curve  $y = e^{\tan x}$  makes with the positive direction of the  $x$ -axis at  $x = \pi$  equals .....
- (a)  $\frac{\pi}{4}$  (b)  $\frac{\pi}{3}$  (c)  $\frac{\pi}{2}$  (d)  $\frac{3\pi}{4}$
- 
- 66 If  $f(x) = \frac{x+1}{g(x)}$ ,  $g(x) \neq 0$  and the curve of  $g(x)$  has a horizontal tangent at the point  $(1, 2)$ , then  $f'(1) = \dots\dots\dots$
- (a) 2 (b) 1 (c) -2 (d)  $\frac{1}{2}$
- 
- 67 If the equation of the normal to the common tangent of the two functions  $f$  and  $g$  at  $x = 1$  is  $y = -\frac{1}{3}x + \frac{3}{2}$ , then  $(f \times g)'(1) = \dots\dots\dots$
- (a) 4 (b) 7 (c) 2 (d) 10
- 
- 68 Length of the tangent segment drawn from the point  $(-4, 0)$  to the curve  $y^2 = 4x$  equals ..... length unit.
- (a)  $4\sqrt{3}$  (b)  $4\sqrt{5}$  (c)  $4\sqrt{6}$  (d) 8
- 
- 69 Length of the perpendicular drawn from the origin to the tangent to the curve  $y = e^x + x$  at the point  $(0, 1)$  equals ..... length unit.
- (a) 1 (b)  $\sqrt{5}$  (c)  $2\sqrt{5}$  (d)  $\frac{1}{\sqrt{5}}$
- 
- 70 Measure of the included angle between the two tangents to the curve  $y = x^2 - 5x + 6$  drawn from the two intersection points with the  $x$ -axis equals .....
- (a)  $\frac{\pi}{2}$  (b)  $\frac{\pi}{3}$  (c)  $\frac{\pi}{4}$  (d)  $\frac{\pi}{6}$
- 
- 71 If  $L_1$  and  $L_2$  are two tangents to the curve  $y = \sqrt{3}e^x$ ,  $y = \sqrt{3}e^{-x}$  at their intersection point, then measure of the included angle between  $L_1$  and  $L_2$  equals .....
- (a)  $\frac{\pi}{2}$  (b)  $\frac{\pi}{3}$   
(c)  $\frac{\pi}{4}$  (d)  $\frac{\pi}{6}$



72 In the opposite figure :


If the straight line  $l$  is a tangent of the function  $f$  at the point  $C$  and cuts  $x$ -axis at the point  $A(-4, 0)$  and if  $B(4, 0)$ ,  $f(4) + f'(4) = 9$ , then the area of  $\triangle ABC = \dots\dots\dots$  square unit.

(a) 9

(b) 64

(c) 32

(d) 16



73 In the opposite figure :


If  $f(x) = g(x) - 3h(x)$ , then  $f'(5) = \dots\dots\dots$

(a) zero

(b) 2

(c) 3

(d) 4



74 In the opposite figure :


The straight line  $L$  is a tangent to the curve  $y = f(x)$  at  $(-2, m)$  and  $g(x) = f(2x)$ , then  $g'(-1) = \dots\dots\dots$

(a) 3

(b) 4

(c) 6

(d) 9




75 The opposite figure represents the function  $f$  and the straight line  $L$  touches the curve of  $f$  at the point  $A(3, 5)$  and  $g(x) = x \cdot f(x)$ , then  $g'(3) = \dots\dots\dots$

(a) 3

(b) 1

(c) 5


(d) 8





**76 In the opposite figure :**


If the straight line  $L$  touches the curve  $y = f(x)$  at  $(-3, 3)$  and  $h(x) = \frac{f(2-x)}{x-2}$ , then  $h(5) = \dots\dots\dots$



- (a) zero (b) -2 (c) 2 (d) 3


**77 (Trial 2021) The opposite figure represents the curve**

$e^{x+y} = 3x - 2$  if the tangent to the curve at the point  $A(1, -1)$  intersects the two coordinate axes  $\vec{OX}$ ,  $\vec{OY}$  at the two points  $B$  and  $C$  respectively, then  $A$  divides  $\vec{BC}$  by the ratio  $\dots\dots\dots$



- (a) 1 : 2 internally. (b) 2 : 1 internally.  
(c) 1 : 2 externally. (d) 2 : 1 externally.


**78** The opposite figure represents the function  $y = \ln x$  and the tangent to the curve at any point  $(l, m)$  intersects the  $y$ -axis at  $C$ , then length of  $\vec{BC} = \dots\dots\dots$  length unit.



- (a)  $\frac{1}{e}$  (b)  $e$   
(c) 1 (d)  $c$

**79 In the opposite figure :**

If the straight line  $\vec{AB}$  is a tangent to the curve  $x = \sqrt{e^y + 2}$  at the point  $A$ , then area of  $\Delta OAB = \dots\dots\dots$  area unit.




- (a)  $\sqrt{3}$   
(b)  $2\sqrt{3}$   
(c)  $3\sqrt{3}$   
(d)  $6\sqrt{3}$



80 In the opposite figure :

If  $y_1 = \sec X$ ,  $y_2 = \csc X$  and  $l_1, l_2$  touch the two curves  $y_1, y_2$  respectively at the two points  $(\frac{11\pi}{6}, \frac{2}{\sqrt{3}}), (\frac{-2\pi}{3}, \frac{-2}{\sqrt{3}})$ , then  $\tan \theta = \dots\dots\dots$

- (a)  $-\frac{2}{3}$  (b)  $\frac{2}{3}$   
(c)  $-\frac{12}{5}$  (d)  $\frac{12}{5}$



#### Fourth Questions on related time rates

Choose the correct answer from the given ones :

- 1 A square with side length 5 cm. , the length of its side start to increase by rate 2 cm./sec. , then the side length of the square after t sec. given by the relation .....  
(a)  $2t$  (b)  $5 + 2t$  (c)  $2t - 5$  (d)  $5 + 4t^2$
- 2 The length of a rectangle is twice its width , if the rate of change of the length is 6 cm./sec. , then the rate of change of its width = ..... cm./sec.  
(a) 12 (b) 3 (c) 6 (d) - 3
- 3 The length of a rectangle is three times its width , if the rate of change of its width is 3 cm./sec. , then the rate of change of its diagonal length = ..... cm./sec.  
(a) 9 (b)  $2\sqrt{10}$  (c)  $3\sqrt{10}$  (d)  $\sqrt{105}$
- 4 If the side length of an equilateral triangle increases by rate 2 cm./sec. , then the perimeter of the triangle increases by rate ..... cm./sec.  
(a) 2 (b) 8 (c) 4 (d) 6
- 5 If the height of an equilateral triangle increases at rate  $\sqrt{3}$  cm./sec. , then the rate of change of its side length equals ..... cm./sec.  
(a) 4 (b) 2 (c)  $\frac{4}{3}$  (d)  $\frac{3}{4}$



- 6 The side length of a cube decreases by rate 3 cm./sec. , then the length of the diagonal of the cube is decreasing at a rate ..... cm./sec.  
 (a)  $3\sqrt{2}$  (b)  $3\sqrt{3}$  (c) 6 (d) 9
- 
- 7 A point moves on the curve  $y = 2x + 1$  , then the ratio between the rate of change of the  $x$ -coordinate with respect to time for a point to the rate of change of the  $y$ -coordinate with respect to time equals .....  
 (a) 2 (b) -2 (c)  $\frac{1}{2}$  (d)  $-\frac{1}{2}$
- 
- 8 If the radius of a circle increases by rate  $\frac{4}{\pi}$  cm./sec. , then the circumference of the circle increases at this instant by rate ..... cm./sec.  
 (a)  $\frac{4}{\pi}$  (b)  $\frac{\pi}{4}$  (c)  $\frac{1}{8}$  (d) 8
- 
- 9 A metal disc , the rate of decreasing of its diameter by cooling is 0.5 cm./sec. , then the decreasing rate of its surface area = ..... cm<sup>2</sup>/sec. when its radius 14 cm. ( $\pi = \frac{22}{7}$ )  
 (a) 5.5 (b) 11 (c) 16 (d) 22
- 
- 10 An empty container , its volume 45 cm<sup>3</sup> , water is poured in it at a rate 5 cm<sup>3</sup>/sec. , the container becomes full after ..... sec.  
 (a) 9 (b) 135 (c) 45 (d) 50
- 
- 11 The side length of a cube increases by rate 5 cm./sec. , then the volume of the cube increases by rate ..... cm<sup>3</sup>/sec. at the side length = 10 cm.  
 (a) 1500 (b) 150 (c) 45 (d) 50
- 
- 12 A cubic water tank of side length 4 m. , water is poured in it by rate  $\frac{1}{2}$  m<sup>3</sup>/min. , then the rate of increasing of water height in the tank = ..... m./min.  
 (a)  $\frac{1}{96}$  (b)  $\frac{1}{32}$  (c)  $\frac{1}{24}$  (d)  $\frac{1}{48}$
- 
- 13 A cubic water tank of side length 5 m. , water is poured in it by rate  $\frac{1}{4}$  m<sup>3</sup>/min. , then the rate of change of the upper surface area of the water = ..... m<sup>2</sup>/min.  
 (a) zero (b) 1 (c)  $\frac{1}{8}$  (d)  $\frac{1}{100}$



- 14 A circle touches the sides of a square internally, then the rate of change of the radius of the circle equals ..... the rate of change of the side length of the square at any instant.  
(a) double (b) half (c) quarter (d) four times
- 15 A square inscribed in a circle, then the rate of change of the radius of the circle equals ..... the rate of change of the square side at any point.  
(a) 1 (b)  $\frac{\sqrt{2}}{2}$  (c)  $\frac{1}{2}$  (d)  $\sqrt{2}$
- 16 A square lamina expands regularly, the rate of increasing in surface area of the lamina  $75 \text{ cm}^2/\text{sec.}$ , then the rate of increasing of side length = .....  $\text{cm./sec.}$  when the side length is 5 cm.  
(a) 2.5 (b) 5 (c) 7.5 (d) 15
- 17 If  $y = x^2 - 3x$ , then  $\frac{dy}{dt} = \frac{dx}{dt}$  at  $x = \dots\dots\dots$   
(a) 1 (b) 2 (c) 3 (d) 4
- 18 If a particle moves on the curve :  $y^2 + x^2 = 10$  such that  $\frac{dy}{dt} = 4$ , then  $\frac{dx}{dt} = \dots\dots\dots$  at the point  $(\sqrt{5}, -\sqrt{5})$   
(a) 2 (b)  $2\sqrt{5}$  (c) 4 (d)  $4\sqrt{5}$
- 19 A point moves on the curve :  $x^2 + y^2 - 5x + 3y - 6 = 0$ , if the rate of change of its  $x$ -coordinate respect to the time  $t$  at the point  $(1, 2)$  equals 3, then the rate of change of its  $y$ -coordinate respect to the time is .....  
(a)  $1\frac{2}{7}$  (b)  $\frac{7}{12}$  (c) 3 (d)  $-\frac{9}{7}$
- 20 A point  $(x, y)$  moves on the curve :  $y = x^2 - \frac{1}{4}$ , then the position of this point at the moment which the rate of change of its  $y$ -coordinate respect to the time equals three times the rate of change of  $x$ -coordinate respect to the time is .....  
(a)  $(6, \frac{143}{4})$  (b)  $(\frac{1}{2}, 0)$  (c)  $(3, \frac{35}{4})$  (d)  $(\frac{3}{2}, 2)$



- 21 A point moves on the curve :  $y = x - \frac{x}{x^2 + 1}$ , the rate of change of its  $x$ -coordinate with respect to time equals 9 at  $x = \sqrt{2}$ , then the rate of change of its  $y$ -coordinate with respect to time at the same point = .....
- (a) 4 (b) 6 (c) 8 (d) 10
- 
- 22 If  $x$  is the radian measure of an angle,  $x \in ]0, \frac{\pi}{2}[$ , then the rate of increasing of tangent of an angle equals 8 times the rate of increasing of sine the same angle at  $x = \dots\dots\dots$
- (a)  $\frac{\pi}{3}$  (b)  $\frac{\pi}{4}$  (c)  $\frac{\pi}{5}$  (d)  $\frac{\pi}{6}$
- 
- 23 A stone fell in still water, then a circular waves is formed whose radius increases at rate of 3 cm./sec., then the rate of increase of surface area of the wave after 4 sec. equals .....  $\text{cm}^2/\text{sec}$ .
- (a)  $8\pi$  (b)  $72\pi$  (c)  $12\pi$  (d)  $24\pi$
- 
- 24 A square lamina, its side length changes by rate 0.2 cm./sec., then rate of change of its surface area .....  $\text{cm}^2/\text{sec}$ . when its diagonal length =  $8\sqrt{2}$  cm.
- (a)  $\frac{1}{25}$  (b) 3.2 (c)  $\frac{4}{10}$  (d) 16
- 
- 25 The radius of a circle increases by rate 2 cm./min., and its area by rate  $20\pi \text{ cm}^2/\text{m}$ , then length of its radius at this moment equals ..... cm.
- (a)  $\frac{5}{2}$  (b) 5 (c) 10 (d) 20
- 
- 26 A cube melt preserving its shape by rate  $1 \text{ cm}^3/\text{sec}$ , then the rate of change of its edge length when its volume  $8 \text{ cm}^3$  is .....  $\text{cm}/\text{sec}$ .
- (a)  $\frac{1}{12}$  (b)  $\frac{1}{192}$  (c)  $\frac{-1}{24}$  (d)  $\frac{-1}{12}$
- 
- 27 If the rate of change of area of circle equals the rate of change of its diameter numerically, then  $r = \dots\dots\dots$
- (a)  $\frac{2}{\pi}$  (b)  $\frac{1}{\pi}$  (c)  $\frac{\pi}{2}$  (d)  $\pi$
- 
- 28 If the side length of an equilateral triangle =  $a$ , and it increases by rate  $k$ , then the rate of increasing of the area of the triangle equals .....
- (a)  $\frac{2}{\sqrt{3}} a k$  (b)  $\sqrt{3} a k$  (c)  $\frac{\sqrt{3}}{2} a k$  (d)  $\frac{\sqrt{2}}{\sqrt{3}} a k$



- 29 The surface area of a sphere increases by constant rate its value  $6 \text{ cm}^2/\text{sec}$ . at the moment which its radius was equal  $30 \text{ cm}$ . , then the rate of increasing of its volume = .....  $\text{cm}^3/\text{sec}$ .

(a) 180 (b) 40 (c) 90 (d)  $90\pi$

- 30 If the rate of change of the volume of a sphere equals the rate of change of its radius numerically , then  $r = \dots\dots\dots$


(a)  $\frac{1}{2\sqrt{\pi}}$  (b)  $\sqrt{2\pi}$  (c)  $\frac{1}{\sqrt{2\pi}}$  (d)  $\frac{1}{\sqrt{2}\pi}$

- 31 If (A) is the area of a circle of radius (r) , the radius changes at constant rate , then .....

(a) A is constant. (b)  $\frac{dA}{dt}$  is constant. (c)  $\frac{dA}{dt} \propto r$  (d)  $\frac{dA}{dt} \propto r^2$

- 32 In the opposite figure :

A circle drawn inside a square if the radius increases by the rate  $\frac{1}{\sqrt{2}} \text{ m./sec}$ . , then



**First :** rate of change of the perimeter of the square  $\equiv \dots\dots\dots \text{ m./sec}$ .

(a)  $\sqrt{2}$  (b) 2 (c)  $2\sqrt{2}$  (d)  $4\sqrt{2}$

**Second :** rate of change of the square area = .....  $\text{m}^2/\text{sec}$ . at  $r = \sqrt{2} \text{ m}$ .

(a)  $2\sqrt{2}$  (b) 4 (c)  $4\sqrt{2}$  (d) 8

**Third :** rate of change of the diagonal of the square = .....  $\text{m./sec}$ .

(a)  $\sqrt{2}$  (b) 2 (c)  $2\sqrt{2}$  (d) 4

**Fourth :** rate of change of the area of the shaded region = .....  $\text{m}^2/\text{sec}$ . at  $r = \sqrt{2} \text{ m}$ .

(a)  $4 - 2\pi$  (b)  $8 - \pi$  (c)  $4 - \pi$  (d)  $8 - 2\pi$

- 33 If the side length of a square decreases by a constant rate  $4 \text{ cm./min}$ . , then rate of change of area of the square = .....  $\text{cm}^2/\text{min}$ . (in terms of perimeter of the square P)

(a)  $-2P$  (b)  $-4P$  (c)  $-6P$  (d)  $-8P$



- 34 A square whose side length  $X$  cm. increases in the rate 1 cm./sec. If the perimeter of this square  $l$  cm. and is diagonal length  $z$  cm. , then  $\frac{dl}{dz} = \dots\dots\dots$


(a)  $\frac{1}{\sqrt{2}}$                       (b)  $\sqrt{2}$                       (c)  $2\sqrt{2}$                       (d)  $4\sqrt{2}$

- 35 In the opposite figure :

A right-angled triangle in which  $\frac{dz}{dt} = 3$

,  $\frac{dX}{dt} = 3 \frac{dy}{dt}$  then  $\frac{dy}{dt} = \dots\dots\dots$

at the instant when  $X = 8$  ,  $y = 6$



(a) 1                      (b) 2                      (c) 3                      (d) 4

- 36 If the base of a triangle decreases in the rate 0.2 cm./sec. and height of the triangle "h" increase in the rate 0.1 cm./sec. If area of the triangle increases when the base length = 3 cm. , then  $\dots\dots\dots$

(a)  $h > 3$                       (b)  $h > 1.5$                       (c)  $h < 1.5$                       (d)  $h < 1$

- 37 If the length of the base of a triangle (b) increases at a rate 3 cm./min. , and its corresponding height (h) decreases at a rate 3 cm./min. , then the area of the triangle (A)  $\dots\dots\dots$

(a) always increases.                      (b) remains constant.  
(c) decreases only when  $b < h$                       (d) decreases only when  $b > h$

- 38 The number of sides of a regular polygon is  $m$  and its side length increases at constant rate  $a$  cm./sec. , then  $\dots\dots\dots$

(a) its perimeter increases at rate  $a$  cm./sec.                      (b) its area increases at rate  $a$  cm<sup>2</sup>/sec.  
(c) its perimeter increases at rate  $m a$  cm./sec.                      (d) its area increases at rate  $m a$  cm<sup>2</sup>/sec.

- 39 The number of sides of a regular polygon is  $m$  and its side length increases at constant rate  $a$  cm./sec. , then the angle at any vertex of the polygon  $\dots\dots\dots$

(a) increases at constant rate  $(a)^{\text{rad.}}/\text{sec.}$   
(b) increases at constant rate  $(m a)^{\text{rad.}}/\text{sec.}$   
(c) increases at non constant rate and can not be determined.  
(d) remains constant.



- 40 A circular segment, the radius of its circle is 10 cm. and measure of its central angle ( $x^\circ$ ) changes in the rate  $3^{\text{rad}}$  per minute, then the rate of increasing in the area of the circular segment at  $x = 60^\circ$  is .....  $\text{cm}^2/\text{min}$ .
- (a) 125                      (b) 75                      (c) 150                      (d) 300
- 
- 41 If  $x + y = \text{constant}$ , then .....
- (a) each of  $x$  and  $y$  increases at the same rate.  
(b) each of  $x$  and  $y$  decreases at the same rate.  
(c) one of them increases and the other decreases at the same rate.  
(d) nothing of the previous.
- 
- 42 A spherical balloon whose radius  $r$ , its volume  $v$ , it is filled with gas but the gas leaks at constant rate, then .....
- (a)  $\frac{dr}{dt} > 0, \frac{dv}{dt} > 0$                       (b)  $\frac{dr}{dt} > 0, \frac{dv}{dt} < 0$   
(c)  $\frac{dr}{dt} < 0, \frac{dv}{dt} > 0$                       (d)  $\frac{dr}{dt} < 0, \frac{dv}{dt} < 0$
- 
- 43 A spherical balloon, its volume  $100\pi \text{ cm}^3$ , full with gass and due to leaking of gass the volume of the balloon decreases at a rate of  $8\pi \text{ cm}^3/\text{min}$ . Preserving its spherical shape Find :
- First :** The rate of change of its radius length when its radius = 4 cm. is ..... cm/min.
- (a)  $-\frac{1}{8}$                       (b)  $-\frac{1}{4}$                       (c)  $-\frac{1}{2}$                       (d) -2
- Second :** The rate of change of its radius length after 8 minutes. from the beginning of leaking = ..... cm./min.
- (a)  $-\frac{3}{8}$                       (b)  $-\frac{2}{9}$                       (c)  $-\frac{1}{9}$                       (d)  $-\frac{2}{3}$
- 
- 44 Triangular right prism its base in the shape of isosceles right-angled triangle in which length of each leg 10 m. If water is poured inside it in the rate  $30 \text{ m}^3/\text{min}$ , then rate of change of the height of water = ..... m./min.
- (a) 0.6                      (b) 1                      (c) 2.4                      (d) 0.35



45 A 10 meter ladder is leaning against a vertical wall and its lower end on a horizontal ground, if the lower end slides 2 m./min., then the rate of change of inclination angle with the horizontal at the moment the lower end at a distance 8 m. equals ..... rad/min.

- (a) 3                      (b) -3                      (c)  $\frac{1}{3}$                       (d)  $-\frac{1}{3}$


46 A 2 meter ladder leaning against a vertical smooth wall with its top and its base on horizontal smooth ground, then the distant between the base to the wall in the moment which the rate of change of sliding of the two ends are equals is ..... m.

- (a) 2                      (b)  $2\sqrt{2}$                       (c)  $\sqrt{2}$                       (d)  $-\sqrt{2}$

47 A 4 meter ladder leaning against a vertical wall with its top and its base on horizontal ground, if its base slides by velocity  $\sqrt{3}$  m./sec. when the ladder inclined with the wall by angle with measure  $60^\circ$ , then the velocity of sliding of the other end on the wall at this moment = ..... m./sec.

- (a)  $\frac{3}{2}$                       (b) 3                      (c)  $\sqrt{3}$                       (d)  $-\frac{\sqrt{3}}{3}$

48 A train moves with velocity 30 km./h. in direction of west towards the station of the train and at the same moment another train moves from the same station with velocity 30 km./h. in direction of north, then the distance between the two trains is .....



- (a) always increasing.  
(b) always decreasing.  
(c) increasing until reach a certain moment, then decreasing.  
(d) decreasing until reach a certain moment, then increasing.

49 A 1.6 meter man walks away from a lamppost at rate of 4 m./sec., the height of the lamppost is 4.8 m. from the ground, then the rate of change of the length of the man's shadow equal ..... m./sec.

- (a) 1                      (b) 2                      (c) 3                      (d) 4



- 50 A metal fine lamina in rectangular form, its length is  $\frac{4}{5}$  of its diagonal, shrinking by cooling uniformly preserving its geometrical shape and same ratio between its dimensions, at a certain moment its diagonal shrunk at a rate 2.5 cm./min. and at the same moment its surface area decreases at a rate 60 cm<sup>2</sup>/min., then the surface area of the lamina at this moment = ..... cm<sup>2</sup>.
- (a) 300 (b) 600 (c) 150 (d) 625
- 
- 51 The radius of a cylindrical tank is 25 cm. and its height 120 cm., oil is poured in it at a rate  $\frac{5000}{L+40}$  m<sup>3</sup>/sec. where L is the height of the oil at any moment then the rate of change of its height in the tank = ..... cm./sec. when its half full.
- (a)  $\frac{4}{25} \pi$  (b)  $\frac{1}{25 \pi}$  (c)  $\frac{2}{25 \pi}$  (d)  $\frac{8}{25 \pi}$
- 
- 52 A right circular cylinder expands preserving its shape, the rate of increasing of its radius is 0.5 cm./sec. and its height (h) increase at a rate of 0.25 cm./sec. then the rate of change of its volume when r = 3 cm., h = 5 cm. equals ..... cm<sup>3</sup>/sec.
- (a)  $\frac{69}{4} \pi$  (b)  $\frac{15}{4} \pi$  (c)  $\frac{13}{2} \pi$  (d)  $\frac{3}{4} \pi$
- 
- 53 An isosceles triangle, the length of each of the two equal sides is 6 cm., and the measure of their included angle equals (X) if X change at a rate of 2° per minute, then the rate of change of its surface area when X = 30° is ..... cm<sup>2</sup>/min.
- (a)  $\frac{\sqrt{3}}{10} \pi$  (b)  $\frac{\pi}{10}$  (c)  $9\sqrt{3}$  (d) 9
- 
- 54 The relation between the vertical displacement (s) and time (t) sec is given by  $s = 49t - 4.9t^2$ , then the maximum displacement can the body reach after ..... sec.
- (a) 9.8 (b) 10 (c) 5 (d) 50
- 
- 55 The slope of the tangent to the curve  $y = f(X)$  at a point equals  $\frac{1}{2}$  and the X-coordinate of this point decreases at a rate 3 unit./sec., then the rate of change of its y-coordinate equals ..... unit./sec.
- (a)  $-\frac{1}{6}$  (b)  $-\frac{3}{2}$  (c)  $\frac{1}{6}$  (d)  $\frac{3}{2}$




56 A man observes a plane flies at 3 km. high horizontally above him and with speed 480 km./h. , then the rate of change of the distance between the man and the plane after 30 sec. later = .....

- (a)  $\frac{320}{3}$  km./h. (b) 384 km./sec. (c) 384 m./sec. (d)  $\frac{320}{3}$  m./sec.

57 A right circular cone its height equals length of its base diameter if the rate of change of the radius of its base =  $\frac{1}{\pi}$  cm./sec. , then the rate of change of the volume of the cone = ..... cm.<sup>3</sup>/sec. when the radius of its base = 5 cm.


- (a)  $50\pi$  (b)  $\frac{250}{3}\pi$  (c) 150 (d) 50

58 The opposite figure represents a circular cone whose vertex is down , water leaks from it in a constant rate , then which of the following does decrease in a constant rate also ?



- (a) radius of the water surface.  
(b) height of water.  
(c) volume of water.  
(d) all the previous.


59 The opposite figure represents a frustum circular right cone , water is pouring inside it from the top , then which of the following is positive ?



- ① rate of change of the radius of the water surface.  
② rate of change of the height of the water.  
③ rate of change of the volume of the water.  
(a) ① only  
(b) ① and ② only  
(c) ② and ③ only  
(d) ① , ② and ③




- 60 The opposite figure represents a vessel in the form of a circular right cylinder, water leaks from it in a constant rate, then which of the following does decrease in a constant rate ?



- ① radius of the water surface.                      ② height of water.  
 ③ volume of water.  
 (a) ① only    (b) ①, ② only  
 (c) ②, ③ only                                        (d) ①, ② and ③

- 61 In the opposite figure :

A square whose side length increases in the rate  $\frac{1}{4}$  cm./sec. , then rate of change of the shaded areas when the side length of the square 16 cm. equals ..... cm<sup>2</sup>/sec.



- (a) 2    (b) 4    (c) 8    (d) 16


- 62 In the opposite figure :

The point A (X, y) moves on the circle whose centre is the origin and its diameter length

10 length unit and  $\frac{dX}{dt} = -2$  length unit/sec.

at the point (3, 4) , then  $\frac{dy}{dt} = \dots\dots\dots$  unit length/sec.

and  $\frac{d\theta}{dt} = (\dots\dots\dots)$  rad./sec.



- (a)  $\frac{3}{2}, \frac{1}{2}$     (b) 6, 2    (c)  $\frac{3}{2}, 2$     (d) 6,  $\frac{1}{2}$

- 63 In the opposite figure :


ABCD is isosceles trapezium drawn inside

a semicircle whose center M and

$m(\angle BAD) = 60^\circ$

If rate of change of the radius =  $\sqrt{3}$  cm./sec.

, then rate of change of area of the trapezium with respect to time = ..... cm<sup>2</sup>/sec. when its radius = 8 cm.




- (a) 36    (b)  $36\pi$     (c) 12    (d)  $12\pi$



## 64 In the opposite figure :

ABC is equilateral triangle inscribed in a circle , if the rate of change of the shaded area =  $(48\pi - 36\sqrt{3}) \text{ cm}^2/\text{sec.}$  at the moment at which the radius of the circle equals 8 cm. , then the rate of change of the radius of the circle = ..... cm./sec.



- (a)  $\sqrt{3}$  (b) 3 (c)  $\sqrt{3}\pi$  (d)  $3\pi$

65 ABC is equilateral triangle in which  $\overline{BE}$  and  $\overline{AD}$  are two medians intersects each other in the point M , if the rate of change of the median length with respect to the time = 2 cm./sec. , then rate of change of area of  $\Delta MDE$  = .....  $\text{cm}^2/\text{sec.}$  when the median length =  $18\sqrt{3}$  cm.

- (a) 3 (b) 6 (c) 18 (d)  $18\sqrt{3}$

66 If y is positive and is increasing then value of y at which rate of increasing in  $y^3$  equals 4 times the rate of increasing in y is .....

- (a)  $\frac{2}{\sqrt{3}}$  (b)  $\frac{\sqrt{3}}{2}$  (c)  $2\sqrt{3}$  (d)  $3\sqrt{2}$

67 In a right-angled triangle the length of two sides of the right angle are 12 cm. , 16 cm. if the length of the first side increases at rate 2 cm./sec. and the length of second side decreases at rate 1 cm./sec. , then the rate of change of its area after 2 seconds = .....  $\text{cm}^2/\text{sec.}$

- (a) 6 (b) 3 (c) 12 (d) 18

68 A water pipe of length 5 m. and its ends are A and B it rests with its end A on a horizontal ground and one of its points D on a vertical fence of height 3 m. If the lower end A slide away from the fence at a rate  $\frac{5}{4}$  m./min. , then the rate of descending the end B when it reaches to the fence top = ..... m./min.

- (a)  $\frac{3}{5}$  (b)  $\frac{3}{4}$  (c)  $\frac{5}{4}$  (d)  $\frac{4}{5}$

69 A car begins to move west at 12 pm. with speed 30 km./h. Another car begins at the same point at 2 pm. to travel North with speed 45 km./h. , then the rate of change of the distance between the two cars at 4 pm. is ..... km./h.

- (a) 49 (b) 51 (c) 53 (d) 55



- 70 The base of metallic cuboid is a square its side length increases at a rate of 1 cm./min. and its height decreases at a rate of 2 cm./min. , then the volume of the cuboid stop increasing after ..... min. from the moment in which the length of the base side is 5 cm. and its height 20 cm.
- (a) 5 (b) 3 (c) 12 (d) 6
- 
- 71 If the point A moves in the positive direction of X-axis starting from the origin (O) with velocity  $\frac{2}{3}$  length unit/min. and B (0 , 2) , C (0 , 4) , then the rate of change in the measure of the angle ( $\angle$  BAC) when A reaches to (2 , 0) is (.....) rad./min.
- (a)  $\frac{1}{3}$  (b)  $\frac{2}{3}$  (c)  $\frac{2}{15}$  (d)  $\frac{1}{30}$
- 
- 72 The length of rectangle 12 cm. and its width 5 cm. the length decreases at a rate 1 cm./min. while the width increases at a rate  $\frac{1}{2}$  cm./min. , then the time the area stop increasing is ..... and the area at this moment is .....
- (a) 1 sec. , 60.5 cm.<sup>2</sup> (b)  $\frac{14}{3}$  sec. , 60.5 cm.<sup>2</sup>  
(c) 1 sec. , 30 cm.<sup>2</sup> (d)  $\frac{14}{3}$  sec. , 30 cm.<sup>2</sup>
- 
- 73 A metal regular quadrilateral pyramid whose height equals its base side length. The volume of the pyramid increases at a rate of 1 cm<sup>3</sup>/sec. when the rate of increasing of both the pyramid's height and its base side length equals 0.01 cm./sec. , then its base side length = ..... cm. at this moment.
- (a) 4 (b) 8 (c) 10 (d) 12
- 
- 74 50 metre rop passes over a pulley which is 24 m. high , one of its ends tied to a heavy mass and the other end tied to a car moves on the ground with velocity 18 m./sec. then the rate of change of height of the mass at the moment when the car at a distance 32 m. from the projection of the pulley = ..... m./sec.
- (a) 7.2 (b) 14.4 (c) 18.8 (d) 21.6




- 75 A right circular conical funnel, its height is 35 cm., and the radius length of its base is 7 cm. fixed such that its axis is vertical while its apex is downward, oil is poured in it at a rate  $5.4 \text{ cm}^3/\text{sec.}$  while its leaking from a hole in the other end at a rate  $3.2 \text{ cm}^3/\text{sec.}$ , then the rate of increasing of the height of oil when the oil height is 25 cm. ( $\pi = \frac{22}{7}$ ) equal ..... cm./sec.

(a) 0.007                      (b) 0.014                      (c) 0.028                      (d) 0.056

76 In the opposite figure :

ABC is an isosceles triangle in which  $AB = AC = 13 \text{ cm.}$ ,  $BC = 10 \text{ cm.}$


If the straight line L moves from point B parallel to  $\overrightarrow{AC}$  in direction of  $\overrightarrow{BC}$  to intersects  $\overline{AB}$ ,  $\overline{BC}$  at D, E respectively where the rate of change in  $BE = \frac{1}{10} \text{ cm./sec.}$ , then the rate of change of the area of  $\Delta DEB$  at  $BE = 1 \text{ cm.}$  equals .....  $\text{cm}^2/\text{sec.}$



(a) 0.48                      (b) 0.12  
(c) 0.96                      (d) 0.24

77 In the opposite figure :

$\overline{AB}$  is tangent segment to the circle,  $\overline{AD}$  is secant to it where  $CD = 5 \text{ cm.}$  If the rate of change of the length of  $\overline{AB}$  equals  $1 \text{ cm./min.}$ , then rate of change of the length of  $\overline{AC}$  when its length 4 cm. equals ..... cm./min.




(a) 2                      (b)  $\frac{1}{2}$   
(c)  $\frac{12}{13}$                       (d)  $\frac{3}{4}$

78 In the opposite figure :

Two concentric circles, their radii are 6 cm., 18 cm.

If the radius of the smaller increases at a rate  $2 \text{ cm./sec.}$  and the radius of the greater increases at a rate  $1 \text{ cm./sec.}$  then the area between the two circles during the time interval  $]0, 12[$  .....




(a) continuously increases.                      (b) continuously decreases.  
(c) increases then decreases.                      (d) decreases then increases.



79 (2<sup>nd</sup> session 2021) In the opposite figure :


Two squares having the same center and their sides lengths are 1 cm. and 4 cm. , if side length of the first square increases by rate 1 cm./sec. and side length of the second square decreases by rate  $\frac{1}{2}$  cm./sec. , then the area of the shaded region after  $\frac{1}{2}$  second .....



- (a) stop increasing instantly. (b) stop decreasing instantly.  
(c) increases at rate  $\frac{1}{4}$  length unit/sec. (d) decreases at rate  $\frac{1}{4}$  length unit/sec.

80 (Trial 2021) In the opposite figure :


A dark barrier of height  $l$  cm. is placed at a distance 3 cm. from a burning candle and at a distance 6 cm. from a vertical wall , if the height of the candle is decreasing by the rate 3 cm./hr. , then the rate of changing of the shadow length ( $y$ ) on the wall equals ..... cm./hr.



- (a) 3 (b) -3 (c) 6 (d) -6


81 In the opposite figure :

M , N , K are three congruent circles , radius of each of them  $r$  , if  $\frac{dr}{dt} = \frac{2}{\pi}$  cm./sec. , then rate of change of area of shaded part = .....  $\text{cm}^2/\text{sec}$ . at  $r = 6$  cm.



- (a) 6 (b) 8  
(c)  $6\pi$  (d)  $8\pi$

82 (1<sup>st</sup> session 2021) ABCDHN is a regular hexagon , the length of its side increases at rate  $\sqrt{3}$  cm./sec. , when the side length is 4 cm. , then the rate of change of the shaded region with respect to time = .....  $\text{cm}^2/\text{sec}$ .




- (a) 12 (b) 8  
(c)  $12\sqrt{3}$  (d)  $8\sqrt{3}$




**83 In the opposite figure :**

ABCDEF is regular hexagon whose centre (M) if its side length increases in the rate  $\sqrt{3}$  cm./sec. , then rate of change in the shaded area = ..... cm<sup>2</sup>/sec. when its side length 4 cm.



- (a) 9                      (b) 18  
(c)  $3\sqrt{3}$               (d)  $9\sqrt{3}$


**84** The opposite figure represents the curve of the function  $y = e^{-\frac{1}{2}x^2}$  if  $\frac{dX}{dt} = e^2$  cm./min. , then rate of change of the area of shaded rectangle = ..... cm<sup>2</sup>/min. when  $X = 2$



- (a) -3  
(b) -4  
(c) -5  
(d) -6

**85 In the opposite figure :**

A particle ( $X, y$ ) is moving along the curve of the function  $y = \sqrt{x}$ . When  $X = 4$  , the  $y$ -component of the position of the particle is increasing at rate 1 length unit/sec.



**First :** The rate of change of the  $X$ -component at this moment = ..... length unit/sec.

- (a) 3                      (b) 4                      (c) 5                      (d) 6

**Second :** The rate of change of the distance from the origin to the particle at the same moment = ..... length unit/sec.

- (a)  $\frac{4\sqrt{5}}{2}$                       (b)  $\frac{5\sqrt{5}}{9}$                       (c)  $\frac{9\sqrt{5}}{5}$                       (d)  $9\sqrt{5}$



**Third :** The rate of change of the angle of inclination  $\theta$  at the same moment = ..... rad./sec.

(a)  $-\frac{1}{5}$

(b)  $-\frac{2}{5}$

(c)  $-\frac{3}{5}$

(d)  $-\frac{4}{5}$

**Fourth :** The rate of change in area of the triangle ABO at the same moment = ..... square units/sec.

(a) 1

(b) 6

(c) 3

(d) 4


- 86** A hemi - sphere tank of water , radius 2 m. , water is poured into it , if the rate of change of the height of water is  $\frac{1}{4}$  m./min. , then the rate of change of the area of the water surface in the tank after 2 min. from the beginning of the pouring water is ..... m<sup>2</sup>/min.

(a)  $\frac{1}{4} \pi$

(b)  $\frac{1}{2} \pi$

(c)  $\frac{3}{4} \pi$

(d)  $\frac{2}{3} \pi$




- 87** Two cylindrical tanks of water , the radius of the smaller is 4 m. and the radius of the bigger one is 8 m. they are being filled simultaneously at the same rate , the rate of increases of water level in the smaller tank is  $\frac{1}{2}$  m./min. , then the rate of increase of water level in the bigger tank = ..... m./min.

(a)  $\frac{1}{2}$

(b)  $\frac{1}{4}$


(c)  $\frac{1}{8}$

(d)  $\frac{1}{16}$




- 88** In the opposite figure :


Water is poured into a right cylindrical tank at a constant rate , which of the following figures represents the relation between the water level (h) in the tank and the time (t) ?




(a)




(b)



(c)



(d)





**Fifth Questions on behavior of the function****1 Choose the correct answer from the given ones :**

- 1 The 1<sup>st</sup> derivative of the decreasing functions is .....  
 (a) positive (b) negative (c) zero (d) otherwise
- 2 The function  $f : f(x) = -x^2$  is increasing on the interval .....  
 (a)  $]0, \infty[$  (b)  $] -\infty, 0[$  (c)  $\mathbb{R} - \{0\}$  (d)  $\mathbb{R}$
- 3 The function  $f : f(x) = -|x| + 1$  is decreasing on the interval .....  
 (a)  $]0, \infty[$  (b)  $] -\infty, 0[$  (c)  $]1, \infty[$  (d)  $] -\infty, 1[$
- 4 The function  $f : f(x) = x^3 + 4x + 2$  is increasing when  $x \in$  .....  
 (a)  $] -4, \infty[$  (b)  $\mathbb{R}$  (c)  $] -\infty, \frac{-4}{3}[$  (d)  $] \frac{-4}{3}, \infty[$
- 5 The function  $f : f(x) = \frac{x}{\ln x}$  is increasing on the interval .....  
 (a)  $]0, \infty[$  (b)  $]0, e[$  (c)  $]e, \infty[$  (d) otherwise.
- 6 If  $f : ] -2, 4[ \rightarrow \mathbb{R}$ ,  $f(x) = x^3 - 3x$ , then the number of the critical points of the function  $f$  equals .....  
 (a) 1 (b) 2 (c) 3 (d) 4
- 7 The function  $f : f(x) = \ln(x^2 - 4)$ , then the number of the critical points = .....  
 (a) zero (b) 1 (c) 2 (d) 3
- 8 If  $f : f(x) = ax^2 + bx + 2$  has a critical point  $(1, 4)$ , then  $a - b =$  .....  
 (a) 2 (b) zero (c) -6 (d) -8
- 9 All the following functions are increasing in their domain except the function  $f : f(x) =$  .....  
 (a)  $2x - 17$  (b)  $e^x$  (c)  $(x - 3)^2$  (d)  $x^3$



- 10 The function  $f : f(x) = (x - 3)^2 + 2$  is increasing in the interval .....  
 (a)  $]2, \infty[$  (b)  $]3, \infty[$  (c)  $] - \infty, 3[$  (d)  $]0, \infty[$
- 
- 11 The function  $f : f(x) = x^3 - 12x$  is decreasing in the interval .....  
 (a)  $[-2, 2]$  (b)  $] - 2, 2[$  (c)  $]0, 12[$  (d)  $]4, \infty[$
- 
- 12 If  $f$  is a continuous function on  $\mathbb{R}$  and the function  $f$  has a critical point at  $x = a$ , so .....  
 (a)  $f'(a) = 0$  (b)  $f'(a)$  is undefined.  
 (c)  $f'(a^+) \neq f'(a^-)$  (d) All the previous.
- 
- 13 If the function  $f : f(x) = 2ax^2 + bx + 3$  has a local extrema at  $(1, 2)$ , then  $a + b =$  .....  
 (a)  $-1$  (b)  $\frac{5}{2}$  (c)  $-\frac{3}{2}$  (d)  $\frac{3}{2}$
- 
- 14 (2<sup>nd</sup> session 2021) If the function  $f(x) = 3ax^3 - bx - 5$  has local maximum value at  $x = 1$ , then  $\frac{b}{a} =$  .....  
 (a)  $-9$  (b)  $9$  (c)  $20$  (d)  $-20$
- 
- 15 (Trial 2021) If  $f(x) = \cot x - m \csc x$  has local maximum value at  $x = \frac{\pi}{4}$ , then the value of the constant  $m =$  .....  
 (a)  $-\sqrt{2}$  (b)  $-1$  (c)  $1$  (d)  $\sqrt{2}$
- 
- 16 If  $f$  is an odd continuous function on  $\mathbb{R}$  and the function has a local minimum value at  $x = 2$ , then the function has .....  
 (a) a local maximum value at  $x = -2$  (b) a local minimum value at  $x = -2$   
 (c)  $f'(2) > f'(-2)$  (d)  $f'(2) < f'(-2)$
- 
- 17 If the function  $f : f(x) = x + \frac{a}{x}$  has local maximum at  $x = -2$ , then  $a =$  .....  
 (a)  $4$  (b)  $2$  (c)  $-2$  (d)  $-4$
- 
- 18 Let  $f$  be a function defined by :  $f(x) = \frac{x}{\ln x}$ , then the local minimum value of  $f$  is .....  
 (a)  $e$  (b)  $\frac{1}{e}$  (c)  $\ln e$  (d)  $-e$



- 19 If  $f$  is a function where  $f(x) = \frac{x^4 + 1}{x^2}$ , then the function is decreasing in .....
- (a)  $]-\infty, -1[$  only. (b)  $]0, 1[$  only.  
 (c)  $]-1, 0[$ ,  $]1, \infty[$  (d)  $]-\infty, -1[$ ,  $]0, 1[$
- 
- 20 If  $f$  is function where  $f(x) = (x^2 - 4)^{\frac{2}{3}}$ , then the function is decreasing on .....
- (a)  $]-\infty, -2[$ ,  $]0, 2[$  (b)  $]-2, 0[$ ,  $]2, \infty[$   
 (c)  $]-\infty, -2[$  only. (d)  $]0, 2[$  only.
- 
- 21 If  $f(x) = x(a - \ln x)$  where  $a$  is constant and the curve of the function has a critical point at  $x = e$ , then  $a =$  .....
- (a) 1 (b) zero (c)  $e$  (d) 2
- 
- 22 If  $f(x) = x \ln x$ , then the function  $f : f(x)$  has a critical point at  $x =$  .....
- (a) zero (b) 1 (c)  $e$  (d)  $\frac{1}{e}$
- 
- 23 The function  $f : f(x) = e^{x^2 - 2x}$  is increasing in the interval .....
- (a)  $]-\infty, 1[$  (b)  $]-\infty, 2[$  (c)  $]2, \infty[$  (d)  $]1, \infty[$
- 
- 24 If the function  $f$  is a polynomial function of fourth degree and its domain is  $\mathbb{R}$ , then the greatest number of the critical points of the function  $f(x)$  is .....
- (a) 1 (b) 2 (c) 3 (d) 4
- 
- 25 If the curve of the function  $f$ , where  $f$  is a polynomial, has a local maximum value at the point  $(a, b)$ , then  $f'(a) =$  .....
- (a)  $b$  (b) zero (c)  $-\frac{b}{a}$  (d) undefined
- 
- 26 Which of the following functions has local minimum value?  $f : f(x) =$  .....
- (a)  $-x^2$  (b)  $x^2 + 2$  (c)  $-x^3$  (d)  $x^3 + 2$
- 
- 27 If the function  $f$  has local maximum value, then  $f(x)$  may be equal .....
- (a)  $x^2 - 3$  (b)  $x^3 + 1$  (c)  $x^3 + 3x$  (d)  $x^4 - 3x^2$



- 28 (2<sup>nd</sup> session 2021) If the function  $f(x) = x^2 + \frac{2a^3}{x}$  where  $a \in \mathbb{R}^-$ , then the function is decreasing in the interval .....
- (a)  $]-\infty, 0[$ ,  $]0, \infty[$  (b)  $]a, \infty[$   
 (c)  $]-\infty, a[$  (d)  $]-\infty, 0[$ ,  $]a, 0[$
- 
- 29 If the function  $f : f(x) = \frac{x}{x^2 + 1}$  is increasing on  $]a, b[$ , then  $b - a = \dots\dots\dots$
- (a) zero (b) 1 (c) 2 (d) 3
- 
- 30 The maximum value of the function  $f : f(x) = 3 - \sin x$  is ..... where  $x \in \mathbb{R}$
- (a) 1 (b) 2 (c) 3 (d) 4
- 
- 31 The function  $f : f(x) = kx - \cos x$  is increasing on its domain, then .....
- (a)  $k \geq 1$  (b)  $k \leq 1$  (c)  $-1 < k < 1$  (d)  $k = \text{zero}$
- 
- 32 (Trial 2021) If the function  $f : [-3, -1] \rightarrow \mathbb{R}$  where  $f(x) = x + \frac{a}{x}$  and the absolute maximum value of  $f$  equals  $-2$  and the function is increasing on  $]-3, -1[$ , then  $a = \dots\dots\dots$
- (a)  $-2$  (b)  $-1$  (c)  $1$  (d)  $2$
- 
- 33 The function  $f$  where  $f(x) = (x - a)^2 + (x - b)^2$  has a local minimum value at  $x = \dots\dots\dots$
- (a)  $\frac{a+b}{2}$  (b)  $\sqrt{ab}$  (c)  $\frac{2}{a+b}$  (d)  $\frac{2ab}{a+b}$
- 
- 34 If the function  $f : f(x) = x^2 + \frac{b}{x}$  has a critical point at  $x = 2$ , then  $b = \dots\dots\dots$
- (a)  $-16$  (b)  $16$  (c)  $-4$  (d)  $2$
- 
- 35 If the function  $f$  is continuous on the interval  $]a, b[$ , if  $c \in ]a, b[$  where  $\tilde{f}(c^+) \neq \tilde{f}(c^-)$ , then  $(c, f(c))$  is called .....
- (a) maximum. (b) minimum. (c) critical. (d) inflection.



- 36 If  $f : f(x) = \sqrt[3]{x-c}$  has critical point at  $(c, 0)$ , then  $f'(c) = \dots\dots\dots$
- (a) undefined.                      (b) zero.                      (c)  $\frac{1}{3}$                       (d)  $\frac{1}{3\sqrt[3]{c^2}}$
- 
- 37 If the curve of the function  $f$  has  $f(5) = 7$ ,  $f'(5) = \text{zero}$ ,  $f''(5) = -4$ , then the point  $(5, 7)$  has  $\dots\dots\dots$
- (a) a local maximum.                      (b) a local minimum.  
(c) undefined.                      (d) an inflection point.
- 
- 38 If the function  $f$  is differentiable and  $f'(x_1) = 0$ ,  $f''(x_1) > 0$ , then  $\dots\dots\dots$
- (a) at  $x = x_1$  the function has maximum point.  
(b) at  $x = x_1$  the function has minimum point.  
(c) the function increases on its domain.  
(d) the function decreases on its domain.
- 
- 39 If  $x \in ]0, \frac{\pi}{2}[$ ,  $f(x) = \sin(x + \frac{\pi}{6}) + \cos(x + \frac{\pi}{6})$  has local maximum value at  $x = \dots\dots\dots$
- (a)  $\frac{\pi}{4}$                       (b)  $\frac{\pi}{6}$                       (c)  $\frac{\pi}{12}$                       (d)  $\frac{\pi}{8}$
- 
- 40 If  $f'(x) = (x-m)(x-k)$  where  $m < k$ , then  $\dots\dots\dots$
- (a)  $f'(m) = f'(k)$   
(b)  $f''(m) < f''(k)$   
(c) The function  $f$  has an inflection point at  $x = \frac{m+k}{2}$   
(d) all the previous correct
- 
- 41 If  $f$  is an even function for all values of  $x \in \mathbb{R}$ , and it has local maximum value at  $x = a$ , then its certainly that  $\dots\dots\dots$
- (a)  $f$  has local maximum value at  $x = -a$   
(b)  $f$  has absolute minimum value at  $x = a$   
(c)  $f'(a) > f'(a)$   
(d)  $f'(a) < f'(a)$



- 42 If  $f$  is a continuous and decreasing function on the interval  $]0, 10[$  and has critical point at  $(4, 2)$ , then which of the following has to be false?
- (a)  $f(4)$  is neither local minimum value nor local maximum value.  
 (b)  $f'(4)$  does not exist.  
 (c)  $f'(4) = 0$   
 (d)  $f'(4) < 0$
- 
- 43 If  $f''(x) = (x - 3)(x + 2)$ , then curve of the function  $f$  is convex upwards at the interval .....
- (a)  $]-\infty, -2[$                       (b)  $]-2, 3[$                       (c)  $]3, \infty[$                       (d)  $]-\infty, 3[$
- 
- 44 The curve of the function  $f$  is convex upwards on a certain interval if ..... on this interval.
- (a)  $f'(x) > 0$                       (b)  $f'(x) < 0$                       (c)  $f''(x) > 0$                       (d)  $f''(x) < 0$
- 
- 45 If  $f'(-1) = f'(3) = 0$  and  $f''(x) > 0$  for all  $x \in ]-2, 2[$ , then .....
- (a)  $f(-1)$  is a local maximum value.                      (b)  $f(-1)$  is a local minimum value.  
 (c)  $f(3)$  is a local maximum value.                      (d)  $f(3)$  is a local minimum value.
- 
- 46 The function  $f : f(x) = x^4 - 4x^2$  has .....
- (a) one local minimum value and two local maximum values.  
 (b) two different local minimum values and one local maximum value.  
 (c) two local minimum values and no local maximum values.  
 (d) two equal local minimum values and one local maximum value.
- 
- 47 If  $f''(x) = (x - 1)(x - 3)$ , then the function  $f$  is decreasing in the interval .....
- (a)  $]1, 3[$                       (b)  $]2, \infty[$                       (c)  $]-\infty, 2[$                       (d)  $\mathbb{R} - [1, 3]$
- 
- 48 If the function  $f : f(x) = 2 - ax^3$  is decreasing on its domain, then .....
- (a)  $a \leq 0$                       (b)  $a > 0$                       (c)  $a \geq 0$                       (d)  $a < 0$



- 49 If  $g$  is increasing function on  $\mathbb{R}$ ,  $h$  is decreasing function on  $\mathbb{R}$  and  $f(x) = 4g(x) - 3h(x)$ , then function  $f$  is ..... on  $\mathbb{R}$
- (a) increasing (b) decreasing (c) constant (d) zero
- 
- 50  $f, g$  are two increasing functions on  $\mathbb{R}$ , which of the following is increasing on its domain?
- (a)  $f + g$  (b)  $f - g$  (c)  $f \cdot g$  (d)  $\frac{f}{g}$
- 
- 51 If the function  $f$  such that  $f'(x) > 0$  on an interval, then the curve of the function is ..... on this interval
- (a) increasing (b) convex upwards  
(c) convex downwards (d) decreasing
- 
- 52 The curve of the function  $f$  is convex downwards on  $\mathbb{R}$  if  $f(x)$  equals .....
- (a)  $3 - x^2$  (b)  $3 - x^3$  (c)  $3 - x^4$  (d)  $3 + x^4$
- 
- 53 The curve of the function  $f$  where  $f(x) = x^3 - 3x^2 + 2$  is convex upwards when  $x \in$  .....
- (a)  $]-\infty, 0[$  (b)  $]-\infty, 1[$  (c)  $]1, 3[$  (d)  $]1, \infty[$
- 
- 54 If  $f(x) = (a - 2)x^2 + 3x - 5$ ,  $x \in \mathbb{R}$ , then the curve of the function  $f$  is concave downward when .....
- (a)  $a > 2$  (b)  $a < 2$  (c)  $a = 2$  (d)  $a = 0$
- 
- 55 The curve of the function  $f$  where  $f(x) = (x - 2)e^x$  is convex downwards on the interval .....
- (a)  $]-\infty, \infty[$  (b)  $]-1, 2[$  (c)  $]0, 2[$  (d)  $]0, \infty[$
- 
- 56 The curve of the function  $f$  where  $f(x) = \begin{cases} x^3 + 2 & , x > 0 \\ 3 - 2x^2 & , x < 0 \end{cases}$  is convex upwards when  $x \in$  .....
- (a)  $]0, \infty[$  (b)  $]-\infty, 0[$  (c)  $\mathbb{R}$  (d)  $\mathbb{R}^*$



- 57 If the curve of the function  $f$  lies above the tangents which drawn from the points on its curve, then the curve of the function is .....  
(a) convex upwards. (b) decreasing.  
(c) increasing. (d) convex downwards.
- 58 If the function  $f$  has  $\lim_{h \rightarrow 0} \frac{f'(x+h) - f'(x)}{h} > 0$  for all values of  $x \in \mathbb{R}$ , then the curve of the function  $f$  is .....  
(a) convex upwards. (b) increasing.  
(c) decreasing. (d) convex downwards.
- 59 If the curve  $y = (2x - c)^3 + 4$  has an inflection point at  $x = 5$ , then  $c =$  .....  
(a) 2 (b) 4 (c) 5 (d) 10
- 60 The function  $f : f(x) = x^3 - 3x - 1$  has an inflection point at .....  
(a) (0, 1) (b) (0, -1) (c) (1, 0) (d) (-1, 0)
- 61 If the point (1, 2) is an inflection point of the curve of the function  $f$  where  $f'(x) = 4x^3 - kx^2$ , then  $k =$  .....  
(a) 24 (b) 6 (c) 4 (d) 12
- 62 The critical point of the function  $f : f(x) = x^3 + 60$  is .....  
(a) an inflection point. (b) a local maximum.  
(c) a local minimum. (d) both (a) and (c)
- 63 If the curve :  $y = x^3 + ax^2 + bx$  has an inflection point at (3, -9), then  $a + b =$  .....  
(a) 6 (b) -9 (c) 15 (d) 24
- 64 The points which separate the convexity up and down parts are called .....  
(a) critical. (b) local minimum.  
(c) local maximum. (d) inflection.



- 65 If the function  $f : f(x) = x^4$ , then the point  $(0, 0)$  is .....
- (a) local minimum. (b) local maximum.  
(c) inflection point. (d) a, c together.
- 
- 66 If the curve of the continuous function has an inflection point, then the maximum number of the points which the curve intersects any straight line equals .....
- (a) 1 (b) 2 (c) 3 (d) 4
- 
- 67 If the function  $f$  is of fourth degree, then the maximum number of inflection points of it is .....
- (a) 2 (b) 1 (c) 3 (d) 4
- 
- 68 If  $f(x) = \sqrt[3]{x-2}$  and if  $f(x)$  has an inflection point at  $(2, 0)$ , then  $f''(2) = \dots\dots\dots$
- (a) undefined (b) zero (c) 1 (d)  $-\frac{2}{9}$
- 
- 69 If the function  $f : f(x) = g(x) - h(x)$  such that :  $g'(2) = h'(2)$  ,  $g''(2) > h''(2)$  , then at  $x = 2$   $f(x)$  has .....
- (a) local minimum value. (b) local maximum value.  
(c) inflection point. (d) absolute maximum value.
- 
- 70 If the function  $f$  is defined on the interval  $[a, b]$  and  $f'(x) < 0$  on the same interval, then the absolute maximum value of  $f(x)$  on this interval = .....
- (a)  $f(a)$  (b)  $f(b)$  (c)  $a$  (d)  $b$
- 
- 71 For the function  $f$  such that  $f'(x) = -2x + 6$ , then all of the following statements are true except .....
- (a) the curve of the function  $f$  convex upwards in the interval  $]-\infty, \infty[$   
(b) the function  $f$  has a local minimum value at  $x = 3$   
(c) the curve of the function  $f$  has no inflection points.  
(d)  $f(x)$  is decreasing in the interval  $]3, \infty[$
- 
- 72 The curve  $y = xe^x$  has at .....
- (a)  $x = -1$  a minimum value. (b)  $x = -1$  a maximum value.  
(c)  $x = 0$  a minimum value. (d)  $x = 0$  a maximum value.



- 73 The curve  $y = e^x - x e^x$  has a local .....
- (a) maximum point at  $x = 1$  (b) minimum point at  $x = 1$   
(c) maximum point at  $x = 0$  (d) minimum point at  $x = 0$
- 
- 74 The absolute maximum value of the function  $f$  where  $f(x) = 10 x e^{-x}$ ,  $x \in [\text{zero}, 4]$  is .....
- (a)  $\frac{10}{e}$  (b) zero (c) 1 (d)  $e$
- 
- 75 The function  $f: \mathbb{R} \rightarrow \mathbb{R}$  where  $f(x) = 2x + \cos x$ , then  $f$  .....
- (a) has minimum value at  $x = \pi$  (b) has maximum value at  $x = 0$   
(c) is a decreasing function. (d) is an increasing function.
- 
- 76 If  $x \in [0, \pi]$ , then the function  $f: f(x) = x \sin x + \cos x$  has absolute minimum value at  $x = \dots$
- (a) zero (b) 1 (c)  $\pi$  (d)  $\frac{\pi}{2}$
- 
- 77 (2<sup>nd</sup> session 2021) The absolute maximum value of the function  $f(x) = x \ln x$  where  $x \in [e^{-2}, e]$  equals .....
- (a)  $-2e^{-2}$  (b)  $e$  (c)  $2e$  (d)  $-e^{-1}$
- 
- 78 If the curve of the function  $f$  lies in the third quadrant and the function is decreasing on  $]a, b[$ , then the function  $g(x) = e^x \times f(x)$  is ..... in the interval  $]a, b[$
- (a) increasing (b) decreasing  
(c) concave upward (d) concave downward
- 
- 79 If  $k$  is a differentiable function on  $\mathbb{R}$  and its range  $\mathbb{R}^-$ , if  $\tilde{f}(x) = (x^2 - 4) \times k(x)$ , then the function  $f$  has .....
- (a) local maximum value at  $x = -2$  and local minimum value at  $x = 2$   
(b) local minimum value at  $x = -2$  and local maximum value at  $x = 2$   
(c) local maximum value at each of  $x = 2, x = -2$   
(d) local minimum value at each of  $x = 2, x = -2$



80 If the curve of the function  $f$  has a local minimum value at  $X = a$  and  $f(X) < 0$  for all values of  $X$ , then which of the following graphs has local minimum value at  $X = a$ ?

- (a)  $f(X) + k$                       (b)  $f(X + k)$                       (c)  $-f(X)$                       (d)  $[f(X)]^2$

81 If  $f$  is increasing function on its domain, then which of the following functions is decreasing on its domain?

- (a)  $g(X) = f(X - 2)$                       (b)  $g(X) = f(X) + 3$   
(c)  $g(X) = -f(X)$                       (d)  $g(X) = |f(X)|$

82 If  $f(X) = (X - 1)(X - 2)^2$ , then curve of the function  $f$  has an inflection point at  $X = \dots\dots\dots$

- (a) 1                      (b) 2  
(c) 1 or 2                      (d) has no inflection point

83 If the curve of the function  $f: f(X) = X^3 + 3X^2 + kX + 4$  has horizontal tangent and inflection point at the same point, then  $k = \dots\dots\dots$

- (a) zero                      (b) -1                      (c) 1                      (d) 3

84 Equation of tangent to the curve  $f(X) = X^3 + 3X^2$  at the point of inflection is  $\dots\dots\dots$

- (a)  $y = -3X - 1$                       (b)  $y = 3X - 1$                       (c)  $y = -3X + 1$                       (d)  $y = 3X + 1$

85 If the function  $f: f(X) = e^X (X^2 - aX + 9)$  has a critical point at  $X = 1$ , then  $\dots\dots\dots$

- (a) the function has local maximum value at  $X = 3$   
(b) the function has local minimum value at  $X = 3$   
(c) the function is increasing at  $X > 1$   
(d) the function is decreasing at  $X < 3$

86 If  $f: ]0, \infty[ \longrightarrow ]0, \infty[$  where

$$f(X) = \frac{\ln X}{X}, \text{ then } \dots\dots\dots$$

- (a)  $f$  is decreasing for each  $X > 1$  and curve of  $f(X)$  convex downward for each  $X > e$   
(b)  $f$  is decreasing for each  $X > 1$  and curve of  $f(X)$  convex upward for each  $X > e$   
(c)  $f$  is increasing for each  $X > 1$  and curve of  $f(X)$  convex downward for each  $X > e$   
(d)  $f$  is increasing for each  $X > 1$  and curve of  $f(X)$  convex upward for each  $X > e$



- 87 (1<sup>st</sup> session 2021)  $f, g$  are polynomial functions,  $f(x) = cx^2 + g(x)$ ,  $g(1) = k$  and  $g'(1) = 6$  where  $c, k$  are constants. If  $(1, 5)$  is an inflection point to the curve of  $f$ , then  $k - c = \dots\dots\dots$
- (a)  $-11$  (b)  $5$  (c)  $11$  (d)  $-5$
- 
- 88 (Trial 2021) If  $f$  is differentiable twice function in the interval  $[-1, 1]$  where  $\hat{f}(x)$  is increasing on  $]-1, 0[$  and  $\hat{f}(x)$  is decreasing on  $]0, 1[$ , then the statement which is certainly correct from the following is  $\dots\dots\dots$
- (a)  $f(0)$  is a local maximum value of the function  $f$   
 (b) the point  $(0, f(0))$  is an inflection point of the function  $f$   
 (c) the function  $f$  is increasing on  $]0, 1[$   
 (d) the function  $f$  is decreasing on  $]0, 1[$
- 
- 89 If  $f(x) = \ln(xe^x)$ , then  $\dots\dots\dots$
- (a) each of  $f(x), \hat{f}(x)$  is increasing on its domain.  
 (b) each of  $f(x), \hat{f}(x)$  is decreasing on its domain.  
 (c)  $f(x)$  is increasing on its domain while  $\hat{f}(x)$  is decreasing.  
 (d)  $f(x)$  is decreasing on its domain while  $\hat{f}(x)$  is increasing.
- 
- 90 The function  $f: f(x) = x^x$  has a stationary point at  $x = \dots\dots\dots$
- (a)  $e$  (b)  $\frac{1}{e}$  (c)  $1$  (d)  $\sqrt{e}$
- 
- 91 If  $f$  is a continuous function in the interval  $[a, b]$  and for every  $x_1, x_2 \in [a, b]$ , then  $\hat{f}(x_2) - \hat{f}(x_1) > 0$  when  $x_2 > x_1$ , then  $\dots\dots\dots$  in the interval  $]a, b[$
- (a) the function  $f$  increases (b) the function  $f$  decreases  
 (c) the curve of  $f$  convex upward (d) the curve of  $f$  convex downward
- 
- 92 If the function  $f$  is defined in  $]3, 8[$  where  $f(x_1) > f(x_2)$  for every  $x_2 > x_1$  and the function  $f$  is defined on  $]-3, 2[$  where  $f(x_1) - f(x_2) > 0$  for every  $x_1 > x_2$ , then all the following statements are true except  $\dots\dots\dots$
- (a)  $\hat{f}(4) < \hat{f}(-2)$  (b)  $\frac{f(5) - f(7)}{\hat{f}(0)} > 0$   
 (c)  $\hat{f}(6) > \hat{f}(-1)$  (d)  $\frac{f(6) - f(4)}{\hat{f}(-1)} < 0$



93 If  $f, g$  are two differentiable twice functions on  $\mathbb{R}$  and  $f''(x) < g''(x)$  for all values of  $x$  and  $h(x) = f(x) - g(x)$ , then  $h$  .....

- (a) increases on its domain. (b) decreases on its domain.  
(c) has a curve convex downwards. (d) has a curve convex upwards.

94 If  $f$  is an even function and continuous on  $\mathbb{R}$  and the function has an inflection point at  $x = -a$ , then the sign of  $f''(a^-) \times f''(a^+)$  is the same as .....

- (a)  $f''(a^-) \times f''(a^+)$  (b)  $f''(a^-) \times f''(a^+)$   
(c)  $f''(-a)$  (d)  $f''(-a^-) \times f''(a^-)$

95 Let  $f$  be an increasing function on its domain, which of the following functions not necessary to be increasing on its domain?

- (a)  $y = \sqrt{f(x)}$  (b)  $y = \sqrt[3]{f(x)}$  (c)  $y = [f(x)]^2$  (d)  $y = e^{f(x)}$

96 If  $f: \mathbb{R}^* \rightarrow \mathbb{R}$  where  $f(x) = x + \frac{1}{x}$  and the function  $f$  has a local maximum value at  $x = a$  and a local minimum value at  $x = b$ , then .....

- (a)  $f(a) > f(b)$  (b)  $f(a) < f(b)$  (c)  $a > b$  (d)  $f'(a) < f'(b)$

97 If  $y = ax + b$  is a tangent to the curve of the function  $f$  at any point on it, and  $f(x) \leq ax + b$ , then  $f(x)$  ..... for all values of  $x$

- (a) increases (b) decreases  
(c) has a curve convex downwards (d) has a curve convex upwards

98 If  $f: f(x) = \cos x$  where  $x \in \left[-\frac{\pi}{2}, \frac{5\pi}{2}\right]$  has an absolute maximum value, then the number of times it reaches to the maximum values is .....

- (a) 1 time (b) 2 times (c) 3 times (d) 4 times

99 If  $f$  is a differentiable function on  $\mathbb{R}$  such that  $f'(x) < 0$  for all values of  $x \in \mathbb{R}$ , then .....

- (a)  $f(x) > f(x-1)$  (b)  $f(x) < f(x+1)$   
(c)  $f(x) < f(x-1)$  (d)  $f(x) + f(x+1) = 1$



- 100 If  $f(x) = x^3 - 6x^2 + 9x + a$  has critical point at  $x_1, x_2$  where  $x_1 < x_2$  and  $f(2), f(x_1), f(x_2)$  are in geometric sequence, then  $a = \dots\dots\dots$
- (a)  $-\frac{8}{3}$  (b)  $\frac{8}{3}$  (c)  $\frac{4}{3}$  (d)  $-\frac{4}{3}$
- 
- 101 If  $f(x) = 1 + \frac{1}{x} + \frac{1}{x^2} + \frac{1}{x^3} + \dots\dots\dots \rightarrow \infty$  where  $|x| > 1$ , then curve of the function  $f$  is convex downward on the interval  $\dots\dots\dots$
- (a)  $]-1, 1[$  (b)  $]1, \infty[$  (c)  $]-\infty, -1[$  (d)  $]-1, \infty[$
- 
- 102 If  $f$  is a polynomial function and  $f'(x) = ax^2 + bx + c$ , then  $f$  is decreasing in its domain if  $\dots\dots\dots$
- (a)  $a > 0, b^2 - 4ac \leq 0$  (b)  $a > 0, b^2 - 4ac \geq 0$   
 (c)  $a < 0, b^2 - 4ac \leq 0$  (d)  $a < 0, b^2 - 4ac \geq 0$
- 
- 103 If  $f$  is a positive increasing functions,  $g$  is a positive decreasing functions and  $z(x) = \frac{f(x)}{g(x)}$ , then  $z$  is  $\dots\dots\dots$
- (a) negative and increasing. (b) negative and decreasing.  
 (c) positive and increasing. (d) positive and decreasing.
- 
- 104 If  $f: \mathbb{R} \rightarrow \mathbb{R}$  where  $f(x) = x^3 + 3x^2 - 9x$  and  $a, b$  are the absolute minimum and maximum values of function  $f$  on the interval  $[-2, 2]$ , then  $b - a = \dots\dots\dots$
- (a) 17 (b) -17 (c) 27 (d) -27
- 
- 105 If  $f: \mathbb{R} \rightarrow \mathbb{R}$  where  $f(x) = \sqrt[3]{x^2}(3x - 7)$  is increasing for every  $x \in ]-\infty, a[$ ,  $x \in ]b, \infty[$ , then  $7a + 15b = \dots\dots\dots$
- (a) 7 (b) 14 (c) 21 (d) 22
- 
- 106 If function  $f: f(x) = \frac{x^2 + mx + 4}{x - 1}$  is an increasing function, where  $m$  is a constant, then  $m \in \dots\dots\dots$
- (a)  $]-\infty, -5]$  (b)  $[-5, \infty[$  (c)  $]-5, 0]$  (d)  $]-5, 0[$
- 
- 107 If  $f: \mathbb{R} \rightarrow \mathbb{R}$  where  $f(x) = x^3 + ax^2 + 12x + 1$ , then the set of values of  $a$  which makes the function  $f$  has no critical points is  $\dots\dots\dots$
- (a)  $-9 < a < 9$  (b)  $-6 < a < 6$  (c)  $-3 < a < 3$  (d)  $0 < a < 9$



**2** By using the opposite table where  $f$  is a polynomial :

Choose the correct answer from the given ones :

$x$	-1	1	2	3	4
$f(x)$	24	0	-3	0	9
$f'(x)$	-18	-6	0	6	12

**1**  $f$  has local maximum value of  $x = \dots\dots\dots$

- (a) -1                      (b) 1                      (c) 2                      (d) 3

**2**  $f$  has local minimum value at  $x = \dots\dots\dots$

- (a) 1                      (b) 2                      (c) 3                      (d) 4

**3**  $f$  is decreasing when  $x \in \dots\dots\dots$

- (a)  $]-\infty, 1[$                       (b)  $]1, 3[$                       (c)  $]3, \infty[$                       (d)  $\mathbb{R}$

**4** The curve of  $f(x)$  is convex upwards when  $x \in \dots\dots\dots$

- (a)  $]-\infty, -1[$                       (b)  $]-\infty, 2[$                       (c)  $]3, \infty[$                       (d)  $]1, \infty[$

**5** The curve  $f(x)$  has inflection point at  $x = \dots\dots\dots$

- (a) -1                      (b) 1                      (c) 2                      (d) 3

**6** If  $g(x) = f(x) + 5$ , then  $g$  has local maximum value when  $x = \dots\dots\dots$

- (a) -1                      (b) 1                      (c) 2                      (d) 3

### Sixth Questions on applications of maxima and minima

Choose the correct answer from the given ones :

**1** The greatest value of the expression  $8x - x^2$  where  $x \in \mathbb{R}$  is  $\dots\dots\dots$

- (a) 8                      (b) 12                      (c) 16                      (d) 20

**2** If  $x \in \mathbb{R}^+$  and  $x + \frac{1}{x} \geq k$ , then  $k = \dots\dots\dots$

- (a) 2                      (b) 4                      (c) 3                      (d) 1

**3** If  $x < 0$ , then the maximum value of  $x + \frac{1}{x}$  equals  $\dots\dots\dots$

- (a) -1                      (b) -2                      (c) -3                      (d) otherwise.

**4** The maximum value of  $(\sin x + \sqrt{3} \cos x)$  at  $x = \dots\dots\dots$

- (a)  $\frac{\pi}{3}$                       (b)  $\frac{\pi}{4}$                       (c)  $\frac{\pi}{6}$                       (d) zero



- 5 If  $x + y = 13$  where  $x$  and  $y$  are positive numbers, then the value of  $x$  which makes the expression  $(2x + 3y + xy)$  has a maximum value is .....  
(a) 2 (b) 4 (c) 6 (d) 8
- 6 If the sum of two numbers is 16 and the sum of their squares is as minimum as possible, then the two numbers are .....  
(a) 8, 8 (b) 7, 9 (c) 6, 10 (d) 5, 11
- 7 The sum of two positive integers is 5 and the sum of cube of the smaller and twice the square of the other is the smallest value, then the two numbers are .....  
(a) 4, 1 (b) 2, 3 (c) 5, zero (d)  $3\frac{1}{2}, 1\frac{1}{2}$
- 8 A rectangle its area is  $50 \text{ m}^2$ , then its perimeter has a minimum value when its dimensions are ..... meters.  
(a) 10, 5 (b)  $5\sqrt{2}, 5\sqrt{2}$  (c)  $15, \frac{10}{3}$  (d) otherwise
- 9 A rectangle with perimeter 14 cm., then the maximum area of the rectangle = .....  $\text{cm}^2$ .  
(a) 10 (b) 12 (c) 12.25 (d) 49
- 10 The rectangles which inscribed in a circle its radius " $r$ ", then the dimensions of the rectangle when its area is maximum are .....  
(a)  $\sqrt{2}r, \sqrt{2}r$  (b)  $r, \sqrt{2}r$  (c)  $2r, \sqrt{2}r$  (d)  $2r, 2r$
- 11 If the perimeter of the circular sector is constant ( $P$ ), then its area has maximum value at  $r =$  .....  
(a)  $\frac{P}{2}$  (b)  $\frac{2}{\sqrt{P}}$  (c)  $\sqrt{P}$  (d)  $\frac{P}{4}$
- 12 A circular sector-piece of metal whose area  $16 \text{ cm}^2$ , then the radius of the sector circle which makes its primeter as minimum as possible = ..... cm.  
(a) 2 (b) 4 (c) 6 (d) 8
- 13 A rectangular piece of land, bounded by a river from one side and of area  $2048 \text{ m}^2$ , such that the length of a fence surrounding its three remaining sides is minimum, then the dimensions are ..... m.  
(a) 16, 128 (b) 32, 64 (c) 8, 256 (d) 4, 512



- 14 The points on the curve  $x^2 - y^2 = 8$  such that their distance from the point  $(0, 2)$  is minimum, then the points are .....
- (a)  $(3, 1), (-3, 1)$  (b)  $(3, -1), (-3, -1)$   
 (c)  $(-3, -1), (-3, 1)$  (d)  $(3, 1), (3, -1)$
- 
- 15 The shortest distance between the straight line  $x - 2y + 10 = 0$  and the curve  $y^2 = 4x$  equals ..... length unit.
- (a)  $6\sqrt{5}$  (b)  $\frac{3\sqrt{5}}{5}$  (c)  $\frac{4\sqrt{5}}{5}$  (d)  $\frac{6\sqrt{5}}{5}$
- 
- 16 The perimeter of an isosceles triangle is equal to 30 cm., then the side lengths of the triangle such that its surface area is maximum are ..... cm.
- (a) 9, 9, 12 (b) 10, 10, 10 (c) 8, 8, 14 (d) 4, 13, 13
- 
- 17 If the hypotenuse length of a right-angled triangle equals 10 cm., then the lengths of the two legs of the right angle when the area of the triangle is as maximum as possible are ..... cm.
- (a) 6, 8 (b)  $4\sqrt{5}, 2\sqrt{5}$  (c)  $5\sqrt{2}, 5\sqrt{2}$  (d)  $2\sqrt{10}, 4\sqrt{15}$
- 
- 18 The area of the largest rectangle that can be inscribed in a circle of radius 4 cm. equals .....  $\text{cm}^2$
- (a) 28 (b) 32 (c) 48 (d) 64
- 
- 19 A thin metallic lamina in the form of a square, then length of its side is 20 cm. A box without a lid in the form of a rectangular parallelepiped is to be made of this lamina by cutting equal squares of its corners, and turning up the sides, then the length of the side of the removed square when the volume of the box is to be maximum equals ..... cm.
- (a)  $2\frac{1}{2}$  (b) 3 (c)  $3\frac{1}{3}$  (d)  $4\frac{1}{2}$
- 
- 20 A box in the form of a rectangular parallelepiped with a square base., then its maximum volume = .....  $\text{cm}^3$  given that its total surface area =  $384 \text{ cm}^2$
- (a) 1000 (b) 128 (c) 256 (d) 512



- 21 A rectangular parallelepiped with a square base, the sum of all its edges equals 240 cm, then the dimensions of the rectangular parallelepiped when its volume is to be maximum, are ..... cm.  
(a) 20, 20, 20      (b) 15, 15, 30      (c) 16, 16, 28      (d) 18, 18, 24
- 22 A rectangular parallelepiped the length of its base is twice its width. If the sum of its three dimensions 180 cm, then these dimensions that make the volume of the rectangular parallelepiped is to be maximum, are ..... cm.  
(a) 50, 60, 70      (b) 40, 60, 80      (c) 35, 65, 80      (d) 30, 70, 80
- 23 A rectangular parallelepiped has a volume of  $576 \text{ cm}^3$  and the ratio between the two lengths of its base is 2 : 1, then the dimensions of the parallelepiped that makes its total surface area minimum are ..... cm.  
(a) 4, 8, 18      (b)  $8, 16, \frac{9}{2}$       (c) 6, 8, 12      (d)  $4, 6\sqrt{2}, 12\sqrt{2}$
- 24 The selling price of each unit of a product is  $(100 - 0.02 X)$  pounds. Where  $X$  is the number of units which are produced. If the cost price of  $X$  units is  $40 X + 1500$  pounds, then the number of units to be produced to satisfy the maximum profit = ..... units.  
(a) 120      (b) 1 400      (c) 1 500      (d) 1 800
- 25 A factory is producing electric appliances. Profits L.E. 30 in every appliance if it produces 50 appliances monthly. When the production increased than this number, then the profit in each one of all produced appliances decreases by 50 piasters for every extra appliance produced, then the number of appliances produced monthly if the profit is to be maximum = ..... appliance  
(a) 50      (b) 55      (c) 60      (d) 65
- 26 The current intensity  $I$  (Ampere) in a circuit for the alternating current at any moment  $t$  (second) is given by the relation  $I = 2 \cos t + 2 \sin t$ , then the maximum value of the current in this circuit = ..... Ampere  
(a) 2      (b) 3      (c)  $2\sqrt{2}$       (d)  $2\sqrt{3}$
- 27 If the sum of the surface area of a sphere and the surface area of a right circular cylinder, whose radius is equal to the radius of the sphere is  $250 \pi \text{ cm}^2$ , then the radius of the sphere if the sum of their volumes is maximum equals ..... cm.  
(a) 2      (b) 3      (c) 4      (d) 5



- 28 The height of a circular cylinder has maximum volume inside a sphere its radius is "r" equals .....
- (a)  $\frac{2r}{\sqrt{5}}$  (b)  $\frac{2r}{\sqrt{3}}$  (c)  $2r$  (d)  $2\sqrt{3}r$
- 
- 29 The height of a cone which has maximum volume inscribed inside a sphere with radius "r" equals .....
- (a)  $\frac{4r}{5}$  (b)  $\frac{4r}{3}$  (c)  $\frac{2r}{3}$  (d)  $\frac{8r}{3}$
- 
- 30 The height of a right cone , which we can put it inside sphere with radius length 9 cm. , such that its volume is maximum equals ..... cm.
- (a) 7 (b) 12 (c) 8 (d) 10
- 
- 31 In a perpendicular coordinate plane ,  $\overleftrightarrow{AB}$  is drawn to pass through the point C (3 , 2) and intersect the positive parts of the coordinate axes at point A and point B , then the minimum area of triangle AOB where O is the origin point (0 , 0) equals ..... square units.
- (a) 10 (b) 11 (c) 12 (d) 13
- 
- 32 If (a , b) is a fixed point in lattice plane where  $a > 0$  ,  $b > 0$  , a straight line is draw passes through (a , b) and cuts the positive part of each X-axis and y-axis at X , y respectively , if "O" is the origin point , then the smallest area of the triangle OXY , equals ..... square unit.
- (a) ab (b) 2 ab (c) 4 ab (d)  $\frac{1}{2} ab$
- 
- 33 The range of the function  $f : f(x) = \sin x + \cos x$  equals .....
- (a)  $[-\sqrt{2}, \sqrt{2}]$  (b)  $[-1, 1]$  (c)  $[0, 1]$  (d)  $[0, \sqrt{2}]$
- 
- 34 If  $\ell$  , m are the two roots of the function  $x^2 - (k - 1)x + k + 2 = 0$  , then the value of k that makes the expression  $\ell^2 + m^2$  as small as possible equals .....
- (a) 2 (b) 3 (c) 1.5 (d) - 1.5
- 
- 35 The minimum distance between the origin point and the curve  $x = 2 \sin t - \sin 2t$  ,  $y = 2 \cos t - \cos 2t$  , equals ..... length unit.
- (a) 1 (b) 2 (c) 3 (d) 4



- 36 ABCD is a square whose side length 10 cm. and  $M \in \overline{BC}$  where  $BM = X$  cm. and  $N \in \overline{CD}$  where  $CN = \frac{3}{2} X$ , then the value of  $X$  which makes the area of  $\Delta AMN$  as minimum as possible equals .....
- (a)  $\frac{7}{3}$  (b)  $\frac{10}{3}$  (c)  $\frac{11}{3}$  (d)  $\frac{13}{3}$
- 
- 37 The maximum area of the trapezium ABCD in which  $\overline{AB} \parallel \overline{CD}$ ,  $AB = AD = BC = 6$  cm. equals .....  $\text{cm}^2$
- (a) 27 (b)  $27\sqrt{3}$  (c)  $27\sqrt{6}$  (d) 81
- 
- 38 A rectangle has one of its sides on the  $X$ -axis, the upper two vertices of the rectangle lie on the curve  $y = 4 - X^2$ , then the dimensions of the rectangle such that its area is maximum are .....
- (a)  $\frac{4\sqrt{3}}{3}, \frac{8}{3}$  (b)  $\frac{2\sqrt{7}}{3}, \frac{8}{3}$  (c)  $\frac{4\sqrt{3}}{3}, \frac{2}{3}$  (d)  $\frac{4\sqrt{3}}{3}, \frac{2\sqrt{7}}{3}$
- 
- 39 ABCD is a trapezoid where  $\overline{AD} \parallel \overline{BC}$ ,  $\overline{AB} \perp \overline{BC}$ ,  $AB = 20$  cm.,  $AD = 10$  cm.,  $BC = 30$  cm., then the dimensions of the rectangle with the largest area which can be drawn inside the trapezoid are .....
- (a) 12, 12 (b) 12, 15 (c) 15, 15 (d) 15, 16
- 
- 40 If a trapezoid is drawn in a semi-circle such that its base is the diameter of the semi-circle, then the measure of the base angle of the trapezoid such that its area is as maximum as possible = ..... $^\circ$
- (a) 30 (b) 40 (c) 50 (d) 60
- 
- 41 A wire of length 68 cm. is cut into two pieces. The first is bent to form a rectangle of width  $X$  cm. and of length twice its width, while the second is bent to form a square., then the value of  $X$  such that the sum of the rectangle and square areas is minimum = ..... cm.
- (a) 5 (b) 6 (c) 7 (d) 8
- 
- 42 If the point  $(a, b) \in$  the curve of the function  $y = -X^2 + 2X + 3$ , then the greatest value of the expression  $a + b =$  .....
- (a)  $\frac{21}{4}$  (b)  $\frac{21}{2}$  (c)  $\frac{23}{4}$  (d)  $\frac{23}{2}$




- 43 Let  $A(0, 9)$ ,  $B(0, 4)$ ,  $C \in \overrightarrow{OX}$ , then the coordinates of  $C$  which make the measure of  $\angle ACB$  is as great as possible are .....
- (a)  $(3, 0)$  (b)  $(4, 0)$  (c)  $(5, 0)$  (d)  $(6, 0)$
- 
- 44 The area of the largest isosceles triangle that can be inscribed in a circle of radius 15 cm. approximately equals .....  $\text{cm}^2$
- (a) 248.04 (b) 284.28 (c) 292.28 (d) 312.24
- 
- 45 A rectangle is drawn such that two adjacent vertices of the rectangle lie on the curve  $y = x^2 - 12$  and the other two vertices lie on the curve  $y = 12 - x^2$ , then the maximum area of this rectangle equals ..... square units.
- (a) 32 (b) 48 (c) 64 (d) 96
- 
- 46  $\overline{AB}$  is a diameter in a circle whose radius  $r$  and  $\overline{CD}$  is a chord in it where  $\overline{CD} \perp \overline{AB}$  and  $\overline{CD} \cap \overline{AB} = \{N\}$ , then the maximum value of the product  $(CN \times NA) = \dots\dots\dots$
- (a)  $\frac{\sqrt{3}}{2} r^2$  (b)  $\frac{3\sqrt{3}}{4} r^2$  (c)  $\frac{3\sqrt{3}}{2} r^2$  (d)  $\sqrt{3} r^2$
- 
- 47 If  $m$  is the slope of the tangent to the curve  $e^y = 4 + x^2$ , then .....
- (a)  $m \leq \frac{1}{2}$  (b)  $m \geq 2$  (c)  $|m| \leq \frac{1}{2}$  (d)  $|m| \geq 2$
- 

48 In the opposite figure :

Greatest area of  $\triangle ABC = \dots\dots\dots$  square unit

- (a)  $\frac{9}{4}$  (b)  $\frac{11}{4}$   
 (c)  $\frac{13}{4}$  (d)  $\frac{15}{4}$




49 In the opposite figure :

$\triangle ABC$  in which  $\overline{AD} \perp \overline{BC}$

and  $AD + BC = 12$  cm.

, then greatest area of  $\triangle ABC = \dots\dots\dots \text{cm}^2$


- (a) 12 (b) 18  
 (c) 24 (d) 36




**50 (2<sup>nd</sup> session 2021) In the opposite figure :**

If  $AB = 5$  cm. , then the area of the figure ABCD is maximum when .....

- (a)  $x = y$   
 (b)  $x = 2y$   
 (c)  $y = 2x$   
 (d)  $3x = 4y$

**51 In the opposite figure :**


ABCD is a rectangle of perimeter = 28 cm.  
 a circle of centre B is drawn and passes through the two points A and E , then the length of  $\overline{AB}$  that makes the area of the shaded part is as maximum as possible equals .....



- (a)  $\frac{14}{4 + \pi}$       (b)  $\frac{14}{2 + \pi}$       (c)  $\frac{28}{4 + \pi}$       (d)  $\frac{28}{2 + \pi}$


**52 The dimensions of the rectangle of largest area that can be inscribed in the right-angled triangle shown in the figure are ..... cm.**

- (a) 1 , 1.5      (b) 1.5 , 2  
 (c) 2 , 2.5      (d) 2.5 , 3

**53 In the opposite figure :**


$\overline{AB}$  is a diameter in a semi-circle ,  $AB = 16$  cm.  
 , then the greatest area of  $\triangle ADC$  = .....  $\text{cm}^2$

- (a)  $12\sqrt{3}$       (b) 24      (c)  $24\sqrt{3}$       (d) 48

**54 In the opposite figure :**

A rectangle is drawn inside the surface of semi-circle with radius 4 cm.  
 , then the dimensions of this rectangle when its area is maximum are ..... cm.

- (a) 4 , 4      (b)  $4\sqrt{2}$  ,  $2\sqrt{2}$       (c)  $4\sqrt{2}$  ,  $4\sqrt{2}$       (d)  $2\sqrt{3}$  ,  $2\sqrt{2}$





**55 In the opposite figure :**


The value of  $\tan \theta$  which makes  $(AB + BC)$  as small as possible is .....

(a) 2

(b) 1

(c)  $\frac{1}{3}$

(d)  $\frac{1}{2}$



**56 In the opposite figure :**

If the equation of the straight line  $\overleftrightarrow{DE}$  is  $y = 3 - x$

, then the greatest area of the rectangle


$ABCO = \dots\dots\dots$  square unit.

(a)  $\frac{9}{8}$

(b)  $\frac{9}{2}$

(c)  $\frac{9}{4}$

(d)  $\frac{3}{2}$



**57 In the opposite figure :**

The length of  $\overline{AM}$  that makes the volume

of cone is as maximum as possible


equals .....

(a)  $\frac{1}{4}$

(b)  $\frac{1}{2}$

(c) 1

(d)  $1\frac{1}{2}$



**58 The opposite figure represents a rectangle**

inscribed in an isosceles right-angled triangle

$AC = 2$  length units then the greatest area of the


rectangle equals .....

(a)  $\frac{1}{2}$

(b) 1

(c)  $1\frac{1}{2}$

(d) 2





59 In the opposite figure :

A quarter of a circle of centre (O)

, B  $\in$  the curve of the circle  $x^2 + y^2 = 9$

, then the greatest area of the rectangle


ABCO = ..... square unit.

(a) 4.5

(c)  $\frac{3\sqrt{2}}{2}$

(b) 9

(d) 27



60 In the opposite figure :

The part is in the first quadrant from the circle


$x^2 + y^2 = r^2$ , then the greatest perimeter of the rectangle ABCO equals ..... length unit.

(a) r

(c) 2r

(b)  $\sqrt{2}r$

(d)  $2\sqrt{2}r$



61 In the opposite figure :

A quarter circle with radius 4 cm.


, the greatest area of the trapezium ABCD equals .....  $\text{cm}^2$

(a) 6

(c) 12

(b)  $6\sqrt{3}$

(d)  $12\sqrt{3}$



62 In the opposite figure :

$\overline{AB}$  is a diameter in circle whose radius r if  $\overline{AD}$ ,  $\overline{BC}$ ,  $\overline{CD}$

are tangents to it, then smallest area of


the trapezium ABCD = ..... square unit

(a)  $r^2$

(c)  $3r^2$

(b)  $2r^2$

(d)  $4r^2$






**63 In the opposite figure :**


The least length of the line segment  $\overline{OA} = \dots\dots\dots$  length unit.

- (a)  $\sqrt{2}$  (b) 2  
(c)  $2\sqrt{2}$  (d) 4

**64 In the opposite figure :**


If  $CD = 2 AF$  and  $FE = ED$   
and the perimeter of the figure  $ABCDEF = 40$  cm.  
, then the maximum area of the figure  $ABCDEF$   
equals  $\dots\dots\dots$   $\text{cm}^2$ .

- (a) 90 (b) 95 (c) 100 (d) 105

**65 In the opposite figure :**


$f(x) = x^3$ , then the maximum area of the rectangle  $ABCD$  equals  $\dots\dots\dots$  square units.

- (a) 2 (b) 8  
(c) 48 (d) 24

**66 In the opposite figure :**

$B \in$  the curve of the function  $y = x^2 - 3x + 5$   
, then the least perimeter of the rectangle  $OABC$  equals  $\dots\dots\dots$  length unit.

- (a) 3 (b) 8  
(c) 12 (d) 16





67 In the opposite figure :

Two straight lines

$$L_1 : y = 4x \quad , \quad L_2 : y = 18 - 2x$$

, then the greatest area of


the shaded rectangle = ..... square unit.

(a) 25

(b) 27

(c) 30

(d) 32



68 In the opposite figure :

If perimeter of  $\triangle ABC = 8$  cm.


, then greatest area of the triangle = .....  $\text{cm}^2$ .

(a) 2

(b) 3

(c) 4

(d) 5



69 In the opposite figure :

length of  $\overline{OA}$  that makes area of  $\triangle ABC$


as great as possible equals ..... unit length.

(a) 3

(b) 4

(c) 6

(d) 8



70 In the opposite figure :

If equation of the curve is  $y = 4x - x^2$

and equation of straight line L is  $y = 2x$  and

the straight line cuts the curve at the two points

O and B and the point A  $(x_1, y_1)$  moves on

the curve where  $0 < x_1 < 2$ , then greatest area


of  $\triangle ABO = \dots\dots\dots$  square unit.

(a) 1

(b) 2

(c) 3


(d) 4





**71 In the opposite figure :**

If the point A ( $x, y$ ) moves on the straight line  $y = -2x + 4$  where  $x \in [0, 1]$  and the point B is the projection of A on the  $x$ -axis, C ( $-2, 0$ ), then the least possible value of area of  $\triangle ABC$  equals ..... square unit.




- (a) 2
- (b) 3
- (c) 4
- (d) 5

**72 In the opposite figure :**

ABCD is a quadrilateral,  $m(\angle A) = 90^\circ$ ,  $m(\angle C) = \theta$   
 $AB = AD$ ,  $BC = 4$  cm.,  $CD = 8$  cm.


If the area of the figure ABCD is maximum  
 , then  $\theta =$  .....



- (a)  $105^\circ$
- (b)  $120^\circ$
- (c)  $135^\circ$
- (d)  $150^\circ$

**73 In the opposite figure :**

The top right corner of a rectangular piece of paper whose dimension 8 cm., 12 cm. is folded to align the lower edge as shown in the figure, then the value of  $x$  which makes  $y$  as minimum as possible equals ..... cm.




- (a) 5
- (b)  $5\frac{1}{2}$
- (c) 6
- (d)  $6\frac{1}{2}$

**74 In the opposite figure :**

If the point  $A \in$  the curve of the quadratic function  $y = (x - 2)^2$ ,  $\overline{AB} \parallel x$ -axis.


, then the coordinates of A which makes the area of triangle OAB as large as possible is .....

- (a)  $(1\frac{2}{3}, \frac{1}{9})$       (b)  $(\frac{2}{3}, \frac{16}{9})$       (c)  $(1\frac{1}{3}, \frac{4}{9})$       (d) (1, 1)

**75 In the opposite figure :**


The greatest area of the rectangle ABCD equals ..... square units.

- (a)  $\frac{400}{27}$       (b)  $\frac{401}{27}$   
(c)  $\frac{403}{27}$       (d)  $\frac{404}{27}$

**76 In the opposite figure :**


$A \in$  the straight line  $x + 3y = 6$ , the greatest area of the isosceles triangle ABO = ..... square units.

- (a) 2      (b) 3      (c) 4      (d) 6

**77 In the opposite figure :**

If  $\overline{AB} \parallel$  the y-axis, then the greatest value of length of  $\overline{AB}$  = ..... cm.

- (a)  $\frac{4\pi - 3\sqrt{3}}{3}$   
(b)  $\frac{4\pi + 3\sqrt{3}}{3}$   
(c)  $\frac{2\pi - 3\sqrt{6}}{3}$   
(d)  $\frac{2\pi + 3\sqrt{3}}{3}$





**Seventh Questions on behavior of the curves represented graphically**

Choose the correct answer from the given ones :

1 The opposite figure represents the curve of the function  $f$ , then

**First :** The function has local minimum value  
at  $x = \dots\dots\dots$

- (a)  $-3$  (b) zero  
(c)  $2$  (d)  $-2$

**Second :** The absolute maximum values  
equals  $\dots\dots\dots$

- (a)  $5$  (b)  $8$   
(c)  $-2$  (d)  $2$

**Third :** The curve of the function is convex  
upwards at  $x \in \dots\dots\dots$


- (a)  $]-3, 0[$  (b)  $]-2, 2[$   
(c)  $]0, 5[$  (d)  $]-3, 5[$

**Fourth :** The curve of the function  $f$  has an  
inflection point is  $\dots\dots\dots$

- (a)  $(-2, 2)$  (b)  $(0, 0)$  (c)  $(2, -2)$  (d)  $(3\frac{1}{2}, 0)$


**Fifth :** The function is decreasing on the interval  $\dots\dots\dots$

- (a)  $]-3, 0[$  (b)  $]-2, 2[$  (c)  $]0, 5[$  (d)  $]-3, 5[$




2 If  $\ddot{f}(x) < 0$ ,  $\ddot{f}(x) > 0$ , for each  $x \in [a, b]$


, then which of the following curves represents the curve of the function  $f$  in  $[a, b]$ ?




(a)



(b)



(c)




(d)




- 3 The opposite figure represents the curve of the function  $f$ , then  $\hat{f}$  is negative in the interval .....

- (a)  $]1, 2[$  (b)  $]0, 3[$   
(c)  $\mathbb{R} - [1, 2]$  (d)  $]0, 2[$




- 4 The opposite figure represents the curve of the function  $y = f(x)$ , then all the following statements are true except .....

- (a)  $\hat{f}(-2) = \text{zero}$  (b)  $\hat{f}(-1) > 0$   
(c)  $f(3) > f(5)$  (d)  $\hat{f}(-4) < \hat{f}(-5)$




- 5 The opposite figure represents the curve of the function  $y = f(x)$ . All the following statements are true except .....

- (a)  $\hat{f}(-3) + \hat{f}(3) < 0$  (b)  $\hat{f}(-1) > \hat{f}(1)$   
(c)  $\hat{f}(-2) + \hat{f}(0) = 0$  (d)  $\hat{f}(-2) < \hat{f}(0)$



- 6 The opposite figure represents the curve of the function  $f$ , then which of the following is not correct?


- (a) The function  $f$  has three critical points  
(b) The function  $f$  has two inflection points  
(c) The function  $f$  has local maximum value  
(d) Curve of the function  $f$  is convex upward on the interval  $]1, 5[$






- 7 The opposite figure represents the function  $f$ , then sign of the expression  $[f'(0) - f'(2)]$  is .....

- (a) negative (b) positive  
(c) zero (d) insufficient givens




- 8 The function in the opposite figure .....

- (a) has absolute minimum value and absolute maximum value.  
(b) has absolute minimum value but not absolute maximum value.  
(c) has absolute maximum value but not absolute minimum value.  
(d) has neither absolute maximum value nor absolute minimum value.



- 9 If the curve of the function  $f$  has two inflection points, then the opposite figure represents the curve of  $y = \dots\dots\dots$

- (a)  $f(x)$   
(b)  $f'(x)$   
(c)  $f''(x)$   
(d) otherwise




- 10 By using the opposite figure which represents the curve of the function  $f'(x)$ , choose the correct answer from the given ones :

**First :**  $f(x)$  has local maximum value at  $x = \dots\dots\dots$

- (a) zero (b) 1  
(c) 2 (d) 3

**Second :**  $f(x)$  has local minimum value at  $x = \dots\dots\dots$

- (a) zero (b) 1 (c) 2 (d) 3





**Third :**  $f(x)$  has an inflection point at  $x = \dots\dots\dots$

- (a) zero (b) 1  
(c) 2 (d) 3

**Fourth :** The curve of  $f(x)$  is convex upwards at  $x \in \dots\dots\dots$

- (a)  $]-\infty, 2[$  (b)  $]2, \infty[$   
(c)  $]1, 3[$  (d)  $\mathbb{R}$

**Fifth :**  $f(x)$  is decreasing when  $x \in \dots\dots\dots$


- (a)  $]-\infty, 2[$  (b)  $]2, \infty[$   
(c)  $]1, 3[$  (d)  $\mathbb{R}$

**Sixth :** The solution set of the inequality  $\hat{f}(x) \geq 0$  is  $\dots\dots\dots$

- (a)  $]-\infty, 2]$  (b)  $[2, \infty[$   
(c)  $[1, 3]$  (d)  $\mathbb{R}$

- 11** The opposite figure represents the curve of the derivative of the function  $f : f(x) = x^3 + 2ax^2 + bx + 1$ , then  $f(1) = \dots\dots\dots$

- (a) -1 (b) -3  
(c) 7 (d) 11



- 12** In the opposite figure :


If the straight line  $L : 4y = 3x + 12$  touches the curve  $\hat{f}(x)$  at  $x = 2$ , then :

**First :**  $\hat{f}(2) = \dots\dots\dots$

- (a)  $\frac{1}{2}$  (b)  $\frac{3}{4}$   
(c) 4.5 (d) 6

**Second :**  $\hat{f}'(2) = \dots\dots\dots$

- (a)  $\frac{1}{2}$  (b)  $\frac{3}{4}$  (c) 4.5 (d) 6






- 13 If the opposite figure represents the curve  $\hat{f}(x)$ , then :

**First :** The curve  $f(x)$  has local maximum value at  $x = \dots\dots\dots$

- (a) -3 (b) -1  
(c) 4 (d) 6


**Second :** The curve  $f(x)$  has local minimum value at  $x = \dots\dots\dots$

- (a) -3 (b) 1 (c) 4 (d) 6




- 14 The opposite figure represents the curve of  $\hat{f}$ , then all the following statements are correct except .....

- (a) At  $x = -5$  there is a maximum local value of  $f$   
(b)  $\hat{f}(1) > \text{zero}$   
(c) At  $x = 7$  there is a minimum local value of  $f$   
(d) The function  $f$  is decreasing on  $]-5, 1[$




- 15 The opposite figure represents the curve  $\hat{f}(x)$ , then the function  $f$  .....

- (a) has a local maximum value but not a local minimum value.  
(b) has a local minimum value but not a local maximum value.  
(c) has a local minimum value and a local maximum value.  
(d) has neither local maximum value nor local minimum value.



- 16 The opposite figure represents the curve of the first derivative of the function  $y = f(x)$ . All the following statements are true except .....


- (a)  $\hat{f}(-4) > \text{zero}$  (b)  $\hat{f}(-1) = \text{zero}$   
(c)  $\hat{f}(1) < \text{zero}$  (d)  $\hat{f}(-2) \cdot \hat{f}(3) > \text{zero}$






- 17 The opposite figure represents  $\hat{f}$ , then the function  $f$  is increasing in the interval .....

- (a)  $]0, \infty[$   
 (b)  $]-\infty, 0[$   
 (c)  $]2, \infty[$   
 (d)  $]-\infty, 2[$



- 18 The opposite figure represents the curve  $\hat{f}(x)$ , the straight line  $L$  touches the curve at  $(a, b)$ , then  $\hat{f}'(a) = \dots\dots\dots$


- (a)  $\tan \theta$   
 (b)  $-\tan \theta$   
 (c) zero  
 (d)  $-\frac{a}{b}$



- 19 (Trial 2021) The opposite figure represents the curve of the first derivative of the function  $f(x)$  which of the following statements is true for sure ?

- (I)  $f(4) < f(3)$   
 (II) The function  $f$  has a local minimum value at  $x = 5$   
 (III) The function  $f$  has a local maximum value at  $x = 1$


- (a) only III  
 (b) I, II  
 (c) II, III  
 (d) only II



- 20 The opposite figure represents the curve of the function  $\hat{f}(x)$ , then :

**First :** The function  $f$  is increasing on .....

- (a)  $]2, 3[$  only  
 (b)  $]4, \infty[$  only  
 (c)  $\mathbb{R}$   
 (d) both (a), (b)






**Second :** The function  $f$  has an inflection point at .....

- (a)  $x = 2$  (b)  $x = 3$   
(c)  $x = 4$  (d) All the previous.

**21** The opposite figure represents the curve of a function  $y = f(x)$  which is differentiable twice , then .....

- (a)  $f(1) < f'(1) < f''(1)$  (b)  $f''(1) < f(1) < f'(1)$   
(c)  $f''(1) < f'(1) < f(1)$  (d)  $f(1) < f''(1) < f'(1)$



**22** The opposite figure represents the function  $f''(x)$  , then :

**First :** The curve of the function  $f$  is convex upwards in the interval .....


- (a)  $]-\infty, 0[$  (b)  $]-\infty, -3[$   
(c)  $]0, 4[$  (d) b , c together

**Second :** The curve of the function is convex downwards in the interval .....

- (a)  $]-3, 0[$  (b)  $]4, \infty[$  (c)  $]0, \infty[$  (d) a , b together

**Third :** The inflection point at  $x =$  .....

- (a)  $-3$  (b) zero  
(c)  $4$  (d) all the previous are true.




**23** The opposite figure represents the curve of the function  $f''(x)$  , then :

**First :** The curve of the function is convex upwards when  $x \in$  .....

- (a)  $]-\infty, 2[$  (b)  $]2, \infty[$   
(c)  $]0, 2[$  (d) a , c together

**Second :** The curve of the function has inflection point at  $x =$  .....

- (a)  $1$  (b)  $2$   
(c)  $3$  (d) zero





**Third :** If  $\ddot{f}(-1) = \ddot{f}(5) = \text{zero}$ , then the function has local maximum value at  $x = \dots\dots\dots$

(a)  $-1$

(b)  $2$

(c)  $3$

(d)  $5$

**Fourth :** The function  $f$  is decreasing for all  $x \in \dots\dots\dots$

(a)  $]-\infty, -1[$

(b)  $]5, \infty[$

(c)  $\mathbb{R}$

(d)  $a, b$  together


- 24** If the opposite figure represents a continuous curve of  $\ddot{f}(x)$  in the interval  $]-2, 4[$ , then  $\ddot{f}(x)$  is increasing in the interval  $\dots\dots\dots$

(a)  $]-2, 4[$

(b)  $]2, 4[$  only

(c)  $]0, 4[$  only

(d)  $]-2, 0[$  only




- 25** If the opposite figure represents the curve of  $\ddot{f}(x)$  of the function  $f$  where  $-2 \leq x \leq 4$ , then the curve of the function  $f$  is convex upwards in  $\dots\dots\dots$

(a)  $-1 < x < 1$

(b)  $0 < x < 2$

(c)  $-2 < x < -1$  only

(d)  $-2 < x < -1$  and  $1 < x < 3$




- 26** The opposite figure represents the curve of the function  $f$ , then  $f(x) > \ddot{f}(x)$  at  $x \in \dots\dots\dots$

(a)  $\mathbb{R}$

(b)  $]-\infty, 1[$


(c)  $]1, \infty[$

(d)  $]2, \infty[$






- 27 The opposite figure represents the curve of the function  $f$  which is polynomial has an inflection point at  $x = 2$ , then  $f(x)$ ,  $f'(x)$ ,  $f''(x)$  have the same sign at  $x \in \dots\dots\dots$




- (a)  $\mathbb{R}$  (b)  $]-\infty, -2[$   
(c)  $]2, 5[$  (d)  $]5, \infty[$

- 28 The opposite figure shows a graphical representation to the curves of the functions  $f(x)$ ,  $f'(x)$ ,  $f''(x)$  where  $f(x)$  is polynomial, then the curves a, b, c respectively




- (a)  $f(x)$ ,  $f'(x)$ ,  $f''(x)$   
(b)  $f'(x)$ ,  $f''(x)$ ,  $f(x)$   
(c)  $f''(x)$ ,  $f(x)$ ,  $f'(x)$   
(d)  $f''(x)$ ,  $f'(x)$ ,  $f(x)$

- 29 (2<sup>nd</sup> session 2021) If  $f$  is a continuous function has the following properties :  $f(3) = 1$ ,  $f'(x) < 0$  when  $x < 3$ ,  $f'(x) > 0$  when  $x > 3$  and  $f''(x) < 0$  for every  $x \neq 3$ , then the curve which can represent this function is .....







- 30 Which of the following figures represents a general curve of the continuous function  $f$  in which  $f(0) = 3$ ,  $\hat{f}(2) = \hat{f}(-2) = 0$  and  $\hat{f}(x) > 0$  when  $-2 < x < 2$ ?




(a)



(b)




(c)



(d)


- 31 The opposite figure represents the curve of function  $f$ , its domain is  $[a, b]$ , then the function  $g : g(x) = x \cdot f(x)$  is ..... in the interval  $]a, b[$

- (a) decreasing  
(b) increasing  
(c) constant  
(d) not possible to determine its monotony




- 32 The opposite figure represents the curve of the function  $f$  where  $f : [a, b] \rightarrow \mathbb{R}^-$ , then the function  $g : g(x) = \dots$  is increasing in the interval  $]a, b[$

- (a)  $[f(x)]^2$   
(b)  $x \times f(x)$   
(c)  $[f(x)]^3$   
(d)  $-2x - f(x)$




- 33 (2<sup>nd</sup> session 2021) The opposite figure represents the curve of the continuous function  $g(x)$  on  $\mathbb{R}$  that has a horizontal tangent at  $x = 1$ , if  $f$  is differentiable function on  $\mathbb{R}$  where  $\hat{f}(x) = |g(x)|$ , then the true statement from the following is .....
- (a) The function  $f$  has inflection point at  $x = -1$   
(b) The function  $f$  has minimum value at  $x = -1$   
(c) The function  $f$  has inflection point at  $x = 1$   
(d) The curve of the function  $f$  convex upward in  $] -1, 1[$






34 (1<sup>st</sup> session 2021) If the opposite figure represents the curve of the first derivative of a continuous function  $f$  whose domain is  $\mathbb{R}$ , then the wrong statement from the following is .....



- (a) the function has inflection point at  $x = 4$
- (b) the function has local maximum value at  $x = 2$
- (c) the curve of the function convex upward in  $]-\infty, 4[$  and convex downward in  $]4, \infty[$
- (d)  $f(-3) < f(-2)$

35 (Trial 2021) The opposite figure represents the curves of the two functions  $h$  and  $k$  in the interval  $[a, b]$  if  $f$  is a function where  $f(x) = (h \circ k)(x)$ , then the correct statement from the following is .....




- (a) The function  $f$  is decreasing on  $]a, b[$
- (b) The function  $f$  is increasing on  $]a, b[$
- (c) The function  $f$  is increasing on  $]a, c[$  only.
- (d) The function  $f$  is decreasing on  $]a, c[$  only.

36 (Trial 2021) If function  $g$  is differentiable twice in the interval  $]0, 1[$  and  $g''(x) < 0$  and if  $f$  is a function where  $f(x) = g(x) + g(1-x)$ , then the correct statement from the following is .....

- (a)  $f$  is decreasing on  $]\frac{1}{2}, 1[$
- (b)  $f$  is increasing on  $]\frac{1}{2}, 1[$
- (c)  $f$  is decreasing on  $]0, 1[$
- (d)  $f$  is increasing on  $]0, 1[$

37 The opposite figure represents the function  $f$ , then curve of the function  $g : g(x) = f^{-1}(x)$  is .....


- (a) increasing and convex upward.
- (b) increasing and convex downward.
- (c) decreasing and convex upward.
- (d) decreasing and convex downward.






- 38 The opposite figure represents the graphs of  $f$  and  $g$   
If  $h'(x) = f'(x) \cdot g(x)$ , then  $h$  is increasing on .....



- (a)  $]-\infty, -2[$   
(b)  $]-2, 1[$   
(c)  $]1, 4[$   
(d)  $]4, \infty[$



- 39 If  $f: \mathbb{R} \rightarrow \mathbb{R}$  and for all values of  $x \in \mathbb{R}$ ,  $f$  is an increasing function, then the figure which represents  $\hat{f}(x)$  is .....








- 40 The opposite figure represents the curve of the function  $y = f(x)$ , which of the following represents the curve  $\hat{f}(x)$ ?










- 41 The opposite figure represents the first derivative of the function  $y = f(x)$  which of the following figures could be the general shape of the function  $y = f(x)$  ?




- 42 The opposite figure represents the graphs of the two functions  $f(x)$ ,  $g(x)$ . Which of the following graphs could represent the curve of the function  $h$  where  $h(x) = f(x) - g(x)$  ?





- 43 The opposite figure represents the curve of the function  $y = f(x)$ . If the equation of the tangent to the curve at any point  $(x, y)$  is  $y = g(x)$  which of the following statements is right .....

- (a)  $g(x) = f(x)$       (b)  $g(x) \leq f(x)$   
(c)  $g(x) \geq f(x)$       (d)  $f(x) + g(x) < 0$




**44 In the opposite figure :**


A jug of water , water is poured in it in a constant rate with respect to the time and  $y$  is the height of water in the jug , then which of the following graph represents height of water ( $y$ ) during filling the jug ?




(a)



(b)



(c)



(d)

**Eighth Questions on the integration**

Choose the correct answer from the given ones :

1  $\int x(x^2 + 3)^5 dx = \dots\dots\dots + c$

(a)  $\frac{1}{6} (x^3 + 3)^6$

(b)  $\frac{1}{12} (x^2 + 3)^6$

(c)  $\frac{1}{4} (x^2 + 3)^4$

(d)  $\frac{1}{8} (x^2 + 3)^4$

2  $\int \frac{x - \frac{1}{2}}{\sqrt{2x-1}} dx = \dots\dots\dots + c$

(a)  $\frac{1}{6} \sqrt{2x-1}$

(b)  $\frac{1}{6} \sqrt{(2x-1)^3}$

(c)  $\frac{1}{6} \sqrt{2x-1}$

(d)  $\frac{1}{2} \sqrt{(2x-1)^3}$

3  $\int \frac{2x+1}{(x^2+x)^2} dx = \dots\dots\dots + c$

(a)  $\frac{-1}{x^2+x}$

(b)  $\frac{1}{x^2+x}$

(c)  $\frac{-2}{(x^2+x)^2}$

(d)  $\frac{2}{(x^2+x)^3}$

4 If  $\int \frac{x^2 dx}{\sqrt{2x^3+1}} = n\sqrt{2x^3+1} + c$ , then  $n = \dots\dots\dots$

(a) 3

(b)  $\frac{1}{3}$

(c) 6

(d)  $\frac{1}{6}$

5 If  $\int 3x^2(x^n+1)^5 dx = \frac{(x^n+1)^6}{6} + c$ , then  $n = \dots\dots\dots$

(a) 2

(b) 3

(c) 5

(d) 6



6  $\int x^n dx = \frac{x^{n+1}}{n+1} + c$ , for every  $n \neq \dots\dots\dots$

(a) zero

(b) 1

(c) 2

(d) -1

7  $\int \frac{(4x^2 - 4x + 1)^7}{(2x - 1)^2} dx = \dots\dots\dots + c$

(a)  $\frac{1}{13} (2x - 1)^{13}$

(b)  $\frac{1}{26} (2x - 1)^{13}$

(c)  $\frac{2}{3} (2x - 1)^{13}$

(d)  $\frac{1}{16} (2x - 1)^8$

8  $\int \frac{(2x + 1)(x - 2)}{\sqrt{x}} dx = \dots\dots\dots + c$

(a)  $2x^{1\frac{1}{2}} - 3x^{\frac{1}{2}}$

(b)  $\frac{(2x + 1)^2 (x - 1)^2}{x^{1\frac{1}{2}}}$

(c)  $2x^{1\frac{1}{2}} - 3x^{\frac{1}{2}} - 2x^{-\frac{1}{2}}$

(d)  $\frac{4}{5} x^{2\frac{1}{2}} - 2x^{1\frac{1}{2}} - 4x^{\frac{1}{2}}$

9  $\int \left(x + \frac{1}{x}\right)^2 dx = \dots\dots\dots + c$

(a)  $\frac{1}{3} \left(x + \frac{1}{x}\right)^3$

(b)  $\frac{1}{3} x^3 + 2x - \frac{1}{x}$

(c)  $\frac{1}{3} x^3 + 2x + \frac{1}{x}$

(d)  $\left(x + \frac{1}{x}\right)^3$

10  $\int \left(x - \frac{1}{x}\right) \left(x + \frac{1}{x}\right) \left(x^2 + \frac{1}{x^2}\right) dx = \dots\dots\dots + c$

(a)  $x^4 - \frac{1}{x^4}$

(b)  $\frac{1}{5} x^5 - \frac{5}{x^5}$

(c)  $\frac{1}{5} x^5 - x^{-3}$

(d)  $\frac{1}{5} x^5 + \frac{1}{3} x^{-3}$

11  $\int 6x^5 \left(1 - \frac{1}{x}\right)^5 dx = \dots\dots\dots + c$

(a)  $6(x - 1)^6$

(b)  $(x - 1)^6$

(c)  $\frac{1}{6} (x - 1)^6$

(d)  $(6x - 1)^6$

12  $\int x \sqrt{x - 2} dx = \dots\dots\dots + c$

(a)  $(x - 2)^{\frac{3}{2}} + 2\sqrt{x - 2}$

(b)  $\frac{2}{3} (x^2 - 2x)^{\frac{3}{2}}$

(c)  $\frac{2}{5} (x - 2)^{\frac{5}{2}} + \frac{2}{3} (x - 2)^{\frac{3}{2}}$

(d)  $\frac{2}{5} (x - 2)^{\frac{5}{2}} + \frac{4}{3} (x - 2)^{\frac{3}{2}}$



13  $\int \frac{x+3}{\sqrt{x-1}} dx = \dots\dots\dots + c$

(a)  $(x-2)^{\frac{3}{2}} - (x-1)^{\frac{1}{2}}$

(b)  $\frac{3}{2}(x-1)^{\frac{3}{2}} + \frac{1}{2}(x-1)^{\frac{1}{2}}$

(c)  $\frac{2}{3}(x-1)^{\frac{3}{2}} + 2(x-1)^{\frac{1}{2}}$

(d)  $\frac{2}{3}(x-1)^{\frac{3}{2}} + 8(x-1)^{\frac{1}{2}}$

14  $\int x(x+2)^8 dx = \dots\dots\dots + c$

(a)  $\frac{1}{9}(x+2)^9$

(b)  $\frac{1}{9}(x^2+2x)^9$

(c)  $\frac{1}{10}(x+2)^{10} - \frac{2}{9}(x+2)^9$

(d)  $\frac{1}{10}(x+2)^{10} + \frac{1}{9}(x+2)^9$

15  $\int x f(1-x^2) dx = \dots\dots\dots + c$

(a)  $-f(1-x^2)$

(b)  $-2f(1-x^2)$

(c)  $-\frac{1}{2}f(1-x^2)$

(d)  $xf(1-x^2)$

16  $\int x \cdot f(x^2) \cdot f(x^2) dx = \dots\dots\dots + c$

(a)  $\frac{1}{4}[f(x^2)]^2$

(b)  $\frac{1}{2}[f(x^2)]^2$

(c)  $[f(x^2)]^2$

(d)  $2[f(x^2)]^2$

17  $\int \sqrt{x^3-3x^2+3x-4}(x-1)^2 dx = \dots\dots\dots + c$

(a)  $\sqrt{\frac{1}{4}x^4 - x^3 + \frac{3}{2}x^2 - 4x}$

(b)  $\frac{2}{9}\sqrt{(x^3-3x^2+3x-3)^3}$

(c)  $\frac{2}{9}\sqrt{x^3-3x^2+3x-4}$

(d)  $\frac{1}{2}(x^3-3x^2+3x-4)^{\frac{2}{3}}$

18  $\int x^3(x^2-1)^5 dx = \dots\dots\dots + c$

(a)  $\frac{1}{6}(x^2-1)^6$

(b)  $\frac{1}{6}x^6 - \frac{1}{4}x^4$

(c)  $\frac{1}{7}(x^2-1)^7 + \frac{1}{6}(x^2-1)^6$

(d)  $\frac{1}{14}(x^2-1)^7 + \frac{1}{12}(x^2-1)^6$

19  $\int \frac{dx}{\sqrt{x}(\sqrt{x}+2)^4} = \dots\dots\dots + c$

(a)  $-\frac{2}{3}(\sqrt{x}+2)^{-3}$

(b)  $-\frac{1}{3}(\sqrt{x}+2)^{-3}$

(c)  $(\sqrt{x}+2)^{-3}$

(d)  $-\frac{1}{2}(\sqrt{x}+2)^{-3}$



20  $\frac{d}{dX} \int (\sin X + 3)^4 dX = \dots\dots\dots$

(a)  $\frac{1}{5} (\sin X + 3)^5 + c$

(b)  $(\sin X + 3)^4$

(c)  $(\sin X + 3)^4 + c$

(d)  $4 (\sin X + 3)^3$

21  $\int \frac{d}{dX} (X^5 + \sqrt{X}) dX = \dots\dots\dots$

(a)  $\frac{1}{6} X^6 + \frac{1}{2\sqrt{X}} + c$

(b)  $X^4 + \frac{1}{2} X^{-\frac{1}{2}}$

(c)  $X^5 + \sqrt{X}$

(d)  $X^5 + \sqrt{X} + c$

22  $\int \cos (3X - 1) dX = \dots\dots\dots + c$

(a)  $\sin (3X - 1)$

(b)  $\frac{1}{3} \sin (3X - 1)$

(c)  $3 \sin (3X - 1)$

(d)  $-\frac{1}{3} \sin (3X - 1)$

23  $\int \csc^2 (3X) dX = \dots\dots\dots + c$

(a)  $-\cot (3X)$

(b)  $\frac{1}{3} \cot (3X)$

(c)  $-3 \cot 3X$

(d)  $-\frac{1}{3} \cot (3X)$

24  $\int \sec y \tan y dy = \dots\dots\dots + c$

(a)  $\sec y$

(b)  $\csc y$

(c)  $\tan y$

(d)  $\cot y$

25  $\int (\sin^2 5X + \cos^2 5X)^{2021} dX = \dots\dots\dots + c$

(a)  $\sin^{-1} X + \cos^{-1} X$

(b) 1

(c)  $X$

(d)  $\frac{1}{5} (\sin^3 5X + \cos^3 5X)$

26  $\int \left( \csc^2 \frac{\pi}{4} + \cos \frac{\pi}{3} \right) dX = \dots\dots\dots + c$

(a)  $-\cot \frac{\pi}{4} + \sin \frac{\pi}{3}$

(b)  $-4 \cot \frac{\pi}{4} + 3 \sin \frac{\pi}{3}$

(c)  $2.5 X$

(d)  $\cot \frac{\pi}{4} - \sin \frac{\pi}{3}$

27  $\int \frac{3}{\sin^2 3X} dX = \dots\dots\dots + c$

(a)  $\frac{-3}{\cos^2 3X}$

(b)  $3 \csc^2 3X$

(c)  $-\cot 3X$

(d)  $-3 \cot 3X$



28  $\int (\sin^2 x + \cos^2 x + \tan^2 x) dx = \dots\dots\dots$

(a)  $\sin x + \cos x + \tan x + c$

(b)  $\tan x + c$

(c)  $\frac{1}{2} x + c$

(d)  $\sec^2 x + c + \cos x \sec x$

29  $\int (\sin x \csc x + \cos x \sec x + \tan x \cot x) dx = \dots\dots\dots + c$

(a) 3

(b)  $3x$

(c) zero

(d)  $-\cot x$

30 If  $\int 4 \cos(ax) dx = 12 \sin(ax) + c$ , then  $a = \dots\dots\dots$

(a)  $\frac{1}{3}$

(b) 3

(c) 6

(d) 12

31  $\int (\cos \frac{x}{2} + \sin \frac{x}{2})^2 dx = \dots\dots\dots + c$

(a)  $1 + \sin x$

(b)  $x$

(c)  $x - \cos x$

(d)  $x + \cos x$

32  $\int (2 \cos^2 x - 1) dx = \dots\dots\dots + c$

(a)  $\frac{1}{2} \sin 2x$

(b)  $\frac{1}{2} \cos 2x$

(c)  $\sin 2x$

(d)  $\cos 2x$

33  $\int (1 + \cot^2 x) dx = \dots\dots\dots + c$

(a)  $\cot x$

(b)  $-\cot x$

(c)  $\tan^2 x$

(d)  $-\cot^2 x$

34  $\int (4 - \csc x \cot x) dx = \dots\dots\dots$

(a)  $4x - \csc x + c$

(b)  $4x + \csc x + c$

(c)  $4x - \cot x + c$

(d)  $4x + \cot x + c$

35  $\int (\sin 3x \cos x - \cos 3x \sin x) dx = \dots\dots\dots + c$

(a)  $\sin 2x$

(b)  $-\frac{1}{2} \cos 2x$

(c)  $\cos 2x$

(d)  $-\frac{1}{4} \cos 4x$

36  $\int \cos x \cos \frac{\pi}{4} - \sin x \sin \frac{\pi}{4} dx = \dots\dots\dots + c$

(a)  $\cos(x + \frac{\pi}{4})$

(b)  $\sin(x + \frac{\pi}{4})$

(c)  $\frac{1}{4} \cos(x + \frac{\pi}{4})$

(d)  $\frac{1}{4} \sin(x + \frac{\pi}{4})$



37  $\int \sin x \cos x \cos 2x \cos 4x \, dx = \dots + c$

- (a)  $-\cos 8x$  (b)  $-\frac{1}{64} \cos 8x$  (c)  $\cos 8x$  (d)  $\frac{1}{64} \cos 8x$

38  $\int \tan^2 x \csc^2 x \, dx = \dots + c$

- (a)  $\tan x$  (b)  $\sec^2 x$  (c)  $\csc^2 x$  (d)  $-\cot x$

39  $\int (\sin^2 x + \sin^2 x \tan^2 x) \, dx = \dots + c$

- (a)  $\sin^2 x + \csc^2 x$  (b)  $\tan x - x$  (c)  $\tan^2 x$  (d)  $\sec x$

40  $\int (\cot^4 x + \cot^2 x) \, dx = \dots + c$

- (a)  $\frac{1}{3} \cot^3 x$  (b)  $\tan x$  (c)  $\log |\sin^2 x|$  (d)  $-\frac{1}{3} \cot^3 x$

41 Sandy and Youssef found the integration  $\int (\tan x + \tan^3 x) \, dx$ ,

Sandy solution was  $\frac{1}{2} \sec^2 x + a$ , while Youssef solution was  $\frac{1}{2} \tan^2 x + b$

, if each of them got the full mark, then  $a - b = \dots$

- (a)  $-1$  (b)  $-\frac{1}{2}$  (c)  $\frac{1}{2}$  (d)  $1$

42  $\int (1 + \tan^2 x) \cos^2 x \, dx = \dots + c$

- (a)  $x$  (b)  $\frac{1}{3} \cos^3 x$  (c)  $\frac{1}{3} \sec^3 x$  (d)  $\frac{1}{3} \tan^3 x$

43  $\int (1 + \csc x)(1 - \csc x) \, dx = \dots + c$

- (a)  $x - \cot x$  (b)  $x + \cot x$   
(c)  $\frac{1}{2} \cot^2 x$  (d)  $\frac{1}{2} x^2 + \cot x$

44  $\int \sec x (\sec x + \tan x) \, dx = \dots + c$

- (a)  $\csc x + \cot x$  (b)  $\sec^2 x + \tan x$  (c)  $\sec x + \tan x$  (d)  $\sec x - \tan x$

45  $\int \frac{1}{1 - \cos^2 x} \, dx = \dots + c$

- (a)  $\cot x$  (b)  $-\cot x$  (c)  $\tan x$  (d)  $-\tan x$



46  $\int \frac{\cos x + \cos^3 x}{\sin^2 x + 2 \cos^2 x} dx = \dots + c$

(a)  $\sin x$

(b)  $\cos x$

(c)  $\tan x$

(d)  $\csc x$

47  $\int \cos^5 x \sin x dx = \dots + c$

(a)  $\frac{1}{6} \sin^6 x$

(b)  $-\frac{1}{6} \sin^6 x$

(c)  $\frac{1}{6} \cos^6 x$

(d)  $-\frac{1}{6} \cos^6 x$

48  $\int x^2 \sec^2(x^3 + 5) dx = \dots + c$

(a)  $\frac{1}{3} x^3 \sec^3(x^3 + 5)$

(b)  $\frac{1}{3} \sec^3(x^3 + 5)$

(c)  $\frac{1}{3} \tan(x^3 + 5)$

(d)  $3 \tan(x^3 + 5)$

49  $\int 10x \sec(5x^2 + 2021) \tan(5x^2 + 2021) dx = \dots + c$

(a)  $\sec^2(5x^2 + 2021)$

(b)  $\cos(5x^2 + 2021)$

(c)  $\sin(5x^2 + 2021)$

(d)  $\sec(5x^2 + 2021)$

50  $\int (\sin x + \cot x)^8 (\cos x - \csc^2 x) dx = \dots + c$

(a)  $\frac{1}{9} (\sin x + \cot x)^9$

(b)  $8 (\sin x + \cot x)^7$

(c)  $\frac{1}{9} \sin^9 x + \frac{1}{9} \cot^9 x$

(d)  $\frac{1}{2} (\cos x - \cot^2 x)^2$

51  $\int \cos(\tan x + 1) \sec^2 x dx = \dots + c$

(a)  $\cos^2(\tan x + 1)$

(b)  $\sin(\tan x + 1)$

(c)  $\frac{1}{2} \sec^3 x \times \sin(\tan x + 1)$

(d)  $\sec(\tan x + 1) \tan(\tan x + 1)$

52  $\int \frac{\sec x}{\sec x + \tan x} dx = \dots + c$

(a)  $\tan x \sec x$

(b)  $\frac{\sec x + \tan x}{\sec x}$

(c)  $\tan x - \sec x$

(d)  $\tan x + \sec x$

53  $\int \cos x f(\sin x) dx = \dots + c$

(a)  $f(\cos x)$

(b)  $f(\sin x)$

(c)  $\frac{1}{2} (f(x))^2$

(d)  $\frac{1}{2} (f(\cos x))^2$



54  $\int [(1 - \cot x)^2 + 2 \cot x] dx = \dots\dots\dots + c$

- (a)  $-\cot x$  (b)  $\csc^2 x$  (c)  $x + \frac{\cot^2 x}{3}$  (d)  $\cot x$

55  $\int \left( \frac{\sin^2 x + 1}{\sin x} \right)^2 dx = \dots\dots\dots + c$

- (a)  $2 \frac{1}{2} x - \frac{1}{2} \sin 2x + \cot x$  (b)  $\frac{1}{2} - \frac{1}{2} \cos 2x - \frac{1}{4} \sin 2x + \cot x$   
 (c)  $2 \frac{1}{2} x - \frac{1}{4} \sin 2x - \cot x$  (d)  $2 \frac{1}{2} x - \frac{1}{4} \sin x - \cot x$

56  $\int \frac{1 + \sin^2 x}{1 - \sin^2 x} dx = \dots\dots\dots + c$

- (a)  $\sec^2 x + \tan^2 x$  (b)  $\tan x + x$  (c)  $\tan x - x$  (d)  $2 \tan x - x$

57  $\int \frac{\sin^3 x + \cos^3 x}{\sin x + \cos x} dx = \dots\dots\dots + c$

- (a)  $1 - \frac{1}{2} \sin 2x$  (b)  $x + \frac{1}{4} \cos 2x$  (c)  $1 + \cos x \sin x$  (d)  $1 - \frac{1}{4} \cos x$

58  $\int \frac{\cos 2x}{\cos x + \sin x} dx = \dots\dots\dots + c$

- (a)  $\sin x - \cos x$  (b)  $\cos x - \sin x$   
 (c)  $\sin x + \cos x$  (d)  $\cos 2x + \sin 2x$

59  $\int 4 \sin^4 x dx = \dots\dots\dots + c$

- (a)  $-\frac{4}{5} \cos^5 x$  (b)  $\frac{4}{5} \cos^5 x$   
 (c)  $x - \sin 2x + \cos 4x$  (d)  $\frac{3}{2} x - \sin 2x + \frac{1}{8} \sin 4x$

60  $3 \int \sin 2x \sin^4 x dx = \dots\dots\dots + c$

- (a)  $\sin^4 x \cos^2 x$  (b)  $\cos^4 x \sin^2 x$  (c)  $\sin^6 x$  (d)  $\cos^6 x$

61  $\int \frac{\sin^6 x}{\cos^8 x} dx = \dots\dots\dots + c$

- (a)  $\tan^7 x$  (b)  $\frac{1}{7} \tan^7 x$  (c)  $\frac{1}{7} \tan 7x$  (d)  $\sec^7 x$



62  $\int (\sec x + \cos x)^2 dx = \dots\dots\dots + c$

(a)  $\frac{1}{3} (\sec x + \cos x)^3$

(b)  $\cot x - \sin^3 x$

(c)  $\frac{5}{2} x - \sin 2x - \tan x$

(d)  $\frac{5}{2} x + \frac{1}{4} \sin 2x + \tan x$

63  $\int \frac{\tan x}{\cos x} dx = \dots\dots\dots + c$

(a)  $\sec^2 x$

(b)  $\sec x$

(c)  $\tan x \sec x$

(d)  $\ln \cos x$

64  $\int \frac{\cos x}{\csc x} dx = \dots\dots\dots + c$

(a)  $\frac{1}{2} \sin 2x$

(b)  $\frac{1}{4} \cos 2x$

(c)  $-\frac{1}{4} \cos 2x$

(d)  $-4 \cos 2x$

65  $\int \tan^2 x dx = \dots\dots\dots$

(a)  $\tan x - x + c$

(b)  $\tan x + x + c$

(c)  $\sec^4 x + c$

(d)  $\frac{1}{3} \tan^3 x + c$

66  $\int \sec^4 x \tan x dx = \dots\dots\dots$

(a)  $\frac{1}{5} \sec^5 x + c$

(b)  $\frac{1}{4} \sec^4 x + c$

(c)  $\frac{1}{3} \tan^3 x + c$

(d)  $-\frac{1}{3} \tan^3 x + c$

67  $\int \frac{\sec^2 x}{1 + \tan^2 x} dx = \dots\dots\dots + c$

(a)  $2 \sec^2 x$

(b)  $\sec^2 x \tan^2 x$

(c)  $2 \sec^2 x$

(d)  $x$

68  $\int \left( \frac{\tan x}{\cot x} + 1 \right) dx = \dots\dots\dots$

(a)  $\tan^2 x + c$

(b)  $\tan x + c$

(c)  $\tan x \sec x + c$

(d)  $\cot x \csc x + c$

69  $\int e^2 dx = \dots\dots\dots + c$

(a)  $e^2 x$

(b)  $\frac{1}{3} e^3$

(c)  $e^2$

(d)  $\frac{1}{3} e^{3x}$

70  $\int x^2 d(x^2) = \dots\dots\dots + c$

(a)  $\frac{1}{3} x^3$

(b)  $\frac{1}{4} x^4$

(c)  $\frac{1}{2} x^3$

(d)  $\frac{1}{2} x^4$



71  $\int e^{\ln x} dx = \dots + c$

- (a)  $\frac{x^2}{2} \ln x$  (b)  $\ln x$  (c)  $\frac{1}{2} x^2$  (d)  $e^x$

72  $\int \left( \sum_{n=0}^{\infty} \frac{x^n}{n} \right) dx = \dots + c$

- (a)  $e^x$  (b)  $e^x$  (c)  $x^{2e}$  (d)  $e^{x^2}$

73  $\int e^{6x} dx = \dots + c$

- (a)  $e^{6x}$  (b)  $\frac{1}{6} e^{6x}$  (c)  $\frac{1}{7} e^{7x}$  (d)  $\frac{1}{7} e^{6x+1}$

74  $\int a^{3 \log_a x} dx = \dots + c$

- (a)  $a^x$  (b)  $e^{x^3}$  (c)  $x^4$  (d)  $\frac{1}{4} x^4$

75  $\int \frac{3}{x} dx = \dots + c$

- (a)  $\frac{-3}{2} x^{-2}$  (b)  $3 \ln |x|$  (c)  $\ln |3x|$  (d)  $\ln |x|$

76  $\int \frac{\log_4 e}{x} dx = \dots + c$

- (a)  $\ln |4-x|$  (b)  $\ln |4+x|$  (c)  $\log_4 |x|$  (d)  $\ln |x|$

77 (2<sup>nd</sup> session 2021) If  $\int \frac{2}{y} dy = \int \frac{1}{x} dx$ , then  $\ln y^2 = \dots + c$  where  $c$  is constant

- (a)  $\ln |x|$  (b)  $|x|$  (c)  $\ln |x+y|$  (d)  $\ln |x-y|$

78 If:  $\int \frac{4x^3 - ax}{x^4 - 2x^2 + 3} dx = \ln |x^4 - 2x^2 + 3| + c$   
then  $a = \dots$

- (a) 1 (b) 2 (c) 3 (d) 4

79 (2<sup>nd</sup> session 2021) If  $z = x + 2$ , then  $\int \frac{x-2}{x^2 + 4x + 4} dx = \dots + c$  where  $c$  is constant

- (a)  $\ln |z| - \frac{4}{z}$  (b)  $\ln |z| + \frac{4}{z}$  (c)  $\frac{-1}{2z^2} - \frac{4}{3z^3}$  (d)  $\frac{-1}{2z^2} + \frac{2}{3z^3}$



80  $\int \left( 2 \sin x + \frac{1}{x} \right) dx = \dots\dots\dots$

(a)  $-2 \cos x + \ln |x| + c$

(b)  $2 \cos x + \ln |x| + c$

(c)  $-\sin x - \frac{1}{x^2} + c$

(d)  $-2 \cos x + \frac{1}{x^2} + c$

81  $\int \frac{e^x + e^{-x}}{e^x} dx = \dots\dots\dots + c$

(a)  $x + e^{-2x}$

(b)  $1 + e^{-2x}$

(c)  $x - e^{-2x}$

(d)  $x - \frac{1}{2} e^{-2x}$

82  $\int \frac{e^{3x} - 1}{e^x - 1} dx = \dots\dots\dots + c$

(a)  $\frac{1}{2} e^{2x} + e^x + x$

(b)  $e^{2x} + e^x + 1$

(c)  $e^{2x} - e^x + 1$

(d)  $\frac{1}{3} e^{3x} - x$

83 If:  $\int e^{f(x)} dx = \frac{1}{2} x^2 + c$ , then  $f(x)$  could be .....

(a)  $x$

(b)  $\log x$

(c)  $\ln x$

(d)  $x^2$

84  $\int \frac{e^{2x} - 1}{e^{2x} + 1} dx = \dots\dots\dots + c$

(a)  $\ln |e^{2x} + 1|$

(b)  $\ln |e^{2x} - 1|$

(c)  $\ln |e^x + e^{-x}|$

(d)  $\ln |e^x + 1|$

85 If  $\int \frac{e^{ax}}{e^{bx}} dx = k \times \frac{e^{ax}}{e^{bx}} + c$ , then  $k = \dots\dots\dots$

(a)  $a + b$

(b)  $a - b$

(c)  $\frac{1}{a - b}$

(d)  $\frac{1}{a + b}$

86  $\int (x^{2e} + e^{3x}) dx = \dots\dots\dots + c$

(a)  $2x^{2e} + 3x^{3e}$

(b)  $\frac{1}{3} x^{3e} + e^{3x}$

(c)  $\frac{1}{3e} x^{3e} + \frac{1}{3} e^{3x}$

(d)  $\frac{x^{2e+1}}{2e+1} + \frac{1}{3} e^{3x}$

87  $\int 4x e^{x^2} dx = \dots\dots\dots$

(a)  $\frac{1}{2} e^{x^2} + c$

(b)  $e^{x^2} + c$

(c)  $2e^{x^2} + c$

(d)  $4e^{x^2} + c$

88  $\int x^2 e^{x^3+1} dx = \dots\dots\dots + c$

(a)  $e^{x^3+1}$

(b)  $3e^{x^3+1}$

(c)  $\frac{1}{3} e^{x^3+1}$

(d)  $3e^{x^4+x}$



89  $\int (13)^x dx = \dots\dots\dots$

- (a)  $\frac{13^x}{\ln 13} + c$       (b)  $(14)^{x+1} + c$       (c)  $(13)^{x+1} + c$       (d)  $14x + c$

90  $\int e^x \cdot \tan(e^x) dx = \dots\dots\dots + c$

- (a)  $\ln |\sec e^x|$       (b)  $\ln |\sin e^x|$       (c)  $\ln |\cos e^x|$       (d)  $\tan e^x$

91  $\int e^{\ln(\sin x)} dx = \dots\dots\dots$

- (a)  $-\cos x + c$       (b)  $\sin x + c$       (c)  $\cos x + c$       (d)  $-\sin x + c$

92  $\int \sin x e^{\cos x} dx = \dots\dots\dots + c$

- (a)  $-e^{\sin x}$       (b)  $-e^{\cos x}$       (c)  $e^{\sin x}$       (d)  $e^{\cos x}$

93 (2<sup>nd</sup> session 2021)  $\int 4 \sin 2x e^{\cos 2x} dx = \dots\dots\dots + c$ , where c is constant

- (a)  $-2e^{\cos 2x}$       (b)  $2e^{\cos 2x}$       (c)  $4e^{\cos 2x}$       (d)  $-4e^{\cos 2x}$

94  $\int e^{(x+e^x)} dx = \dots\dots\dots + c$

- (a)  $e^x$       (b)  $e^{e^x}$       (c)  $e^{x+e^x}$       (d)  $x+e^x$

95  $\int \frac{x^3 dx}{x^4 + 3} = \dots\dots\dots + c$

- (a)  $\frac{1}{4}(x^4 + 3)$       (b)  $\frac{1}{4} \ln |x^4 + 3|$       (c)  $\ln |x^4 + 3|$       (d)  $\frac{1}{4}(x^4 + 3)^{-1}$

96  $\int \frac{\ln x}{x} dx = \dots\dots\dots + c$

- (a)  $\ln x^2$       (b)  $(\ln x)^2$       (c)  $\frac{1}{2} \ln x^2$       (d)  $\frac{1}{2} (\ln x)^2$

97  $\int \frac{\ln x^2}{\ln x} dx = \dots\dots\dots$

- (a)  $\frac{x}{2} + c$       (b)  $\frac{1}{x} + c$       (c)  $2x + c$       (d)  $\ln |x| + c$

98  $\int \frac{1}{x \ln x^3} dx = \dots\dots\dots$

- (a)  $3 \ln |x| + c$       (b)  $3 \ln |\ln x| + c$   
(c)  $\frac{1}{3} \ln |x| + c$       (d)  $\frac{1}{3} \ln |\ln x| + c$



99  $\int \frac{\ln x^5}{x \ln x^3} dx = \dots + c$

- (a)  $\frac{5}{3} \ln |x|$  (b)  $\frac{3}{5} \ln |x|$  (c)  $\ln (\ln |x|)$  (d)  $\ln \left(\frac{5}{3} |x|\right)$

100  $\int \frac{x+3}{x-1} dx = \dots + c$

- (a)  $1 + \ln |x+3|$  (b)  $x + \ln |x-1|$  (c)  $1 + \ln |x-1|$  (d)  $x + 4 \ln |x-1|$

101  $\int \frac{x^2-25}{x^2-5x} dx = \dots + c$

- (a)  $\ln |x+5|$  (b)  $x + 5 \ln |x|$   
(c)  $5x + \ln |x|$  (d)  $x + \ln |x+5|$

102 (1<sup>st</sup> session 2021)  $\int \frac{x}{x^2-2x+1} dx = \dots + c$ , where  $c$  is constant.

- (a)  $\ln |x-1| - \frac{1}{x-1}$  (b)  $\ln |x-1| + \frac{1}{x-1}$   
(c)  $\ln |x-1| + (x-1)^2$  (d)  $\ln |x^2-2x+1| + \frac{1}{x-1}$

103  $\int \frac{1+\ln x}{8+x \ln x} dx = \dots + c$

- (a)  $\ln |x \ln x|$  (b)  $\ln |8+x \ln x|$  (c)  $8 + \ln |x|$  (d)  $\ln 8 + \ln x$

104  $\int \frac{6}{x+x \ln x} dx = \dots + c$

- (a)  $6(1+\ln x)$  (b)  $6 \ln |1+\ln x|$   
(c)  $\frac{6}{\ln |1+\ln x|}$  (d)  $\frac{6}{\ln |x+\ln x|}$

105 If  $\int \frac{\ln(x)^a}{x} dx = 5(\ln x)^2 + c$ , then  $a = \dots$

- (a)  $\frac{2}{5}$  (b)  $\frac{5}{2}$  (c) 5 (d) 10

106  $\int \tan \theta d\theta = \dots$

- (a)  $-\ln |\cos \theta| + c$  (b)  $-\ln \cos \theta + c$  (c)  $\ln \cos \theta + c$  (d)  $|\ln \cos \theta| + c$



$$107 \int \frac{2 \tan x}{1 - \tan^2 x} dx = \dots + c$$

(a)  $\frac{1}{2} \tan 2x$

(b)  $2 \sec^2 2x$

(c)  $-\ln |\cos x|$

(d)  $-\frac{1}{2} \ln |\cos 2x|$

$$108 \int \frac{\sin 2x}{1 + \sin^2 x} dx = \dots + c$$

(a)  $(1 + \sin^2 x)^{-2}$

(b)  $x + \frac{1}{3} \sin^3 x$

(c)  $\ln |1 + \sin^2 x|$

(d)  $-\frac{1}{2} \cos 2x + \frac{1}{3} \sin^3 x$

$$109 \int \frac{\cos x - \sin x}{\cos x + \sin x} dx = \dots + c$$

(a)  $\ln \cos x - \ln \sin x$

(b)  $\ln \cos x + \ln \sin x$

(c)  $\ln |\cos x - \sin x|$

(d)  $\ln |\cos x + \sin x|$

$$110 \int \csc x dx = \dots + c$$

(a)  $-\csc x \cot x$

(b)  $-\ln |\csc x + \cot x|$

(c)  $\sin^{-1} x$

(d)  $\frac{1}{2} \csc^2 x$

$$111 \int \frac{x^2}{x+1} dx = \dots + c$$

(a)  $\frac{1}{2} x^2 - x + \ln |x+1|$

(b)  $\ln |x+1|$

(c)  $\frac{1}{2} x^2 + \frac{1}{3} x^3$

(d)  $(x-1)^2 + \ln |x+1|$

$$112 \int \frac{6}{x} (\ln x)^5 dx = \dots + c$$

(a)  $(\ln x)^6$

(b)  $\frac{1}{6} (\ln x)^6$

(c)  $\ln x^6$

(d)  $\frac{1}{6} \ln x^6$

$$113 \int \frac{\sqrt{5 + \ln x}}{x} dx = \dots + c$$

(a)  $(5 + \ln x)^{\frac{3}{2}}$

(b)  $\frac{2}{3} (5 + \ln x)$

(c)  $\frac{1}{2} \sqrt{5 + \ln x}$

(d)  $\frac{2}{3} (5 + \ln x)^{\frac{3}{2}}$

$$114 \int \sqrt{1 - \cos^2 x} dx = \dots + c \text{ where } x \in ]0, \pi[$$

(a)  $\csc x$

(b)  $-\cos x$

(c)  $\cos^2 x$

(d)  $\sin^2 x$



115  $\int \cos^{10} x \sec^9 x \, dx = \dots\dots\dots + c$

(a)  $\frac{1}{\pi} \cos^{11} x$

(b)  $\frac{6}{10} \sec^{10} x$

(c)  $\cos x$

(d)  $\sin x$

116  $\int \cos^9 x \sec^{10} x \, dx = \dots\dots\dots + c$

(a)  $\sec x$

(b)  $\frac{1}{2} \sec^2 x$

(c)  $\ln |\sec x + \tan x|$

(d)  $\sec x \tan x$

117  $\int (1 + \tan^2 x) e^{1 + \tan x} \, dx = \dots\dots\dots + c$

(a)  $e^{\cot x}$

(b)  $e^{1 + \tan x}$

(c)  $e^{1 + \cot x}$

(d)  $e^{\sec^2 x}$

118  $\int \ln x \, dx = \dots\dots\dots + c$

(a)  $\frac{1}{2} (\ln x)^2$

(b)  $x \ln x$

(c)  $x \ln x - x$

(d)  $x \ln x - 1$

119  $\int x \ln x \, dx = \dots\dots\dots + c$

(a)  $x^2 \ln x - x^2$

(b)  $x^2 - \ln x$

(c)  $x - \frac{1}{2} x^2 \ln x^2$

(d)  $\frac{1}{2} x^2 \ln x - \frac{1}{4} x^2$

120 (Trial 2021)  $\int \frac{9}{2} \sqrt{x} \ln x \, dx = \dots\dots\dots + c$

(a)  $x^{\frac{3}{2}} (\ln x^3 - 2)$

(b)  $x^{\frac{3}{2}} (\ln x - 2)$

(c)  $x^{\frac{3}{2}} (\ln x^3 + 2)$

(d)  $x^{\frac{3}{2}} (\ln x + 2)$

121  $\int \frac{1}{\sin^2 x \cos^2 x} \, dx = \dots\dots\dots + c$

(a)  $\tan x - \cot x$

(b)  $\frac{1}{\sin x \cos x}$

(c)  $\frac{9}{\tan^3 x \cot^3 x}$

(d)  $\frac{1}{9} \csc^3 x \sec^3 x$

122  $\int (3x + 2) \sin x \, dx = \dots\dots\dots + c$

(a)  $(3x + 2) \cos x + 3 \sin x$

(b)  $-(3x + 2) \cos x + 3 \sin x$

(c)  $(3x + 1) \cos x + 2 \sin x$

(d)  $-(x + 1) \cos x - 3 \sin x$



123 If  $\int (2x+3) \ln x \, dx = yz - \int z \, dy$ , then  $yz$  equals .....

- (a)  $3x \ln x$  (b)  $(2x+3)$   
 (c)  $\frac{1}{2} (2x+3) \ln x$  (d)  $x(x+3) \ln x$

124 If  $\int (2x-1) e^{2x+3} \, dx = yz - \int z \, dy$ , then  $\int z \, dy = \dots + c$

- (a)  $e^{2x+3}$  (b)  $\frac{1}{2} e^{2x+3}$   
 (c)  $-e^{2x+3}$  (d)  $-\frac{1}{2} e^{2x+3}$

125 If each of  $y, z$  is a function of  $x$ , then  $\int y \, dz + \int z \, dy = \dots$

- (a)  $d(yz)$  (b)  $\int yz \, dx$  (c)  $yz+c$  (d)  $y+z+c$

126  $\int x e^x \, dx = \dots + c$

- (a)  $\frac{1}{2} x^2 e^x$  (b)  $e^x(x-1)$  (c)  $\frac{1}{2} x^2 e^{x+1}$  (d)  $e^x(x+1)$

127  $\int x^3 e^x \, dx = e^x \times (\dots) + c$

- (a)  $\frac{1}{4} x^4$  (b)  $x^3 + 3x^2$   
 (c)  $x^3 - 3x^2 + 6x - 6$  (d)  $\frac{1}{4} x^4 + x^3 + 3x^2$

128  $\int \frac{dx}{e^x + e^{-x} + 2} = \dots + c$

- (a)  $\frac{1}{e^x + 1}$  (b)  $-\frac{1}{e^x + 1}$  (c)  $\frac{2}{e^x + 1}$  (d)  $-\frac{2}{e^x + 1}$

129  $\int x \cos x^2 \, dx = \dots + c$

- (a)  $-\frac{1}{2} \sin^2 x$  (b)  $\frac{1}{2} \sin^2 x$  (c)  $-\frac{1}{2} \sin x^2$  (d)  $\frac{1}{2} \sin x^2$

130  $\int \frac{dx}{1 - \sin x} = \dots + c$

- (a)  $\tan x - \sec x$  (b)  $\tan x + \sec x$   
 (c)  $\sec x - \tan x$  (d)  $\csc x$



131  $\int x \cos x \, dx = \dots\dots\dots + c$

(a)  $x \sin x + \cos x$

(b)  $x \sin x - \cos x$

(c)  $\sin x (x^{-1})$

(d)  $x \sin x + \sin x$

132  $\int \cos^3 x \sin^5 x \, dx = \dots\dots\dots + c$

(a)  $\frac{1}{6} \sin^6 x - \frac{1}{8} \sin^8 x$

(b)  $\sin^6 x - \sin^8 x$

(c)  $\frac{1}{3} \sin^3 x - \frac{1}{6} \sin^6 x$

(d)  $\frac{1}{4} \cos^4 x + \frac{1}{6} \sin^6 x$

133  $\int \frac{\sin(\ln x)}{x} \, dx = \dots\dots\dots + c$

(a)  $-\cos x (\ln x)$

(b)  $\cos(\ln x)$

(c)  $-\cos(\ln x)$

(d)  $\frac{1}{2} [\sin(\ln x)]^2$

134  $\int e^x (\sin x + \cos x) \, dx = \dots\dots\dots + c$

(a)  $e^x \sin x$

(b)  $e^x \cos x$

(c)  $-e^x \sin x$

(d)  $e^{2x} \sin x$

135  $\int e^x (1 + \tan x + \tan^2 x) \, dx = \dots\dots\dots + c$

(a)  $e^x \tan x$

(b)  $e^x \sec x$

(c)  $e^x \sin x$

(d)  $e^x \cos x$

136  $\int e^x (1 - \cot x + \cot^2 x) \, dx = \dots\dots\dots + c$

(a)  $e^x \cot x$

(b)  $-e^x \cot x$

(c)  $e^x \csc x$

(d)  $-e^x \cos x$

137  $\int e^x \left( \frac{1 + x(\ln x)}{x} \right) \, dx = \dots\dots\dots + c$

(a)  $\frac{e^x}{x}$

(b)  $e^x \ln x$

(c)  $x e^x \ln x$

(d)  $\frac{1}{x} + \ln x$

138  $\int e^x (f'(x) + f''(x)) \, dx = e^x \times \dots\dots\dots + c$

(a)  $x$

(b)  $f(x)$

(c)  $f'(x)$

(d)  $f''(x)$

139 If  $\int e^{2x} \left( \ln x + \frac{a}{x} \right) \, dx = \frac{1}{2} e^{2x} \ln x + c$ , then  $a = \dots\dots\dots$

(a)  $\frac{1}{2}$

(b)  $\frac{1}{4}$

(c) 1

(d) 2



140 If  $\sin^2 x \, dy = y \, dx$ , then .....

- (a)  $|y| = e^{-\cot x + c}$       (b)  $|y| = e^{\cos x + c}$       (c)  $|y| = e^{\cot x + c}$       (d)  $|y| = e^{\csc x + c}$

141 If  $I_2 = \int x^2 e^x \, dx$ ,  $I_1 = \int x e^x \, dx$ , then .....

- (a)  $I_2 = I_1$       (b)  $I_2 - I_1 = x e^x$   
 (c)  $I_2 + I_1 = x e^x$       (d)  $I_2 + 2 I_1 = x^2 e^x + c$

142  $\int (1 + 4x^4) e^{x^4} \, dx = \dots + c$

- (a)  $x e^{x^4}$       (b)  $\frac{4}{5} x e^{x^4}$       (c)  $4 x e^{x^4}$       (d)  $(x + \frac{4}{5} x^5) e^{x^4}$

143  $\int e^{2x} \cos x \, dx = \dots + c$

- (a)  $\frac{1}{2} e^{2x} \sin x$       (b)  $2 e^{2x} (\sin x + \cos x)$   
 (c)  $\frac{1}{4} e^{2x} (\sin x + \cos x)$       (d)  $\frac{1}{5} e^{2x} (\sin x + 2 \cos x)$

144  $\int \tan^4 x \, d(\tan x) = \dots + c$

- (a)  $\frac{1}{5} \tan^5 x$       (b)  $\frac{1}{3} \tan^3 x - \tan x$   
 (c)  $\frac{1}{5} \tan^5 x + \tan x$       (d)  $\frac{1}{3} \tan^3 x - \tan x + x$

145  $\int \tan^4 x \, dx = \dots + c$

- (a)  $\frac{1}{5} \tan^5 x$       (b)  $\frac{1}{3} \tan^3 x - \tan x$   
 (c)  $\frac{1}{5} \tan^5 x + \tan x$       (d)  $\frac{1}{3} \tan^3 x - \tan x + x$

146  $\int \sin\left(\frac{\pi}{4} + 2x\right) \sin\left(\frac{\pi}{4} - 2x\right) \, dx = \dots + c$

- (a)  $\frac{1}{8} \sin 4x$       (b)  $\frac{1}{2} \cos 2x$       (c)  $\frac{1}{4} \sin 2x$       (d)  $-\frac{1}{4} \cos 4x$

147  $\int \frac{\sin\left(x + \frac{\pi}{3}\right)}{\sin x} \, dx = \dots + c$

- (a)  $\frac{1}{2} x + \frac{\sqrt{3}}{2} \ln |\cos x|$       (b)  $\frac{1}{2} x + \frac{\sqrt{3}}{2} \ln |\sin x|$   
 (c)  $x + \ln |\tan x|$       (d)  $x + \ln |\sec x|$



148  $\int \frac{x^{e-1} + e^{x-1}}{x^e + e^x} dx = \dots\dots\dots + c$

- (a)  $\frac{1}{e} \ln |x^e + e^x|$   
 (b)  $\ln |x^e + e^x|$   
 (c)  $\ln |x^{e-1} + e^{x-1}|$   
 (d)  $\ln |x + x^e|$

149  $\int \frac{1}{x^3} (\ln x^x)^2 dx = \dots\dots\dots + c$

- (a)  $\frac{1}{3} x^3 (\ln x) + x$  (b)  $\frac{1}{3} (\ln x)^3$   
 (c)  $3 \ln |\ln x|$  (d)  $\frac{1}{x} (\ln x)^2$

150 If  $f(x) = x^3 + 1$ , then  $\int f^{-1}(x) dx = \dots\dots\dots + c$


- (a)  $\frac{4}{3} (x-1) \sqrt[3]{x-1}$   
 (b)  $\frac{4}{3} (x+1) \sqrt[3]{x+1}$   
 (c)  $\frac{3}{4} (x-1) \sqrt[3]{x-1}$   
 (d)  $\frac{3}{4} (x+1) \sqrt[3]{x+1}$

151 In the opposite figure :

If  $m(\angle ADE) = m(\angle C)$

, then  $\int y dx = \dots\dots\dots + c$

- (a)  $\tan x + \frac{1}{3} \tan^3 x$   
 (b)  $\tan x + \frac{1}{2} \tan^2 x$   
 (c)  $\tan x + \frac{1}{9} \tan^3 x$   
 (d)  $\frac{1}{3} \sec^3 x + \frac{1}{3} \tan^3 x$





**Ninth Questions on the definite integration**

Choose the correct answer from the given ones :

1  $\int_a^b f(x) dx = \dots\dots\dots$

- (a)  $f(b) - f(a)$       (b)  $\int_b^a f(x) dx$       (c)  $-\int_b^a f(x) dx$       (d)  $b - a$

2 If  $f$  is continuous on the interval  $[a, b]$ , then  $f(a) + \int_a^b f'(x) dx = \dots\dots\dots$

- (a)  $f'(b)$       (b)  $f(b)$       (c)  $f(a)$       (d)  $f'(a)$

3  $\int_a^b x^2 dx + \int_b^a y^2 dy = \dots\dots\dots$  (where  $a \neq b$ )

- (a)  $2 \int_a^b x^2 dx$       (b) zero      (c)  $2 \int_b^a y^2 dy$       (d)  $b - a$

4 If  $\int_2^4 f(x) dx = 7$ ,  $\int_4^2 g(x) dx = 2$ , then  $\int_2^4 [2f(x) - 3g(x) - 5] dx$  equals  $\dots\dots\dots$

- (a)  $-18$       (b)  $-8$       (c)  $10$       (d)  $14$

5 If  $\int_2^5 f(x) dx = 4$ , then  $\int_2^5 [3f(x) - 1] dx$  equals  $\dots\dots\dots$

- (a)  $9$       (b)  $11$       (c)  $12$       (d)  $-8$

6 If  $f$  is continuous function on the interval  $[2, 7]$ , then  $\int_2^7 f(x) dx + \int_7^4 f(x) dx = \dots\dots\dots$

- (a)  $\int_2^4 f(x) dx$       (b) zero  
(c)  $\int_4^2 f(x) dx$       (d)  $2 \int_2^4 f(x) dx$

7 If  $f$  is a continuous function on  $\mathbb{R}$ ,  $\int_{-1}^3 f(x) dx = 7$ ,  $\int_5^3 f(x) dx = -11$ , then  $\int_{-1}^5 f(x) dx$  equals  $\dots\dots\dots$

- (a)  $-4$       (b)  $18$       (c)  $-18$       (d)  $77$

8 If  $\int_1^4 f(x) dx + \int_2^8 f(x) dx = \int_1^8 f(x) dx$ , then  $b$  may be equal  $\dots\dots\dots$

- (a)  $2$       (b)  $4$       (c)  $1$       (d)  $8$



- 9 If  $f$  is an even function, and is continuous on the interval  $[-4, 4]$   
 $_{-4} \int^4 f(x) dx = 20$ ,  $_0 \int^2 f(x) dx = 6$ , then:  $_{-4} \int^2 f(x) dx = \dots\dots\dots$   
(a) 8 (b) 14 (c) 16 (d) 26
- 10 If  $f$  is an odd function and continuous on  $\mathbb{R}$  and  $_{-6} \int^4 f(x) dx = -3$   
 $_0 \int^6 f(x) dx = 11$ , then  $_0 \int^4 f(x) dx = \dots\dots\dots$   
(a) 5 (b) 6 (c) 8 (d) 10
- 11 If  $_1 \int^3 f(x) dx = 5$ ,  $_3 \int^4 f(x) dx = 2$ ,  $_2 \int^4 f(x) dx = 6$   
then  $_2 \int^1 f(x) dx = \dots\dots\dots$   
(a) 1 (b) 13 (c) -3 (d) -1
- 12 If  $_2 \int^{10} f(x) dx + _8 \int^{11} f(x) dx + _{10} \int^8 f(x) dx = 9$   
then  $_2 \int^{11} f(x) dx = \dots\dots\dots$   
(a) 4.5 (b) 9 (c) 12 (d) 18
- 13 If  $_0 \int^3 f(x) dx = 5$  then  $_0 \int^2 (1 + f(x)) dx + _2 \int^3 f(x) dx = \dots\dots\dots$   
(a) 5 (b) 7 (c) 9 (d) 11
- 14  $_{-1} \int^1 \frac{x^3}{x^4 + \cos x} dx = \dots\dots\dots$   
(a) -1 (b) zero (c) 1 (d) 4
- 15  $_{-2} \int^2 (ax^3 + bx + c) dx$  depends on  $\dots\dots\dots$   
(a) value of  $b$  (b) value of  $c$  (c) value of  $a$  (d) values of  $a, b$
- 16 If  $_{-2} \int^2 f(x) dx = \text{zero}$ , then  $f(x)$  may be  $\dots\dots\dots$   
(a)  $x^2 + 1$  (b)  $x$  (c)  $x + 1$  (d)  $x - 1$
- 17  $\frac{d}{dx} \left( _2 \int^3 x^2 \sqrt{x^2 + 1} dx \right) = \dots\dots\dots$   
(a) -1 (b) zero (c) 1 (d) 2



18  $\int_0^2 (2 - |x|) dx = \dots\dots\dots$

- (a) 4                                      (b) 2                                      (c) 1                                      (d) zero

19  $\int_{-8}^8 \frac{1}{\sqrt{1+|x|}} dx = \dots\dots\dots$

- (a) 6                                      (b) -8                                      (c) -6                                      (d) 8

20  $-\pi \int_{-\pi}^{\pi} (4 + \pi \cos 2x) dx = \dots\dots\dots$

- (a)  $\pi$                                       (b)  $2\pi$                                       (c)  $4\pi$                                       (d)  $8\pi$

21  $\int_0^1 2 \sin \pi z dz = \dots\dots\dots$

- (a)  $4\pi$                                       (b)  $\frac{2}{\pi}$                                       (c)  $\frac{4}{\pi}$                                       (d)  $8\pi$

22  $\int_0^{\frac{\pi}{2}} \cos^2 x dx = \dots\dots\dots$

- (a)  $\frac{\pi}{3}$                                       (b)  $\frac{\pi}{2}$                                       (c)  $\frac{\pi}{4}$                                       (d) zero

23  $\frac{\pi}{6} \int_{\frac{\pi}{6}}^{\frac{\pi}{2}} \frac{\cos x}{\sin^2 x} dx = \dots\dots\dots$

- (a) 0                                      (b) 1                                      (c) 2                                      (d) -1

24  $\int_0^1 x e^{x^2} dx = \dots\dots\dots$

- (a)  $\frac{e+1}{2}$                                       (b)  $\frac{e-1}{2}$                                       (c)  $\frac{e}{2}$                                       (d)  $\frac{1}{2}$

25  $\frac{\pi}{4} \int_{\frac{\pi}{4}}^{\frac{\pi}{2}} \frac{e^{\cot x}}{\sin^2 x} dx = \dots\dots\dots$

- (a) 1                                      (b)  $e-1$                                       (c)  $e+1$                                       (d)  $e$

26  $\int_0^{\frac{\pi}{6}} \sin^4 x dx + \frac{\pi}{6} \int_0^{\frac{\pi}{6}} \cos^4 x dx = \dots\dots\dots$

- (a)  $\frac{\sqrt{3}}{4}$                                       (b)  $-\frac{\sqrt{3}}{4}$                                       (c)  $\frac{1}{4}$                                       (d)  $-\frac{1}{4}$

27  $\int_e^{e^2} \frac{dx}{x} = \dots\dots\dots$

- (a) 1                                      (b) 2                                      (c)  $e$                                       (d)  $e^2$



28  $\int_2^3 x(x-3)^5 dx = \dots\dots\dots$

(a)  $-\frac{1}{6}$

(b)  $-\frac{5}{14}$

(c)  $-\frac{2403}{308}$

(d)  $-\frac{729}{14}$

29  $\int_0^6 |3-x| dx = \dots\dots\dots$

(a) 9

(b) -9

(c)  $\frac{9}{2}$

(d)  $-\frac{9}{2}$

30  $\int_{-3}^3 |x^2-4| dx = \dots\dots\dots$

(a)  $7\frac{2}{3}$

(b) zero

(c)  $15\frac{1}{3}$

(d) -6

31  $\int_1^3 (x-1)|x-2| dx = \dots\dots\dots$

(a)  $\frac{2}{3}$

(b) 1

(c) -1

(d)  $-\frac{2}{3}$

32 If  $\int_2^a (4x-1) dx = 9$ ,  $a \in \mathbb{Z}^+$ , then  $a = \dots\dots\dots$

(a) 3

(b) 4

(c) 1

(d) -3

33 If  $\int_k^3 2x dx = 5$ , then  $k = \dots\dots\dots$

(a) 13

(b) 5

(c) 1

(d)  $\pm 2$

34 If  $a < 2 < b$ , and  $\int_a^b |x-2| dx = 4$ , then  $\frac{a^2+b^2}{a+b} = \dots\dots\dots$

(a)  $\frac{1}{2}$

(b) 4

(c) 2

(d) -4

35 If  $\int_{\ln b}^{\ln a} e^x dx = 2$ ,  $a^2 - b^2 = 12$ , then  $a = \dots\dots\dots$

(a) 8

(b) 12

(c) 4

(d) 6

36  $\int_0^{\ln 3} (e^{2x} + e^x) dx = \dots\dots\dots$

(a) 12

(b) 6

(c)  $7\frac{1}{2}$

(d)  $5\frac{1}{2}$

37  $\int_1^e (\ln x) dx = \dots\dots\dots$

(a)  $\frac{1}{e}$

(b)  $e$

(c) 1

(d) -1

38  $\int_0^{20\pi} |\sin x| dx = \dots\dots\dots$

(a) 20

(b)  $20\pi$

(c) 40

(d)  $40\pi$



39  $\int_0^2 \sqrt{4-x^2} dx = \dots\dots\dots$

- (a) zero (b) 2 (c)  $\pi$  (d)  $\frac{\pi}{2}$

40 If  $f(x) = \begin{cases} 2x-1 & -1 \leq x \leq 2 \\ 3 & 2 < x < 5 \end{cases}$ , then  $\int_{-1}^4 f(x) dx = \dots\dots\dots$

- (a) 4 (b) 5 (c) 6 (d) 7

41 If  $f(x) = \begin{cases} |x-1| & x \leq 1 \\ x^2-1 & x > 1 \end{cases}$ , then  $\int_{-2}^2 f(x) dx = \dots\dots\dots$

- (a)  $\frac{3}{2}$  (b)  $-\frac{35}{6}$  (c)  $\frac{35}{6}$  (d)  $-\frac{3}{2}$

42 If  $\int_0^a |x| dx = 32$ , then  $a = \dots\dots\dots$

- (a) 8 (b) -8 (c)  $\pm 8$  (d) zero

43  $\int_{-1}^1 \sqrt{16-16x^2} dx = \dots\dots\dots$  square units.

- (a)  $16\pi$  (b)  $\pi$  (c)  $2\pi$  (d)  $4\pi$

44  $\int_0^4 4^{\log_2 x} dx = \dots\dots\dots$

- (a)  $\frac{64}{3}$  (b)  $\frac{16}{3}$  (c) 4 (d)  $\frac{11}{3}$

45 If  $\int_k^{k+1} (9)^{\log_3 \sqrt{x}} dx = \frac{5}{2}$ , then  $k = \dots\dots\dots$

- (a) zero (b) 1 (c) 2 (d) 3

46  $\int_{\frac{1}{\pi}}^{\frac{2}{\pi}} \frac{\sin(\frac{1}{x})}{x^2} dx = \dots\dots\dots$

- (a) -2 (b) -1 (c) zero (d) 1

47 If  $\int_0^{\pi} \frac{\cos x}{1+x^{10}} dx = m$ , then  $\int_{-\pi}^{\pi} \frac{3 \cos x}{1+x^{10}} dx = \dots\dots\dots$

- (a) 2m (b) 3m (c) 6m (d) 12m

48  $\int_e^{e^3} \frac{(\ln x)^3}{x} dx = \dots\dots\dots$

- (a) 20 (b) 16 (c) 12 (d) 10



- 49 If  $\int_{-2}^2 (x^7 + k) dx = 16$ , then  $k = \dots\dots\dots$   
(a) -12 (b) -4 (c) zero (d) 4
- 50 If  $f(x)$ ,  $\hat{f}(x)$  are two continuous functions and if  $f(6) = 4$ ,  $\hat{f}(6) = 3$ ,  $f(7) = 14$ ,  $\hat{f}(7) = 5$ , then  $\int_6^7 \hat{f}(x) dx = \dots\dots\dots$   
(a) 10 (b) 2 (c) 18 (d) 8
- 51  $\int_{-1}^0 e^{-x} dx = \dots\dots\dots$   
(a)  $\frac{1}{e} - 1$  (b)  $1 - e$  (c)  $e - 1$  (d)  $1 - \frac{1}{e}$
- 52 If the function  $f$  is continuous,  $f(5) = 9$ ,  $f(1) = 4$ , then  $\int_1^5 3\sqrt{f(x)} \hat{f}(x) dx = \dots\dots\dots$   
(a) 19 (b) 38 (c)  $10\sqrt{3} - 2$  (d) 50
- 53  $\int_0^\pi x^2 \cos x dx = \dots\dots\dots$   
(a)  $-\pi$  (b)  $-2\pi$  (c)  $\pi$  (d)  $2\pi$
- 54 If  $\int_0^\theta \frac{\sec^2 x}{1 + \tan x} dx = \ln 2$  where  $0 < \theta < \frac{\pi}{2}$ , then  $\theta = \dots\dots\dots$   
(a)  $\frac{\pi}{12}$  (b)  $\frac{\pi}{6}$  (c)  $\frac{\pi}{4}$  (d)  $\frac{\pi}{3}$
- 55 If  $(\int_0^a x dx)^3 = \int_0^a x^3 dx$ ,  $a \in \mathbb{R}^+$ , then  $a = \dots\dots\dots$   
(a) 2 (b)  $\sqrt{2}$  (c) 3 (d)  $\sqrt{3}$
- 56 If  $\int_0^k (3x^2 - 1) dx = k^3 - 2$ , then  $k = \dots\dots\dots$   
(a) 6 (b) 4 (c) 2 (d) 1
- 57 If  $\int_{-\frac{\pi}{2}}^a \sin^3 x dx = \text{zero}$ , then  $a = \dots\dots\dots$   
(a) zero (b)  $\pi$  (c)  $-\pi$  (d)  $\frac{\pi}{2}$
- 58 If  $\int_{2k-3}^{k+2} (x^6 + x^4 - 2) dx = \text{zero}$ , then  $k = \dots\dots\dots$   
(a) 2 (b) 3 (c) 4 (d) 5



59 If  $\int_{-1}^a f(z) dz = 4a^3 - 2a + c$ , then  $c = \dots\dots\dots$

(a) -1

(b) 1

(c) 2

(d) -2

60 If  $m, n \in \mathbb{R}$  and  $\int_0^1 (mX + nX) dX = 15$ , then  $\int_0^1 (mX^2 + nX^2) dX = \dots\dots\dots$

(a) 10

(b) 15

(c) 20

(d) 30

61 If  $f$  is a differentiable function on  $\mathbb{R}$  and  $f(1) = 4$ ,  $f(3) = 10$ , then  $\int_1^3 [f(X) + Xf'(X)] dX = \dots\dots\dots$

(a) 22

(b) 24

(c) 26

(d) 28

62  $\pi \int^{2\pi} \sin^2 X dX + 2\pi \int^{\pi} (3 - \cos^2 X) dX = \dots\dots\dots$

(a)  $2\pi$

(b)  $\pi$

(c)  $-\pi$

(d)  $-2\pi$

63 (Trial 2021) If  $a, b \in ]0, \frac{\pi}{2}[$  then  $\int_a^b \tan^2 X dX + \int_b^a \sec^2 X dX = \dots\dots\dots$

(a)  $a - b$

(b)  $b - a$

(c) 1

(d)  $\tan b - \tan a$

64 If  $0 < a < \frac{\pi}{2}$  and  $\int_a^{\frac{\pi}{2}} \frac{\sin X (1 + \cos 2X)}{\sin 2X} dX = \frac{1}{2}$  then  $a = \dots\dots\dots$

(a)  $\frac{\pi}{2}$

(b)  $\frac{\pi}{6}$

(c)  $\frac{\pi}{4}$

(d)  $\frac{\pi}{3}$

65 (2<sup>nd</sup> session 2021) If  $f$  is an even continuous function in  $\mathbb{R}$  and  $\int_0^3 f(X) dX = 5$ , then  $\int_{-3}^3 [f(X) - f(X)] dX = \dots\dots\dots$

(a) 15

(b) -10

(c) -15

(d) 10

66 If  $f(X)$  is defined on  $\mathbb{R}$  and  $f(-X) = f(X)$  and  $\int_{-5}^1 f(X) dX = 24$ ,  $\int_1^5 f(X) dX = 16$ , then  $\int_0^1 f(X) dX = \dots\dots\dots$

(a) -8

(b) 4

(c) 8

(d) 20

67 If  $f$  is an even function and  $\int_{-5}^{-3} f(X) dX = 10$ ,  $\int_0^3 f(X) dX = 8$ , then  $\dots\dots\dots$

(a)  $\int_0^5 f(X) dX = 2$

(b)  $\int_0^5 f(X) dX = -2$

(c)  $\int_{-5}^3 f(X) dX = 26$

(d)  $\int_{-5}^5 f(X) dX = 18$



68  $\int_2^5 \frac{1}{x+2} dx = \dots\dots\dots$

(a)  $\int_0^3 \frac{1}{x+4} dx$

(b)  $-\int_1^2 \frac{1}{x+5} dx$

(c)  $\int_3^6 \frac{1}{x+1} dx$

(d) All the previous

69  $\int_0^2 (x+6)^2 e^{x^2} dx - \int_0^2 (x-6)^2 e^{x^2} dx = \dots\dots\dots$

(a)  $12e^2$

(b)  $12e^4$

(c)  $12(e^4 - 1)$

(d)  $24(e^4 - 1)$

70 If  $\int_{\sin \theta}^{\cos \theta} \frac{x}{2} dx = \frac{-\sqrt{3}}{8}$ , then one of the values of  $\theta$  equals  $\dots\dots\dots$

(a)  $30^\circ$

(b)  $45^\circ$

(c)  $60^\circ$

(d)  $75^\circ$

71  $\int_{2019}^{2021} (x-2020)^2 dx = \dots\dots\dots$

(a)  $\frac{1}{3}$

(b)  $\frac{2}{3}$

(c) 1

(d)  $\frac{4}{3}$

72 If  $\int_3^9 f(x) dx = 20$ ,  $\int_5^{11} f(x) dx = 25$ , then  $\int_3^5 f(x) dx - \int_9^{11} f(x) dx = \dots\dots\dots$

(a) -10

(b) -5

(c) zero

(d) 5

73 If  $\int_a^b (3x^2 + 2) dx = 96$ ,  $\int_a^b dx = 4$ , then  $a \times b = \dots\dots\dots$

(a) 1

(b) 1.5

(c) 2

(d) 2.5

74 If  $f(x) = \frac{3x^{2021} + 1}{x^{2023} + 1}$ , then  $\int_0^1 f(x) dx = \dots\dots\dots$

(a) zero

(b) 1

(c) 2

(d) 3

75  $\lim_{h \rightarrow 0} \left[ \frac{1}{h} \int_3^{3+h} \sqrt{x^2 + 16} dx \right] = \dots\dots\dots$

(a) 3

(b) 5

(c) 9

(d) 16

76 If  $\int_0^a (3x^2 - 2x) dx \leq 2$  where  $a \in \mathbb{R}^+$ , then  $a \in \dots\dots\dots$

(a)  $[0, 2]$

(b)  $]0, 2]$

(c)  $]2, \infty[$

(d)  $]1, \infty[$



77 If  $f$  is a function where  $\int_6^{12} f(2x) dx = 10$ , then which of the following is correct?

(a)  $\int_{12}^{24} f(t) dt = 5$

(b)  $\int_{12}^{24} f(t) dt = 20$

(c)  $\int_6^{12} f(t) dt = 5$

(d)  $\int_6^{12} f(t) dt = 20$

78 If  $f$  is a differentiable twice function in the interval  $[a, b]$  and the slope of the tangent of the function  $f$  at  $x = a$  equals 1 and the slope of the tangent of the function  $f$  at  $x = b$  equals  $\sqrt{2}$ , then  $\int_a^b 2f'(x)f''(x) dx = \dots\dots\dots$

(a) -2

(b) -1

(c) 1

(d) 2

79 If  $k$  is one of the roots of the equation:  $x^4 - x^2 - 6 = 0$  then  $\int_{-1}^k (4x^3 - 2x) dx = \dots\dots\dots$

(a) 3

(b) 4

(c) 5

(d) 6

80  $-\frac{\pi}{2} \int_{\frac{\pi}{2}}^{\pi} \ln\left(\frac{2 - x \cos x}{2 + x \cos x}\right) dx = \dots\dots\dots$

(a)  $\ln 1$

(b)  $\ln 2$

(c)  $\ln e$

(d)  $\ln \pi$

81 If  $I_1 = \int_a^b |f(x)| dx$ ,  $I_2 = \left| \int_a^b f(x) dx \right|$  then  $\dots\dots\dots$

(a)  $I_1 = I_2$

(b)  $I_1 \geq I_2$

(c)  $I_1 \leq I_2$

(d) can not be determined.

82 If  $f$  is continuous on the interval  $[0, \pi]$  and integrable,  $\int_0^{\pi} f(x) dx = 2 \int_0^{\frac{\pi}{2}} f(x) dx$ , then  $\dots\dots\dots$

(a)  $f(x)$  is an odd function.

(b)  $f(x)$  is an even function.

(c) The curve of  $f(x)$  is symmetric about the point  $\left(\frac{\pi}{2}, 0\right)$

(d) The curve of  $f(x)$  is symmetric about the straight line  $x = \frac{\pi}{2}$


**83 In the opposite figure :**the curve of the function  $y = f(x)$ , then  $\int_0^2 f'(x) dx = \dots\dots\dots$ 

(a) 12


(b) -2

(c) -4

(d) 4

**84 In the opposite figure :** $\int_0^2 [f(x)]^2 f'(x) dx = \dots\dots\dots$ (a)  $-\frac{64}{3}$ (b)  $\frac{64}{3}$ (c)  $\frac{8}{3}$ 

(d) -64

**85 In the opposite figure :**

The straight line L is a tangent to the curve


 $y = f(x)$  at the point A (1, 2), then  $\int_{-3}^1 f'(x) dx = \dots\dots\dots$ 

(a) -3


(b) -2

(c) -1

(d) 1

**86 In the opposite figure :** $\int_{-4}^3 \frac{f'(x)}{f(x)} dx = \dots\dots\dots$ 

(a) 1

(b)  $\ln \frac{3}{2}$ (c)  $\ln \frac{2}{3}$ (d)  $\log_3 2$ 



87 In the opposite figure :

If L is the tangent to the curve  $y = f(x)$  at the point

A (3, 3), then


$$\int_0^3 x f'(x) dx = \dots\dots\dots$$

(a) -2

(b) 2

(c) 4

(d) 8



88 The opposite figure represents the curve of the function  $y = f(x)$  and  $g(x) = x \cdot f(x)$


$$\text{, then } \int_1^3 g'(x) dx = \dots\dots\dots$$

(a) -1

(b) -2

(c) -3

(d) -4



89 In the opposite figure :

If ABCD is a square

$$\text{, } m(\angle AED) = 90^\circ$$


$$\text{, then } \int_0^1 (EC)^2 dx = \dots\dots\dots$$

(a)  $\frac{82}{3}$

(b)  $\frac{91}{3}$

(c)  $\frac{100}{3}$

(d)  $\frac{109}{3}$



**Tenth Questions on the applications of integration****Choose the correct answer from the given ones :**

- 1 If  $\dot{f}(x) = 1 - \sin 2x$ ,  $f(0) = \frac{1}{2}$ , then  $f\left(\frac{\pi}{2}\right) = \dots\dots\dots$   
 (a)  $\frac{\pi}{2} - \frac{1}{2}$  (b)  $-\frac{1}{2}$  (c) 1 (d)  $\frac{\pi}{2}$
- 
- 2 If  $\frac{dy}{dx} = \csc^2 x$ ,  $y = 2$  at  $x = \frac{\pi}{4}$ , then  $y = \dots\dots\dots$   
 (a)  $-2 - \cot x$  (b)  $2 - \cot x$  (c)  $-3 - \cot x$  (d)  $3 - \cot x$
- 
- 3 If  $\frac{dy}{dx} = x + \frac{1}{x}$ ,  $y = \frac{1}{2}$  at  $x = 1$ , then  $y = \dots\dots\dots$  when  $x = e$   
 (a)  $\frac{1}{2}e^2 + 1$  (b)  $e + \frac{1}{e}$  (c)  $e^2$  (d)  $e^2 + 1$
- 
- 4 If  $f(x) = \int \frac{dx}{\sqrt{2x+1}}$ ,  $f(4) = 7$ , then  $f(x) = \dots\dots\dots$   
 (a)  $\sqrt{2x+1}$  (b)  $2\sqrt{2x+1} + 6$  (c)  $\frac{1}{2}\sqrt{2x+1}$  (d)  $\sqrt{2x+1} + 4$
- 
- 5 If  $f(x) = \int (x+1)(2x^2 + 4x - 1) dx$ ,  $f(-2) = 1$ , then  $f(3) = \dots\dots\dots$   
 (a) 54 (b) 86 (c) 98 (d) 106
- 
- 6 If  $\int f(x) dx = x^3 - x^2$ , then  $\dot{f}(1) - f(1) = \dots\dots\dots$   
 (a) 4 (b) 3 (c) zero (d) 2
- 
- 7 If  $\int \frac{f(x)}{x} dx = \ln|x| + x^2 + c$ , then  $f(x) = \dots\dots\dots$   
 (a)  $x^2 + x + 1$  (b)  $x^3 + x^2 + 1$  (c)  $2x^2 + 1$  (d)  $x^2 + 1$
- 
- 8 The equation of the curve which intercepted 7 units, from negative direction of y-axis and slope of its tangent at any point on it  $= 3x^2 + 2$ , is  $\dots\dots\dots$   
 (a)  $y = x^3$  (b)  $y - x^3 - 2x = 0$   
 (c)  $y = x^3 - 7$  (d)  $y = x^3 + 2x - 7$
- 
- 9 If  $f(x) = \sin x + \cos x$ , then  $\dot{f}(x) + \int f(x) dx = \dots\dots\dots$   
 (a) zero (b) constant (c)  $2 \sin x$  (d)  $2 \cos x$



- 10 If  $f(x) = \int \frac{1}{x} dx$ , then  $f'(2) = \dots\dots\dots$   
 (a) does not exist (b)  $\frac{1}{x} + c$  (c) 2 (d)  $\frac{1}{2}$
- 
- 11 If  $\int f(x) dx = g(x) + c_1$ ,  $\int g(x) dx = \ln \frac{1}{x} + c_2$ , then  $f(2) + g(2) = \dots\dots\dots$   
 (a)  $-\frac{1}{4}$  (b)  $-\frac{1}{2}$  (c) 2 (d) 4
- 
- 12 (1<sup>st</sup> session 2021) If  $x \frac{dy}{dx} - y = 3$  and  $x = 1$  when  $y = -2$ , then the relation between  $x, y$  is  $\dots\dots\dots$   
 (a)  $|y| = |x + 3|$  (b)  $|y + 3| = |x|$  (c)  $y + x = -1$  (d)  $y + 2x = 0$
- 
- 13 (2<sup>nd</sup> session 2021) If  $3y^2 \frac{dy}{dx} = 2x + 1$  and  $x = 1$  when  $y = -1$ , then the relation between  $x, y$  is  $\dots\dots\dots$   
 (a)  $y^3 = x^2 + x - 3$  (b)  $y^3 = x^2 - 2$   
 (c)  $y^3 + x^2 = 0$  (d)  $y^3 + 1 = x^2 - x$
- 
- 14 If  $f'(x) = \ln x$  and  $f(1) = 1$ , then  $f(2) = \dots\dots\dots$   
 (a)  $\ln \sqrt{2}$  (b)  $\ln 2$  (c)  $\ln 4$  (d)  $\ln 8$
- 
- 15 (Trial 2021) If  $f(x) = \int \frac{1 - (\ln x)^2}{x} dx$  where  $f(1) = 0$ , then  $f(e) = \dots\dots\dots$   
 (a)  $-\frac{2}{3}$  (b)  $-\frac{1}{3}$  (c)  $\frac{1}{3}$  (d)  $\frac{2}{3}$
- 
- 16 If  $\frac{dy}{dx} = y \cos x$  and  $y = 1$  when  $x = 0$  then  $\dots\dots\dots$   
 (a)  $y = e^{\sin x}$  (b)  $y = e^{\sin x} + 1$  (c)  $y \sin x = 1$  (d)  $y \sec x = 1$
- 
- 17 If  $f''(x) = e^x$  and  $f'(0) = 3$ ,  $f(0) = 5$ , then  $f(x) = \dots\dots\dots$   
 (a)  $e^x + 2x + 4$  (b)  $e^x + 3x + 4$  (c)  $e^x + 4$  (d)  $e^x + x^2 + 4$
- 
- 18 If slope of the tangent to curve of the function  $y = f(x)$  at  $x = 1$  equals  $-\frac{1}{2}$  and  $f''(x) = x + 1$ , then  $f'(2) = \dots\dots\dots$   
 (a) -2 (b) -1 (c) 1 (d) 2



- 19 If  $\ddot{f}(x) = 6x - 4$ ,  $\dot{f}(1) = 2$ ,  $f(0) = -4$ , then  $f(x) = \dots\dots\dots$
- (a)  $-2x^2 + 3x - 4$  (b)  $x^3 - 2x^2 + 3x$   
 (c)  $3x^2 - 4x - 4$  (d)  $x^3 - 2x^2 + 3x - 4$
- 
- 20 If  $\ddot{f}(x) = \frac{1}{2}[e^x + e^{-x}]$ ,  $f(0) = 1$ ,  $\dot{f}(0) = 0$ , then  $f(x) = \dots\dots\dots$
- (a)  $-f'(x)$  (b)  $f'(x)$  (c)  $-\ddot{f}(x)$  (d)  $\ddot{f}(x)$
- 
- 21 If the slope of the tangent to a curve at any point on it  $(x, y)$  equals  $4e^{2x}$ ,  $f(0) = 2$ , then  $f(-2) = \dots\dots\dots$
- (a) 4 (b)  $4e^{-4}$  (c)  $2e^{-4}$  (d)  $2e$
- 
- 22 If the slope of the tangent to the curve:  $y = f(x)$  at any point on it equals  $6x + a$ , where  $a$  is constant, if the equation of the tangent to the curve at the point  $(1, -1)$  is  $y = 4 - 5x$ , then the equation of the curve is  $\dots\dots\dots$
- (a)  $y = 3x^2 - 4$  (b)  $y = 3x^2 - 11x + 7$   
 (c)  $y = 3x^2 - 4x + 7$  (d)  $y = 3x^2 - 11x - 4$
- 
- 23 If the slope of the tangent to the curve  $y = f(x)$  at any point on it equals  $\sec^2 x - \sin x$ , the curve passes through the point  $(\frac{\pi}{4}, \frac{1}{\sqrt{2}})$ , then the equation is  $\dots\dots\dots$
- (a)  $y = \frac{1}{3} \sec^3 x + \cos x - 1$  (b)  $y = \tan x - \cos x - 1$   
 (c)  $y = \tan x + \cos x - 1$  (d)  $y = \tan x + \cos x$
- 
- 24 If the slope of the tangent to the curve of the function  $f$  at any point on it  $(x, y)$ , equals  $\frac{5x+3}{x}$ , and the curve passes through the point  $(e, 5e + 7)$ , then the equation of the curve is  $\dots\dots\dots$
- (a)  $y = 5x + 3 \ln |x| + 4$  (b)  $y = 5x + \ln |x| + 6$   
 (c)  $y = 5x + 3 \ln |x| - 3$  (d)  $y = x + 7 \ln |x| + 4e$



25 The equation of the curve passes through the point  $(1, 0)$  and the slope of its tangent at any point on it equals  $X e^y$  is .....

(a)  $\frac{1}{2} X^2 + e^y = 1 \frac{1}{2}$

(b)  $e^{-y} + \frac{1}{3} X^3 = 1 \frac{1}{3}$

(c)  $e^{-y} + \frac{1}{2} X^2 = 1 \frac{1}{2}$

(d)  $X + e^y = 2$

26 If the slope of the tangent to the curve of the function  $y = f(X)$  at any point on it equals  $\frac{1}{X\sqrt{3+\ln X}}$  where  $X > 0$  and the curve passes through  $(e, 4)$ , then the relation between  $X, y$  is .....

(a)  $y = \sqrt{3 + \ln X}$

(b)  $y = 2\sqrt{3 + \ln X}$

(c)  $y = 3 + \ln X$

(d)  $y = 2\sqrt{3 + \ln X} - 5$

27 If the slope of the tangent to the curve of the function  $f$  at any point  $(X, y)$  lying on it is given by the relation  $g(X) = X e^{-X}$  and the curve passes through the point  $(-1, 3)$ , then the equation of the curve is .....

(a)  $f(X) = -e^{-X}(X+1) + 3$

(b)  $f(X) = -X e^{-X} + X - e$

(c)  $f(X) = -e^X(X^2 + 1)$

(d)  $f(X) = e^{-X}(X+1) + 3$

28 If the slope of the tangent to a curve at any point on it  $(X, y)$  equals  $X(\sqrt{X+1})$  and the curve passes through  $(0, \frac{11}{15})$ , then the equation of the curve is  $y = \dots\dots\dots$

(a)  $\frac{X}{\sqrt{X+1}} + \frac{11}{15}$

(b)  $2\sqrt{X+1} - \frac{19}{15}$

(c)  $\frac{2}{5}(X+1)^{\frac{5}{2}} - \frac{2}{3}(X+1)^{\frac{3}{2}} + 1$

(d)  $\frac{2}{5}(X+1)^{\frac{5}{2}} + \frac{1}{3}$

29 The slope of the tangent to the curve of a function  $f$  equals  $\frac{1}{X-2}$  and the curve passes through the point  $(3, 0)$ , then  $f(e^2 + 3) = \dots\dots\dots$

(a)  $e^2 + 1$

(b)  $\ln(e^2 + 1)$

(c)  $\frac{1}{e^2 + 1}$

(d)  $2$

30 The slope of the normal to the curve at any point on it  $(X, y)$  equals  $3 - 2X$ , then the equation of the curve given it passes through the point  $(1, 1)$  is .....

(a)  $y = \frac{1}{2} \ln |2X - 3| + 2$

(b)  $y = \frac{1}{2} \ln |2X - 3| + 1$

(c)  $y = \ln |2X - 3| + 4$

(d)  $y = \ln |2X - 3| + 1$



- 31 If the slope of the tangent to the curve of the function  $f$  at any point  $(X, y)$  lying on it is given by the relation  $g(X) = \frac{Xe^X}{(X+1)^2}$ , then the equation of the curve if it passes through the point  $(1, 2e)$  is .....

(a)  $y = \frac{-Xe^X}{X+1} + \frac{1}{2}e$

(b)  $y = \frac{-Xe^X}{X+1} + e^X + \frac{3}{2}e$

(c)  $y = \frac{Xe^X}{X+1} + e^X - \frac{3}{2}e$

(d)  $y = \frac{Xe^X}{X+1} - \frac{1}{2}e$

- 32 If the curve of the function  $y = f(X)$  passes through the point  $(0, 2)$ ,  $\frac{dy}{dX} = \frac{-X}{ye^{X^2}}$  and  $\frac{dy}{dX} > 0$  for all values of  $X \in \mathbb{R}$ , then  $f(X) = \dots\dots\dots$

(a)  $3 + e^{-X^2}$

(b)  $\sqrt{3 + e^{-X^2}}$

(c)  $\sqrt{3 + e^{-X^2}}$

(d)  $\sqrt{3 + e^{X^2}}$

- 33 If the slope of the tangent to the curve at any point on it is given by the relation  $\frac{dy}{dX} = 2Xy$  and  $y = 1$  when  $X = 1$  then the equation of this curve is .....

(a)  $y = 2e^{X-1}$

(b)  $y = e^X$

(c)  $y = e^{X^2-1}$

(d)  $y = e^{X^2+1}$

- 34 If the slope of the tangent to the curve :  $y = f(X)$  at any point on it  $(X, y)$  equals :  $\frac{3+4X}{6y}$ , then the equation of the curve, if we know that it passes through the point  $(1, 1)$  is .....

(a)  $3y^2 = 3X + 2X^2$

(b)  $y^2 = 2X^2$

(c)  $3y^2 = 3X - 2$

(d)  $3y^2 = 3X + 2X^2 - 2$

- 35 If the slope of the normal to the curve  $y = f(X)$  at any point on it is  $(2y + 1) \csc X$  if we know that the curve passes through the origin point, then its equation is .....

(a)  $y^2 + y = \sin X - 1$

(b)  $y^2 + y = \cos X - 1$

(c)  $y^2 + y \csc y \cot y = 0$

(d)  $y^2 + y = (\sin X)^{-2}$

- 36 The slope of the tangent at any point  $(X, y)$  on the curve  $y = f(X)$  is equal to  $3X^2 - 6X - 9$  and the local maximum value of the function  $f$  is 17, then the local minimum value of the function  $f$  equals .....

(a) -17

(b) -15

(c) 7

(d) 15



- 37 If  $y = f(x)$ ,  $\frac{d^2 y}{dx^2} = ax + b$  where  $a$  and  $b$  are two constants and the curve has an inflection point at the point  $(0, 2)$  and a local minimum value at the point  $(1, 0)$ , then the local maximum value to this curve = .....
- (a) 3 (b) 4 (c) 5 (d) 6
- 
- 38 If  $y = f(x)$ , and  $\frac{d^2 y}{dx^2} = \frac{2}{x^3}$  and the equation of its tangent at the point  $(2, \frac{5}{2})$  which lies on the curve is:  $3x - 4y + 4 = 0$ , then the equation of the curve is .....
- (a)  $y = x^2 + 2$  (b)  $y = \frac{1}{x} + x$  (c)  $y = \frac{6}{x} + x$  (d)  $y = x - \frac{1}{x}$
- 
- 39 If  $y = f(x)$ , and:  $\frac{d^2 y}{dx^2} = 6(1 - x)$  and the curve has a local minimum value at the point  $(0, -6)$ , then the equation of the curve is .....
- (a)  $y = x^3 - 3x^2$  (b)  $y = 6x - 3x^2 - 6$   
 (c)  $y = 3x^2 - x^3 - 6$  (d)  $y = x^3 + 6x^2 - 6$
- 
- 40 (2<sup>nd</sup> session 2021) If  $y = f(x)$ ,  $\frac{d^2 x}{dy^2} = e^{y+1}$  and the tangent to the curve of the function  $f$  at the point  $(x, -1)$  which lies on it is parallel to the straight line:  $y = x - 3$ , then the slope of the tangent to the curve of the function  $f$  at any point  $(x, y)$  lies on it equals .....
- (a)  $e^{y+1}$  (b)  $e^{2y+2}$  (c)  $e^{-y-1}$  (d)  $e^{-2y-2}$
- 
- 41 (1<sup>st</sup> session 2021) If  $P = f(n)$  where  $P, n \in \mathbb{R}^+$  and the rate of change of  $P$  with respect to  $n$  varies inversely with  $n$ ,  $f(1) = 200$  and  $f(\sqrt[3]{e}) = \frac{700}{3}$ , then  $f(3) = \dots\dots\dots$
- (a)  $100 \ln(3e^2)$  (b)  $200 \ln(9e)$   
 (c)  $100 \ln(e^2 + 3)$  (d)  $200 \ln(9 + e)$
- 
- 42 If the slope of the tangent at any point  $(x, y)$  on the curve of the function  $f$  is inversely proportional to  $x$  and the slope of the tangent equals 2 when  $x = 4$  and  $y = 2$ , then  $y = \dots\dots\dots$
- (a)  $8 \ln|x| + 2$  (b)  $8 \ln|x| + 2 - 8 \ln 4$   
 (c)  $x^{-2} + \frac{1}{32}$  (d)  $4 \ln|x| - 2$



- 43 If the rate of change of the slope of the tangent to a curve at any point on it is equal to  $6x - 2$  and the slope of the tangent at the point  $(3, 1)$  that lies on the curve equals 2, then the equation of this curve is .....
- (a)  $y = x^3 - x^2 - 18$   
(b)  $y = x^3 - x^2 - 2x - 12$   
(c)  $y = 3x^2 - 2x - 19$   
(d)  $y = x^3 - x^2 - 19x + 40$
- 
- 44 The capacity of an empty vessel is  $1400 \text{ cm}^3$ , water is poured in it at a rate  $(2t + 50) \text{ cm}^3/\text{sec}$ , where  $t$  is the time in seconds, then need the time to fill the vessel = ..... sec.
- (a) 28 (b) 20  
(c) 70 (d) 700
- 
- 45 A liquid is leaking from a small hole in the bottom of a vessel filled with the liquid. If the volume of the liquid changes at the rate of  $(0.4t - 40) \text{ cm}^3/\text{sec}$ , where  $t$  represents the time in seconds and the volume of the liquid is  $980 \text{ cm}^3$  after 30 seconds from the start of leaking, then the capacity of the vessel = .....  $\text{cm}^3$ .
- (a) 1 000 (b) 2 000  
(c) 3 000 (d) 4 000
- 
- 46 If the rate of change of area of a lamina is  $A$  (in square centimeters), with respect to  $t$  (in seconds) by the relation  $\frac{dA}{dt} = e^{-0.2t}$ , if the area of the lamina at the beginning of changing is  $140 \text{ cm}^2$ , then area of the lamina after  $\frac{1}{3}$  minutes equals .....  $\text{cm}^2$ .
- (a) 145 (b)  $145 - 5e^{-4}$   
(c)  $145 - e^{-\frac{1}{5}}$  (d)  $145 - e^{-\frac{1}{15}}$



**Eleventh Questions on the areas and the volumes of revolution solids****Choose the correct answer from the given ones :**

- 1 The area bounded by the straight line :  $y = x$  ,  $x = 1$  ,  $y = 0$  equals .....  
 (a)  $\frac{1}{2}$  (b) 2 (c) 1 (d)  $\frac{1}{4}$

---

- 2 The area of the planar region bounded by the curve :  $y = x^2$  and the two straight line  $y = 0$  ,  $x = 3$  equals .....  
 (a) 6 (b) 7 (c) 8 (d) 9

---

- 3 The area of planar region bounded by the curve :  $y = x^3$  and the straight lines :  $x = -1$  ,  $x = 1$  ,  $y = 0$  equals .....  
 (a) zero (b)  $\frac{1}{2}$  (c)  $\frac{1}{4}$  (d) 6

---

- 4 The area of the planar region bounded by the curve :  $y = x^2 + 4$  and  $x$ -axis and the two straight lines  $x = -1$  ,  $x = 2$  equals .....  
 (a) 15 (b) 9 (c)  $14\frac{1}{3}$  (d)  $12\frac{1}{3}$

---

- 5 The area of the region bounded by the curve :  $y = 2x - x^2$  and  $x$ -axis equals .....  
 (a)  $\frac{8}{3}$  (b)  $\frac{4}{3}$  (c)  $\frac{7}{3}$  (d)  $\frac{3}{4}$

---

- 6 The area of the planar region bounded by the two curves :  $y = x^2$  ,  $y = x^3$  is ..... square unit.  
 (a) 1 (b)  $\frac{7}{12}$  (c)  $\frac{1}{12}$  (d) 2

---

- 7 The area of the planar region bounded by the two curves :  $y^2 = x$  ,  $y = x^3$  equals .....  
 (a)  $\frac{5}{12}$  (b)  $\frac{5}{6}$  (c)  $\frac{12}{5}$  (d)  $\frac{6}{5}$

---

- 8 The area of the planar region bounded by the two curves :  $y^3 = x$  ,  $y = x$  equals .....  
 (a)  $\frac{1}{2}$  (b)  $\frac{1}{4}$  (c)  $\frac{3}{4}$  (d)  $\frac{1}{2}$

---

- 9 The area of the planar region bound by the curves :  $y = x^2 - 2x + 1$  ,  $y = x + 1$  equals .....  
 (a)  $-\frac{9}{2}$  (b)  $\frac{9}{2}$  (c)  $\frac{3}{2}$  (d)  $-\frac{3}{2}$



- 10 The area of the planar region bounded by the curve :  $y = \sqrt{x-1}$  and the straight line  $y = x - 3$  and  $x$ -axis equals .....
- (a)  $3\frac{1}{3}$  (b) zero (c)  $5\frac{1}{3}$  (d) 2
- 
- 11 If  $f$  is a continuous function on the interval  $[a, b]$  and  $A$  is the area bounded by the curve of the function  $y = f(x)$ , the  $x$ -axis and the two straight lines  $x = a$ ,  $x = b$ , then  $A = \dots\dots\dots$
- (a)  $\int_a^b |y| dx$  (b)  $|\int_a^b y dx|$  (c)  $\int_a^b y dx$  (d)  $\int_a^b |x| dy$
- 
- 12 The area of the region bounded by the curve  $xy = 4$  and  $x$ -axis and the two straight lines  $x = 1$ ,  $x = 3$  equals .....
- (a)  $2 \ln 3$  (b)  $4 \ln 3$  (c)  $3 \ln 3$  (d)  $3 \ln 4$
- 
- 13 If the area bounded by the curve  $y = x^3$  and the two straight lines  $y = 0$ ,  $x = a$  where  $a \in \mathbb{R}^+$  equals 4 square units, then  $a = \dots\dots\dots$
- (a) 8 (b) 4 (c) 2 (d) 1
- 
- 14 If the area of the region between the curve  $y = x^2$  and the straight line  $y = kx$  (where  $k > 0$ ) equals 4.5 square units then  $k = \dots\dots\dots$
- (a)  $\frac{3}{2}$  (b) 3 (c)  $\frac{9}{2}$  (d) 2
- 
- 15 The area of the region bounded by the curve  $(x-2)^2 = y-1$  and its tangent at the point  $(3, 2)$  and the  $y$ -axis equals ..... area units.
- (a) 4 (b) 6 (c) 9 (d) 12
- 
- 16 The area of the region bounded by the curve  $x = 2 - y^2$  and the  $y$ -axis equals ..... square units
- (a)  $\frac{2\sqrt{2}}{3}$  (b)  $\frac{4\sqrt{2}}{3}$  (c)  $\frac{8\sqrt{2}}{3}$  (d)  $\frac{16\sqrt{2}}{3}$
- 
- 17 (2<sup>nd</sup> session 2021) The area of the region bounded by the curve  $y - 1 = e^x$  and  $x$ -axis and the straight lines  $x = 0$ ,  $x = k$  where  $k > 0$  equals ..... square unit
- (a)  $e^k - k + 1$  (b)  $e^k + k + 1$  (c)  $e^k + k - 1$  (d)  $e^k - k - 1$



18 (2<sup>nd</sup> session 2021) The area bounded by the curve  $y = f(x)$  where  $y = n - \frac{n^2}{4}$ ,  $x = \frac{n}{2}$  and  $x$ -axis equals ..... square unit.

(a)  $\frac{7}{3}$

(b)  $\frac{3}{4}$

(c)  $\frac{4}{3}$

(d)  $\frac{3}{7}$

19 The area of the region bounded by the curve whose parametric equations  $y = 3t^2$ ,  $x = 6t$  and the  $x$ -axis and the two straight lines  $x = 0$ ,  $x = 12$  equals ..... square units.

(a) 48

(b) 96

(c) 132

(d) 192

20 The area of the region bounded by the curve  $y = \sqrt{4 - x^2}$  and  $x$ -axis estimated by square units equals .....

(a) 2

(b) 4

(c)  $2\pi$

(d)  $4\pi$

21 An architect has designed an arc -like entryway of a hotel whose equation  $y = -\frac{1}{2}(x-1)(x-7)$  where  $x$  in metres. How much does the glass cost if this entryway is covered by the glass which costs L.E. 1 500 per square metre = L.E. ....

(a) 9 000

(b) 27 000

(c) 54 000

(d) 63 000

22 An advertising company produces a poster to market an item. If the poster is shaped as an area bounded by the curve of the two functions  $f$  and  $g$  where  $f(x) = 2x^2$  and  $g(x) = x^4 - 2x^2$ , then the area needed of adhesive paper to produce 1 000 posters for this item = ..... square units.

(a)  $\frac{25\,600}{9}$

(b)  $\frac{12\,800}{3}$

(c)  $\frac{25\,600}{3}$

(d)  $\frac{12\,800}{9}$

23 In the opposite figure :


If  $f(x) = x^2 - 4x + 3$  then the area of the shaded region = ..... square units.

(a)  $\frac{8}{3}$

(b) 4

(c)  $\frac{16}{3}$

(d) 8






24 In the opposite figure :

represents the curve of a quadratic function  
then the area of the shaded region  
= ..... square units.


- (a)  $\frac{32}{3}$
- (b)  $\frac{40}{3}$
- (c)  $\frac{64}{3}$
- (d)  $\frac{80}{3}$



25 In the opposite figure :

The area of the shaded region = ..... square units.


- (a)  $\frac{2}{3}$
- (b)  $\frac{4}{3}$
- (c)  $\frac{5}{3}$
- (d) 2



26 In the opposite figure :

The area of the shaded region = ..... square units.

- (a) 4
- (b) 8
- (c) 12
- (d) 16





27 In the opposite figure :


The area of the shaded region = ..... square units.

(a) 20

(b)  $\frac{45}{2}$

(c) 25

(d)  $\frac{55}{2}$



28 In the opposite figure :


If  $f(x) = \frac{1}{x}$ , then the area of the shaded region = ..... square units.

(a) 1

(b) e

(c)  $e^2$

(d) 2



29 In the opposite figure :


If  $f(x) = 3x^2 + 2$  and the area of the shaded region = 33 square units, then  $k =$  .....

(a) 1

(b) 2

(c) 3

(d) 4




30 The opposite figure represents a quadratic function, its vertex is  $(k, 9)$ , then the area of the shaded region = ..... square units.

(a) 6

(b) 9


(c) 12

(d) 18



**31 In the opposite figure :**

The area of the region bounded by the two curves :  $y = x^2$  and  $y = |x|$  equals = .....



(a)  $2 \int_{-1}^0 (x^2 - x) dx$


(b)  $\int_0^1 (x - x^2) dx$

(c)  $2 \int_0^1 (x - x^2) dx$

(d)  $\int_{-1}^1 (x - x^2) dx$

**32 In the opposite figure :**

The area of the region bounded by the curve  $y = x^3$  and the straight line  $y = x$  equals .....



(a)  $\int_{-1}^1 (x^3 - x) dx$


(b)  $2 \int_0^1 (x^3 - x) dx$

(c)  $\int_0^1 (x - x^3) dx$

(d)  $2 \int_0^1 (x - x^3) dx$

**33 In the opposite figure :**

The area of the shaded region = ..... square units.



(a) 1


(b)  $\frac{1}{2}$

(c) 2

(d)  $\frac{3}{2}$

**34 In the opposite figure :**

The area of the shaded region = ..... square units.



(a) 2

(b)  $\frac{7}{3}$

(c) 3

(d)  $\frac{10}{3}$



35 In the opposite figure :


If the region A between the curve  $f(x)$  and the  $x$ -axis equals 8 square units, then  $\int_0^5 (1 - f(x)) dx = \dots\dots\dots$

(a) 12

(b) 13

(c) 14

(d) 15




36 The opposite figure represents the curve of the function  $f : f(x) = -x(x-4)$ , then all of the following are true except  $\dots\dots\dots$

(a)  $\int_0^1 f(x) dx = \int_3^4 f(x) dx$

(b)  $\int_0^4 f(x) dx = 2 \int_0^2 f(x) dx$

(c)  $\int_{-1}^0 |f(x)| dx = \int_0^1 f(x) dx$

(d)  $\int_{-1}^0 f(x) dx = \int_4^5 f(x) dx$




37 The shaded area in the figure is equal to  $\dots\dots\dots$

(a)  $\pi + 2$

(b)  $\pi - 2$

(c) 2

(d)  $\pi$




38 The opposite figure represents the curve of the function  $f$ , then the area included between the curve of the function  $f$  and  $x$ -axis equals  $\dots\dots\dots$

(a)  $\int_0^{x_2} f(x) dx$

(b)  $\int_0^{x_1} f(x) dx + \int_{x_1}^{x_2} f(x) dx$

(c)  $\int_0^{x_1} f(x) dx - \int_{x_1}^{x_2} f(x) dx$

(d)  $\left| \int_0^{x_2} f(x) dx \right|$



**39 In the opposite figure :**

A part of the curve  $f(x)$  is drawn in interval  $[a, b]$

, if the area M equals 5 square units

, and the area N equal 3 square units


, then  $\int_a^b f(x) dx = \dots\dots\dots$

(a) -5

(b) -2

(c) 2

(d) 8

**40 In the opposite figure :**

If  $A_1 = 5$  square units ,  $A_2 = 2$  square units

,  $A_3 = 8$  square units


, then  $\int_{-2}^5 f(x) dx + \int_{-2}^5 |f(x)| dx = \dots\dots\dots$

(a) 15

(b) 20

(c) 22

(d) 26

**41 In the opposite figure :**

If  $\int_{-3}^4 f(x) dx = 12$

and area of the shaded part = 28 square units


, then  $\int_1^4 f(x) dx = \dots\dots\dots$

(a) -16

(b) -8

(c) 8

(d) -20

**42 If each of  $f$  and  $g$  is a continuous function and**

the graph illustrates the curve of  $f(x)$  and  $g(x)$

and  $A_1 = 3$  square units ,  $A_2 = 2$  square units

,  $A_3 = 4$  square units


which of the following statements is not true ?

(a)  $\int_a^c [f(x) - g(x)] dx = 1$

(b)  $\int_b^d [g(x) - f(x)] dx = -2$

(c)  $\int_a^d [f(x) - g(x)] dx = 5$

(d)  $\int_a^c [f(x) - g(x)] dx = 4$






43 In the opposite figure represents

the curve of the function  $y = f(x)$

and  $\int_{-3}^5 f(x) dx = 8$ ,  $\int_{-3}^5 |f(x)| dx = 12$

then  $\frac{A_1}{A_2} = \dots\dots\dots$

- (a) 5 (b) 4  
(c) 3 (d) 2




44 In the opposite figure :

If the area of the shaded region = 10 square units.

, then  $\int_0^5 [f(x) + f(x)] dx = \dots\dots\dots$

- (a) -4 (b) -18  
(c) -12 (d) -8




45 The opposite figure represents

the graph of the function  $y = f(x)$

If the area of the shaded region = 7 square units.

, then  $\int_5^{11} f(x) dx = \dots\dots\dots$  square units.

- (a) 25 (b) 23  
(c) 21 (d) 19




46 In the opposite figure :

If  $\int_0^4 f(x) dx = 8$ ,  $\int_4^6 f(x) dx = 3$

, then the area of the shaded region =  $\dots\dots\dots$  square units.

- (a) 4  
(b) 5  
(c) 6  
(d) 8





47 (2<sup>nd</sup> session 2021) In the opposite figure :


If  $f(x) = 3(x-1)^2$ ,  $\int_{-1}^4 g(x) dx = 150$ ,  
then the area of shaded part equals .....  
square units

(a) 131

(b) 123

(c) 119

(d) 115



48 In the opposite figure :


If the area of shaded region =  $\frac{8}{3}$  square units  
, then  $a = \dots\dots\dots$

(a)  $\frac{1}{2}$

(b) 1

(c)  $\frac{3}{2}$

(d) 2



49 In the opposite figure :


If the area of the shaded region = 2 square unit  
, then  $k = \dots\dots\dots$

(a) 2

(b) 3

(c) 4

(d) 5



50 In the opposite figure :


ABC is a right-angled triangle at B  
 $\overline{AC}$  touches the curve  $y = 1 - x^2$   
at  $x = -\frac{1}{2}$  then the area of  
the shaded region = ..... area units.

(a)  $\frac{81}{32}$

(b)  $\frac{115}{32}$


(c)  $\frac{115}{96}$

(d)  $\frac{81}{64}$






- 51 (1<sup>st</sup> session 2021) The opposite figure represents the curve of the two functions :  $y = a\sqrt{x} - 2$  ,  $y = a\sqrt{x}$   
If the area of shaded region equals  $\frac{13}{6}$  square units.  
then the value of the constant  $a = \dots\dots\dots$



- (a)  $\frac{1}{2}$   
(b) 1  
(c)  $\frac{1}{3}$   
(d) 2

- 52 In the opposite figure :


If  $y_1$  is a curve of a quadratic function and  $y_2$  is a curve of a circle of center (O) , then the area of the shaded region  
= ..... square units.



- (a) 9  
(b) 12  
(c) 15  
(d) 18

- 53 In the opposite figure :

The straight line  $l$  is tangent to the curve  $y = x^3 + 1$  at  $(1, k)$   
The area of the shaded region  
= ..... square units



- (a) 3  
(b) 2  
(c)  $\frac{4}{3}$   
(d)  $\frac{3}{2}$

**54 In the opposite figure :**


If  $a > 0$  , then the area of

the region bounded by the curve

$y = \frac{1}{x}$  and the  $x$ -axis and between

the two straight lines  $x = a$  and  $x = 2a$  is .....

- (a) increasing when  $a < \frac{1}{2}$  and decreasing when  $a > \frac{1}{2}$
- (b) decreasing when  $a < \frac{1}{2}$  and increasing when  $a > \frac{1}{2}$
- (c) increasing as  $a$  increasing and decreasing as  $a$  decreasing
- (d) constant , it does not depend on the value of  $a$

**55 In the opposite figure :**

The area of the region bounded by


the curve of the function  $f$  and the two

straight lines  $y_1$  and  $y_2$  where :

$f(x) = x^3$  ,  $y_1 = x + 6$  ,  $y_2 = -\frac{1}{2}x$

equals .....


- (a) 11
- (b) 16
- (c) 22
- (d) 27

**56 In the opposite figure :**

If the area of the shaded region = 3 square units

, then  $\int_0^4 f(x) dx = \dots\dots\dots$

- (a) 3
- (b) 4
- (c) 5
- (d) 6





57 The opposite figure represents the curve

of the function  $f : f(x) = x^2$


, then  $\frac{A_1}{A_2} = \dots\dots\dots$

(a)  $\frac{2}{7}$

(b)  $\frac{3}{7}$

(c)  $\frac{4}{7}$

(d)  $\frac{5}{7}$



58 In the opposite figure :

$A_1 = 2$  square units ,

$A_2 = 7$  square units


, then  $\int_{-2}^2 f(x) dx = \dots\dots\dots$

(a) 5

(b) 9

(c) 15

(d) 19



59 In the opposite figure :

If  $A_2$  is the region bounded by the two curves


$y^2 = 4x$  ,  $x^2 = 4y$  , then  $A_1 : A_2 : A_3 = \dots\dots\dots$

(a) 2 : 1 : 2

(b) 1 : 2 : 1

(c) 1 : 1 : 1

(d) 3 : 2 : 3



60 The opposite figure represents the curves

of two functions  $f, g$  in the interval  $[0, 9]$

If  $\int_0^a f(x) dx = \int_0^a g(x) dx$


, then  $a = \dots\dots\dots$

(a) 3

(b) 4

(c) 5

(d) 8






61 In the opposite figure :

$$f(x) = x^2 + 1$$

, then  $k$  which makes

$$A_1 = A_2 \text{ equals } \dots\dots\dots$$

- (a) 3
- (b) 4
- (c) 5
- (d) 8




62 In the opposite figure :

If the curve of the function  $f(x)$  is continuous and convex downward in the interval  $[2, 8]$

, then  $\int_2^8 f(x) dx$  can not be  $\dots\dots\dots$

- (a) 28
- (b) 30
- (c) 32
- (d) 34



63 In the opposite figure :

$f$  is a function defined on


the interval  $[2, 6]$  and  $f(x)$

is a continuous and one-to-one

If the shaded area = 13 square units.

, then  $\int_2^6 f^{-1}(x) dx = \dots\dots\dots$

- (a) 18
- (b) 19
- (c) 20
- (d) 21




64 In the opposite figure :

The area of the shaded region

=  $\dots\dots\dots$  square units.

- (a)  $\frac{22}{3}$
- (b)  $\frac{23}{3}$
- (c) 8
- (d)  $\frac{25}{3}$






65 The opposite figure represents the curve of the function  $f(x)$  in the interval  $[0, 6]$

If  $A_1$  and  $A_2$  are the shaded regions

$A_1 = 20$  square units.

,  $A_2 = 7$  square units,  $f(0) = 2$ , then the greatest value of

the function  $f$  in the interval  $[0, 6]$  equals .....



(a) 13

(b) 15

(c) 22

(d) 29

66 If  $x \in [1, 4]$  and  $f(x) \in [5, 8]$ , then  $\int_1^4 f(x) dx \in$  .....

(a)  $[1, 4]$

(b)  $[5, 8]$

(c)  $[6, 12]$

(d)  $[15, 24]$

67 If the region bounded by the curve  $y = x^2$  and the straight line  $y = 2$  revolved a complete revolution about y-axis, then the volume of the solid generated by revolving equals .....

(a)  $\int_0^2 y dy$

(b)  $\pi \int_0^2 y dy$

(c)  $\pi \int_0^2 x dx$

(d)  $\pi \int_0^2 x^2 dx$

68 If the region bounded by the curve  $y = x^2$  and the straight line  $y = 2$  revolves a complete revolution about the x-axis, then the volume of the generated solid equals .....

(a)  $\pi \int_0^2 x^4 dx$

(b)  $\pi \int_{-\sqrt{2}}^{\sqrt{2}} (4 - x^4) dx$

(c)  $\pi \int_{-2}^2 (4 - x^4) dx$

(d)  $\pi \int_{-\sqrt{2}}^{\sqrt{2}} x^4 dx$

69 The volume of the solid generated by revolving the region bounded by the two curves  $y = x^2$ ,  $y = 1$  a complete revolution about y-axis is .....

(a)  $\pi$

(b)  $\frac{1}{2} \pi$

(c)  $\frac{1}{4} \pi$

(d)  $-\pi$



- 70 The volume of the solid generated by revolving the region bounded by the two curves  $y = x^2$  ,  $y = 1$  a complete revolution about  $X$ -axis equals .....
- (a)  $\frac{8}{5} \pi$                       (b)  $\frac{-8}{5} \pi$                       (c)  $\frac{4}{5} \pi$                       (d)  $\frac{-4}{5} \pi$
- 
- 71 The volume of the solid generated by revolving the region bounded by the curve  $f(x) = x^2$  and  $X$ -axis , and the two straight lines  $x = -2$  ,  $x = 2$  a complete revolution about  $X$ -axis equals .....
- (a)  $\frac{16 \pi}{5}$                       (b)  $\frac{32 \pi}{5}$                       (c)  $\frac{64 \pi}{5}$                       (d)  $4 \pi$
- 
- 72 The volume of the solid generated by revolving the region bounded by the curve  $f(x) = x^2$  ,  $X$ -axis ,  $x = a$  where  $a \in \mathbb{R}^+$  a complete revolution about  $X$ -axis equals .....
- (a)  $\frac{1}{2} \pi a^4$                       (b)  $\pi a^2$                       (c)  $\frac{1}{5} \pi a^5$                       (d)  $\frac{1}{3} \pi a^3$
- 
- 73 The volume of the solid generated by revolving  $\Delta ABC$  such that  $A(-2, 0)$  ,  $B(1, 5)$  ,  $C(4, 0)$  a complete revolution about  $X$ -axis = ..... cubic unit.
- (a)  $25 \pi$                       (b)  $50 \pi$                       (c)  $75 \pi$                       (d)  $90 \pi$
- 
- 74 The volume of the solid generated by revolution the region bounded by the curve  $y^2 = 2ax$  and the straight line  $x = b$  where  $a, b \in \mathbb{R}^*$  half revolution about  $X$ -axis equals ..... cubic unit.
- (a)  $\pi a b^2$                       (b)  $\pi b a^2$                       (c)  $\pi a b$                       (d)  $\pi a^2 b^2$
- 
- 75  $\int_0^r \pi x dx = \dots\dots\dots$
- (a) Perimeter of a circle whose radius length  $r$   
 (b) half the volume of a sphere whose radius length  
 (c) half the perimeter of a circle whose radius length  $r$   
 (d) half the area of a circle whose radius length  $r$



76  $\int_0^h \pi r^2 dX = \dots\dots\dots$

- (a) volume of a circular cylinder whose height (h) and its base radius is (r)
- (b) the area of a sphere whose radius length (h)
- (c) the lateral area of a right circular cylinder whose height (h) and its base radius is (r)
- (d)  $\frac{1}{3} \pi h^3 + c$

77 The volume of the solid generated by revolving the region bounded by the straight line  $y = X + 1$  and the two straight lines  $X = 0$  ,  $y = 2$  a complete revolution about  $X$ -axis equals .....

- (a)  $\frac{5}{3} \pi$
- (b) 4
- (c)  $\frac{7}{3} \pi$
- (d)  $3 \pi$

78  $\pi \int_{-2}^2 (4 - X^2) dX$  is the volume of .....

- (a) a sphere whose radius length is 4 units.
- (b) a right circular cone whose height is 4 units.
- (c) a sphere whose radius length is 2 units.
- (d) a right circular cylinder whose height is 4 units.

79 The volume of a solid generated by revolving the region bounded by the curve  $y = X(X - 2)$  and  $X$ -axis a complete revolution about  $X$ -axis equals .....

- (a)  $\frac{-4}{3} \pi$
- (b)  $\frac{4}{3} \pi$
- (c)  $\frac{16}{15} \pi$
- (d)  $\frac{16}{15}$

80 The volume of the solid generated by revolving the region enclosed by the curve  $y = 2X^2$  and the line  $y = 8X$  a complete revolution about the  $X$ -axis is equal to .....

- (a)  $\pi \int_0^8 (8X - 2X^2)^2 dX$
- (b)  $\pi \int_0^4 (8X - 2X^2)^2 dX$
- (c)  $\pi \int_0^4 (64X^2 - 4X^4) dX$
- (d)  $\pi \int_0^4 (4X^4 - 64X^2)^2 dX$



- 81 When the region bounded by the curve  $x = \frac{1}{\sqrt{y}}$ ,  $1 \leq y \leq 4$  and y-axis revolves a complete revolution about y-axis, then the volume of the solid generated measured by cubic units equals .....
- (a)  $\frac{2}{3} \pi$  (b)  $3\sqrt{2} \pi$  (c)  $2 \pi \ln 2$  (d)  $\frac{2}{3} \pi \log 3$
- 
- 82 The volume of the solid generated by revolving the region bounded by the curve :  $y = \sec x$  and the two straight lines :  $x = 0$ ,  $x = \frac{\pi}{3}$  a complete revolution about x-axis = ..... cubic unit
- (a)  $\sqrt{3}$  (b)  $\frac{1}{\sqrt{3}}$  (c)  $\frac{\pi}{\sqrt{3}}$  (d)  $\sqrt{3} \pi$
- 
- 83 The solid generated by revolving the region bounded by the two curves  $y = \tan x$ ,  $y = \sec x$  and the two straight lines  $x = \frac{\pi}{6}$ ,  $x = \frac{\pi}{3}$  a complete revolution about x-axis measured by cubic units equals .....
- (a)  $\frac{\pi^2}{6}$  (b)  $\frac{\pi^2}{3}$  (c)  $\frac{2\pi^2}{5}$  (d)  $2\pi^2$
- 
- 84 The volume of the solid generated by revolving the region bounded by the curve  $f(x) = \sqrt{25 - x^2}$  and  $g(x) = 3$  a complete revolution about x-axis = ..... cubic unit.
- (a)  $\frac{232}{3} \pi$  (b)  $\frac{244}{3} \pi$  (c)  $\frac{256}{3} \pi$  (d)  $\frac{268}{3} \pi$
- 
- 85 The volume of the solid generated by revolving the region bounded by the two curves  $y = \sin x$  and  $y = \cos x$  and the y-axis where  $x \in [0, \frac{\pi}{4}]$  a complete revolution about x-axis equals ..... cubic unit.
- (a)  $\frac{1}{4} \pi$  (b)  $\frac{1}{2} \pi$  (c)  $\frac{3}{4} \pi$  (d)  $\pi$
- 
- 86 ABCD is a trapezium in which A (0, 0), (2, 0), C (2, 5), D (0, 3), then the volume of the solid generated by revolving the trapezium ABCD a complete revolution about x-axis = ..... cubic unit
- (a)  $\frac{98}{3} \pi$  (b)  $\frac{160}{3} \pi$  (c)  $\frac{223}{3} \pi$  (d)  $\frac{226}{3} \pi$




- Scanned with CamScanner

**92 In the opposite figure :**


The volume of the solid generated by revolving the shaded region a complete revolution about the y-axis equals ..... cube units.

- (a)  $4\pi$  (b)  $6\pi$   
(c)  $8\pi$  (d)  $10\pi$

**93 In the opposite figure :**


The volume of the solid generated by revolving the shaded area a complete revolution about the X-axis = ..... cube units.

- (a)  $\frac{\pi}{3}$  (b)  $\frac{\pi}{2}$   
(c)  $\frac{2\pi}{3}$  (d)  $\pi$

**94 In the opposite figure :**


The volume of the solid generated by revolving the shaded area a complete revolution about the y-axis = ..... cube units.

- (a)  $\frac{4}{3}\pi$  (b)  $\frac{8}{3}\pi$   
(c)  $6\pi$  (d)  $\frac{16}{3}\pi$

**95 In the opposite figure :**

The straight line L is a tangent to the curve  $y = \sqrt{2x - 2}$  at  $(3, 2)$ , then the volume of the solid generated by revolving the shaded region a complete revolution about the X-axis equals ..... cube units.

- (a)  $\frac{4}{3}\pi$  (b)  $\frac{3}{4}\pi$   
(c)  $\frac{1}{3}\pi$  (d)  $\frac{2}{3}\pi$





96 In the opposite figure :


The axis of a right cone lies along the  $X$ -axis and its vertex at the origin , then its volume = .....

(a)  $\pi_0 \int_0^h \left(\frac{r}{h} x\right) dx$

(b)  $\pi_0 \int_0^r \left(\frac{h}{r} y\right) dy$

(c)  $\pi_0 \int_0^h \left(\frac{r^2}{h^2} x^2\right) dx$


(d)  $\pi_0 \int_0^r \left(\frac{h^2}{r^2} y^2\right) dy$



97 The opposite figure represents the curve :

$y = \frac{\ln x}{\sqrt{x}}$  and the line  $x = e$  , then the volume of the generated solid

by revolving the shaded region a complete revolution about the  $X$ -axis = ..... cube units.



(a)  $\frac{1}{3} \pi$


(b)  $\pi$

(c)  $3 \pi$

(d)  $9 \pi$

98 The opposite figure represents the curve  $y = x^3$  and the line  $x = 1$

, then the volume of the solid generated by revolving the shaded region a complete revolution about the  $y$ -axis = ..... cube units.



(a)  $\frac{1}{7} \pi$


(b)  $\frac{2}{5} \pi$

(c)  $\frac{3}{5} \pi$

(d)  $\frac{3}{4} \pi$

99 In the opposite figure :

The volume of the solid generated by revolving the shaded region a complete revolution about  $X$ -axis = ..... cubic units.



(a)  $\frac{14}{3} \pi$

(b)  $\frac{15}{2} \pi$


(c)  $\frac{15}{2}$

(d)  $\frac{14}{3}$

**100 In the opposite figure :**


If the volume of the solid generated by revolving the shaded area a complete revolution about  $X$ -axis on the interval  $[a, b]$  equals  $8\pi$ , then  $b^2 - a^2 = \dots\dots\dots$

- (a) 8 (b) 12  
(c) 16 (d) 20

**101 In the opposite figure :**


If the volume of the solid generated by revolving the shaded region a complete revolution about  $X$ -axis, equals  $\frac{\pi}{2}(e^{10} - e^{-2})$  cubic unit, then :  $k = \dots\dots\dots$

- (a) 5 (b) 10  
(c) 20 (d) -5

**102 (2<sup>nd</sup> session 2021) In the opposite figure :**


If the volume of the solid generated by revolving the shaded region complete revolution about  $y$ -axis equals  $8\pi$  cubic units, then  $a = \dots\dots\dots$

- (a) 8 (b) 16  
(c) 24 (d) 4

**103 (1<sup>st</sup> session 2021) In the opposite figure :**

If the volume of the solid generated by revolving the shaded region included between the curve  $xy = 3$ , the two straight lines  $y = 1$ ,  $y = k$  and  $y$ -axis complete revolution about  $y$ -axis equals  $6\pi$  cubic units, then  $k = \dots\dots\dots$

- (a) 4 (b) 2  
(c) 3 (d) 1.5






**104 In the opposite figure :****First :** The area of the shaded region

= ..... square units

- (a)  $\frac{4}{3}$  (b)  $\frac{8}{3}$   
 (c)  $\frac{16}{3}$  (d) 4

**Second :** The value of the solid generated by revolving the shaded region a complete revolution about y-axis equals ..... cubic unit.

- (a)  $32\pi$  (b)  $\frac{64}{7}\pi$  (c)  $\frac{128}{7}\pi$  (d)  $\frac{64}{5}\pi$


**Third :** The volume of the solid generated by revolving the shaded region a complete revolution about X-axis equals ..... cubic units.

- (a)  $9.6\pi$  (b)  $10.8\pi$  (c)  $11.2\pi$  (d)  $12.8\pi$

**105 In the opposite figure :**

If  $\int_0^{12} \pi (f(x))^2 dx = 135\pi$ , then the volume of the solid generated by revolving the shaded region about the X-axis = ..... cubic units.


- (a)  $35\pi$  (b)  $25\pi$   
 (c)  $20\pi$  (d)  $15\pi$

**106 In the opposite figure :**

If the volume of the solid generated by revolving the shaded area about the X-axis = a cube units and about the y-axis = b cube units.


, then which of the following statements is true ?

- (a)  $a = b$  and the two solids are congruent.  
 (b)  $a = b$  and the two solids are not congruent.  
 (c)  $a > b$   
 (d)  $a < b$







- 107 Which of the following figures, the volume of the solid generated by revolving about the  $x$ -axis equals the volume of the solid generated by revolving about the  $y$ -axis ?




(a)



(b)



(c)




(d)

- 108 In the opposite figure : The curve  $f(x) = 6x^2$ , then the value of  $k$  which makes the shaded region as small as possible equals .....

(a)  $1\frac{1}{4}$ (b)  $1\frac{3}{4}$ 

(c) 2


(d)  $2\frac{1}{4}$ 

- 109 In the opposite figure :  
The area of the shaded region when the area of rectangle ABCD is as big as possible equals ..... square unit.


(a)  $\frac{8}{3}$ 

(b) 4

(c) 5

(d)  $\frac{32}{3}$ 

- 110 In the opposite figure :  
If  $B = (12, 0)$ , then the greatest volume of the solid generated by revolving the shaded region a complete revolution about  $x$ -axis = ..... cubic unit.


(a)  $48\pi$ (b)  $36\pi$ (c)  $24\pi$ (d)  $18\pi$ 



# Practice Exams

on

## Differential & Integral calculus







Answer the following questions :

1  $\lim_{x \rightarrow 0} x\sqrt{1+x} = \dots\dots\dots$

(a) 0

(b) 1

(c)  $\frac{1}{e}$

(d) e

2 The length of the radius of a circle increase at a rate  $\frac{4}{\pi}$  cm./sec. , then the rate of increase of its circumference at this moment is ..... cm./sec.

(a)  $\frac{4}{\pi}$

(b)  $\frac{\pi}{4}$

(c)  $\frac{1}{8}$

(d) 8


3 If the volume of the solid generated by revolving the shaded region included between the curve  $xy = 3$  , the two straight lines  $y = 1$  ,  $y = k$  and y-axis complete revolution about y-axis equals  $6\pi$  cubic units , then  $k = \dots\dots\dots$

(a) 4

(b) 2

(c) 3

(d) 1.5



4 If  $y = \cot 5x$  , then  $\frac{dy}{dx} + 5y^2 = \dots\dots\dots$

(a) 5

(b)  $-5y$

(c)  $5y$

(d)  $-5$

5 If  $f(x) = x^3 - 3x - 4$  , then the function  $f$  is decreasing when .....

(a)  $|x| < 1$

(b)  $|x| > 1$

(c)  $x > 1$

(d)  $x < -1$

6 If  $y^2 = 1 - \frac{1}{x^2}$  , then  $y \frac{d^2y}{dx^2} + \left(\frac{dy}{dx}\right)^2 = \dots\dots\dots$

(a)  $3x^4$

(b)  $-3x^{-4}$

(c)  $3x^{-4}$

(d)  $-3x^{-2}$

7  $\int \frac{x}{x^2 - 2x + 1} dx = \dots\dots\dots + c$  (where  $c$  is constant)

(a)  $\ln|x-1| - \frac{1}{x-1}$

(b)  $\ln|x-1| + \frac{1}{x-1}$


(c)  $\ln|x-1| + (x-1)^2$

(d)  $\ln|x^2 - 2x + 1| + \frac{1}{x-1}$



- (a) 100                      (b) 50                      (c) 200                      (d) 400

(a) 131                      (b) 123  
(c) 119                      (d) 115



- (a) parallel to y-axis.      (b) parallel to x-axis.
- (c) parallel to the straight line  $y = x$       (d) parallel to the straight line  $y = -x$

- (a) 1                      (b) 10                      (c) 55                      (d)  $(55)^2$

- (a)  $\frac{\pi}{2}$       (b)  $\frac{\pi}{3}$       (c)  $\frac{\pi}{4}$       (d)  $\frac{\pi}{6}$

- (a)  $\frac{x}{a} - \frac{y}{b} = 1$ 
 (b)  $a x + b y = 1$ 
 (c)  $a x - b y = 1$ 
 (d)  $\frac{x}{a} + \frac{y}{b} = 1$

- (a)  $-2e^{\cos 2x}$       (b)  $2e^{\cos 2x}$       (c)  $4e^{\cos 2x}$       (d)  $-4e^{\cos 2x}$



15 The derivative of  $\sin^2 X$  with respect to  $\cos^2 X$  is .....

- (a)  $-\tan^2 X$  (b)  $\tan^2 X$   
(c)  $-(\sin^2 X + \cos^2 X)$  (d)  $\cos^2 X - \sin^2 X$

16  $\int (1 + \cos X)^2 dX = \dots + c$

- (a)  $(1 + \sin X)^2$  (b)  $\frac{1}{3} (1 + \cos X)^3$   
(c)  $\frac{3}{2} X + 2 \sin X + \frac{1}{4} \sin 2X$  (d)  $\frac{3}{2} + 2 \cos X + \frac{1}{2} \cos 2X$

17 If  $\int \frac{e^{ax}}{e^{bx}} dX = k \frac{e^{ax}}{e^{bx}} + c$ , then  $k = \dots$

- (a)  $a + b$  (b)  $a - b$  (c)  $\frac{1}{a - b}$  (d)  $\frac{1}{a + b}$

18 If the function  $f : f(X) = 3aX^3 - bX - 5$  has local maximum value at  $X = 1$ , then  $\frac{b}{a} = \dots$

- (a)  $-9$  (b)  $9$  (c)  $20$  (d)  $-20$

19 If the perimeter of a triangle is 8 cm., the length of one of its sides is 3 cm., then the maximum area for this triangle  $= \dots \text{ cm}^2$

- (a) 3 (b) 2 (c) 4 (d) 5

20 If  $a$  and  $b$  are the absolute extrema of the function  $f : f(X) = \begin{cases} X^2 + X - 2 & , X \leq 2 \\ 5X - 6 & , X > 2 \end{cases}$  in the interval  $[-1, 3]$ , then  $a + b = \dots$

- (a) 6 (b)  $\frac{28}{9}$  (c)  $4\frac{1}{2}$  (d) 6.75

21 If  $\int (2X - 1) e^{2X+3} dX = yz - \int z dy$ , then  $\int z dy = \dots$

- (a)  $e^{2X+3} + c$  (b)  $\frac{1}{2} e^{2X+3} + c$  (c)  $-e^{2X+3} + c$  (d)  $-\frac{1}{2} e^{2X+3} + c$

22 If  $f$  is a function of fourth degree, then the greatest possible number of the inflection points equals .....

- (a) 2 (b) 1 (c) 3 (d) 4



23 If  $f(x) = a \ln |x| + b x^2 + x$  has critical point at  $x = -1$ ,  $x = 2$ , then  $a, b = \dots\dots\dots$

(a) 1

(b) -1


(c) 2

(d)  $\frac{1}{2}$ 


24 If  $y = f(x)$ ,  $\frac{d^2 x}{d y^2} = e^{y+1}$  and the tangent to the curve of the function  $f$  at the point  $(x, -1)$  which lies on it is parallel to the straight line  $y = x - 3$ , then the slope of the tangent to the curve of the function  $f$  at any point  $(x, y)$  lies on it equals  $\dots\dots\dots$

(a)  $e^{y+1}$ (b)  $e^{2y+2}$ (c)  $e^{-y-1}$ (d)  $e^{-2y-2}$ 


25 If  $f$  is a continuous function has the following properties :  $f(3) = 1$ ,  $f'(x) < 0$  when  $x < 3$ ,  $f'(x) > 0$  when  $x > 3$  and  $f''(x) < 0$  for every  $x \neq 3$ , then the curve which can represent this function is  $\dots\dots\dots$




(a)



(b)



(c)



(d)



Answer the following questions :

1  $\lim_{x \rightarrow 0} \frac{e^{5x} - 1}{x} = \dots\dots\dots$

(a) 5

(b)  $e^5$

(c)  $\ln 5$

(d)  $5e$

2 If  $f(x) = \sin 2x$ , then  $f\left(\frac{\pi}{4}\right) = \dots\dots\dots$

(a) zero

(b) -2

(c) -4

(d) -6

3  $\int \sec x \, dx = \dots\dots\dots + c$

(a)  $\sec x \tan x$

(b)  $\ln |\sec x + \tan x|$

(c)  $\cos^{-1} x$

(d)  $\frac{1}{2} \sec^2 x$

4 The equation of the curve passes through the point (0, 1) and the slope of its tangent at any point on it (x, y) equals  $x\sqrt{x^2 + 1}$  is .....

(a)  $y = \frac{1}{3} (x^2 + 1)^{\frac{2}{3}} + \frac{2}{3}$

(b)  $y = \frac{1}{3} (x^2 + 1)^{\frac{3}{2}} + \frac{2}{3}$

(c)  $y = \frac{1}{3} (x^2 + 1)^{\frac{3}{2}} - \frac{2}{3}$

(d)  $y = \frac{4}{3} (x + 1)^{\frac{2}{3}} - \frac{2}{3}$


5 The opposite figure represents the curve of the function  $f : f(x) = x^3 + a$ , if the area of the shaded region equals 68 square unit, then  $a = \dots\dots\dots$

(a) 1

(b) 2

(c) 3

(d) 4



6 In the opposite figure :


The length of the square increases by rate  $\frac{1}{4}$  cm./sec., then the rate of change of the shaded areas when the length of the square side equals 16 cm. equals .....  $\text{cm}^2/\text{sec}.$

(a) 2

(b) 4

(c) 8

(d) 16





7 If  $\sin X \cos y - \sin y \cos X = 1$ , then  $\frac{dy}{dX} = \dots\dots\dots$  where  $X, y \in ]0, 2\pi[$

- (a)  $\cos(X - y)$  (b)  $\sin(X - y)$  (c) 1 (d) -1


8 If  $f: \mathbb{R} \rightarrow \mathbb{R}$ ,  $f(X) = X^3 + aX^2 + 12X + 1$  and the function has no critical points, then  $a \in \dots\dots\dots$

- (a)  $]-6\sqrt{2}, 6\sqrt{2}[$  (b)  $]-3, 3[$  (c)  $]-12, 12[$  (d)  $]-6, 6[$

9 In the opposite figure :

If the volume of the solid generated by revolving the shaded region complete revolution about y-axis equals  $8\pi$  cubic units, then  $a = \dots\dots\dots$

- (a) 8 (b) 16  
(c) 24 (d) 4



10 The absolute minimum value of the function  $f(X) = \sqrt{X^2 + 9}$  where  $X \in [-3, 3]$  equals  $\dots\dots\dots$

- (a)  $3\sqrt{2}$  (b)  $-3\sqrt{2}$  (c) 3 (d) -3

11 The equation of the tangent to the curve :  $Xy = X^y$  at the point (1, 1) which lies on it is  $\dots\dots\dots$

- (a)  $X - 1 = 0$  (b)  $y + X = 2$  (c)  $y - 1 = 0$  (d)  $X - y = 0$

12 If  $f\left(\frac{1}{2}X\right) = |X|^3$ , then  $f'(-1) = \dots\dots\dots$

- (a) -48 (b) 48 (c) 14 (d) 1

13 If  $y = X^{n+1} + nX^{n-1} + 1$ , then  $\frac{d^n y}{dX^n} = \dots\dots\dots$

- (a)  $\underline{n+1}$  (b)  $X \underline{n+1}$  (c)  $X \underline{n}$  (d)  $X^{-1} \underline{n}$

14 If  $z = X + 2$ , then  $\int \frac{X-2}{X^2+4X+4} dX = \dots\dots\dots + c$  (where c is constant)

- (a)  $\ln|z| - \frac{4}{z}$  (b)  $\ln|z| + \frac{4}{z}$  (c)  $\frac{-1}{2z^2} - \frac{4}{3z^3}$  (d)  $\frac{-1}{2z^2} + \frac{2}{3z^3}$



- 15 If the function  $f : f(x) = \sin x + \frac{1}{2} \sin 2x, x \in ]0, \frac{\pi}{2}[$ , then the function  $f$  .....
- (a) has a local minimum value at  $x = \frac{\pi}{4}$   
 (b) has a local maximum value at  $x = \frac{\pi}{6}$   
 (c) has not a local maximum value at this interval  
 (d) has a local maximum value at  $x = \frac{\pi}{3}$

16 If  $y = \frac{\log_x 2e}{\log_x 3e}$ , then  $\frac{dy}{dx} = \dots\dots\dots$


- (a) 0 (b) 1 (c) e (d)  $\frac{1}{e}$

- 17 If the function  $f : f(x) = x^2 + \frac{2a^3}{x}$  where  $a \in \mathbb{R}^-$ , then the function is decreasing in the interval .....

- (a)  $]-\infty, 0[$ ,  $]0, \infty[$  (b)  $]a, \infty[$   
 (c)  $]-\infty, a[$  (d)  $]-\infty, 0[$ ,  $]a, 0[$

- 18 The opposite figure represents the curve of the continuous function  $g : g(x)$  on  $\mathbb{R}$  that has a horizontal tangent at  $x = 1$ , if  $f$  is differentiable function on  $\mathbb{R}$  where  $f'(x) = |g(x)|$ , then the true statement from the following is .....

- (a) The function  $f$  has inflection point at  $x = -1$   
 (b) The function  $f$  has minimum value at  $x = -1$   
 (c) The function  $f$  has inflection point at  $x = 1$   
 (d) The curve of the function  $f$  convex upward in  $]-1, 1[$



19  $-\frac{\pi}{4} \int_4^{\frac{\pi}{4}} \frac{\tan x}{x^2 + \cos x} dx = \dots\dots\dots$

- (a)  $-\sqrt{3}$  (b)  $-\frac{\sqrt{3}}{3}$  (c) zero (d)  $\frac{\sqrt{3}}{3}$

- 20 The area of the region bounded by the curve  $y = 6 - x^2$  and the straight line passing through the two points  $(3, -3)$ ,  $(-2, 2)$  equals ..... square units.

- (a)  $\frac{56}{3}$  (b)  $\frac{55}{3}$  (c)  $\frac{95}{6}$  (d)  $\frac{125}{6}$



21 In the opposite figure :


$C \in$  the line  $x + 3y = 6$ , then the greatest area of the isosceles triangle  $OBC = \dots\dots\dots$  square unit.

(a) 2

(b) 3

(c) 6

(d) 9



22 By discussion the convexity intervals upward and downward and point of inflection (if exist) to the curve of the function  $f$  where  $f(x) = (x-1)^4 + 3$  the curve is .....

(a) convex downward on the interval  $]-\infty, 1[$  and convex upward on the interval  $]1, \infty[$  and it has no inflection point.

(b) convex downward on the intervals  $]-\infty, 1[$  and  $]1, \infty[$  and it has an inflection point at  $x = 1$

(c) convex downward on the intervals  $]-\infty, 1[$  and  $]1, \infty[$  and it has no inflection point.

(d) convex upward on the interval  $]-\infty, 1[$  and convex downward on the interval  $]1, \infty[$  and it has an inflection point at  $x = 1$

23 In the curve equation  $y = f(x)$  if  $\frac{d^2 y}{dx^2} = ax + b$  where  $a, b$  are constant and the curve has an inflection point  $(0, 2)$  and local minimum value at the point  $(1, 0)$ , then  $2a + b = \dots\dots\dots$

(a) 6

(b) 12

(c) -12

(d) -60

24 If  $a > 1 > b > 0$ ,  $\int_1^a \frac{1}{x} dx = \int_b^1 \frac{1}{x} dx$ , then .....

(a)  $a = b$

(b)  $a = -b$

(c)  $a = \frac{1}{b}$

(d)  $a = \frac{1}{\ln b}$


25 The opposite figure represents the function  $f$  and the straight line  $L$  touches the curve of  $f$  at the point  $A(3, 5)$  and  $g(x) = x \cdot f(x)$ , then  $g'(3) = \dots\dots\dots$

(a) 3

(b) 1

(c) 5

(d) 8





Answer the following questions :

1 If  $\lim_{x \rightarrow 0} \frac{\ln(1+x)^{4a}}{x} = a^2 + 4$ , then the value of constant  $a = \dots\dots\dots$

(a) -2

(b) -4

(c) 2

(d) 4

2  $-\pi \int \frac{4x + \sin x}{x^2 + \cos x} dx = \dots\dots\dots$

(a)  $-\pi$

(b) 0

(c)  $\pi$

(d)  $2\pi$

3 If  $f$  is a function,  $f : f(x) = x^2 + ax + b$  has local minimum value = 3 at  $x = 1$ , then  $ab = \dots\dots\dots$

(a) -48

(b) -8

(c) 2

(d) -12

4 The equation of the normal to the curve  $y = \ln(\tan x)$  at the point which lies on the curve and its  $x$ -coordinate equals  $\frac{\pi}{4}$  is  $\dots\dots\dots$

(a)  $4x - 8y = \pi$

(b)  $8y + 4x = \pi$

(c)  $4x + 2y = \pi$

(d)  $4x - 2y = \pi$

5 If the area of the region bounded by the two curves  $y = x^2$ ,  $y = kx$  equals  $\frac{9}{2}$  square units, where  $k > 0$ , then  $k = \dots\dots\dots$

(a) 9

(b) 3

(c) 6

(d) 12

6 The slope of the tangent to the curve of the function  $y = f(x)$  at a particular point is  $\frac{1}{2}$  and the  $x$ -coordinate of this point decreases at a rate 3 units/sec., then the rate of change of its  $y$ -coordinate equals  $\dots\dots\dots$  unit/sec.

(a)  $-\frac{1}{6}$


(b)  $-\frac{3}{2}$

(c)  $\frac{1}{6}$

(d)  $\frac{3}{2}$



7 If the opposite figure represents the curve of the first derivative of a continuous function  $f$  whose domain is  $\mathbb{R}$ , then the wrong statement from the following is .....



- (a) The function has inflection point at  $x = 4$
- (b) The function has local maximum value at  $x = 2$
- (c) The curve of the function convex upward in  $]-\infty, 4[$  and convex downward in  $]4, \infty[$
- (d)  $f(-3) < f(-2)$

8 If  $x = \tan(1 + y)$ , then  $\frac{dy}{dx} = \dots$  at  $x = 2$

- (a) 0.5
- (b) 0.2
- (c) 1
- (d) 5


9 The derivative of  $\frac{ax+b}{cx+d}$  with respect to  $\frac{ax+b}{cx+d}$  equals .....

- (a) 0
- (b) 1
- (c)  $\frac{a}{(cx+d)^2}$
- (d)  $\frac{b}{(ax+b)^2}$

10 If  $y = e^{x^3-4}$  and  $2\dot{y} = m x^2 y$  where  $x \neq 0$ , then  $m = \dots$

- (a) 6
- (b) 4
- (c) 3
- (d) 2

11 The opposite figure represents the curve of the function  $f: f(x) = \frac{4}{x^2}$  in the first quadrant, then the coordinate of the point A which makes the perimeter of the rectangle ABOC has minimum value is .....



- (a)  $(\frac{1}{2}, 16)$
- (b)  $(2, 1)$
- (c)  $(1, 4)$
- (d)  $(\frac{1}{4}, 64)$

12 If  $f(x) = (x+3)e^x$ , then the curve of the function  $f$  convex downward in the interval .....

- (a)  $]-\infty, -5[$
- (b)  $]-6, \infty[$
- (c)  $]-5, \infty[$
- (d)  $]-\infty, -3[$



13 If  $\int \frac{4x^3 - ax}{x^4 - 2x^2 + 3} dx = \ln |x^4 - 2x^2 + 3| + c$ , then  $a = \dots\dots\dots$

(a) 1

(b) 2

(c) 3


(d) 4

14 If the slope of tangent to the curve  $y = f(x)$  at any point  $(x, y)$  on it equals  $(e^x - e^{-x})$  then the equation of this curve is  $\dots\dots\dots$  given that the curve passes through the point  $(0, 4)$ .

(a)  $y = e^x + e^{-x} + 4$ (b)  $y = e^x + e^{-x} + 2$ (c)  $y = e^x + e^{-x} - 4$ (d)  $y = e^x + e^{-x} - 2$ 

15 In the opposite figure :

Two squares having the same center and their side lengths are 1 cm. and 4 cm. , if side length of the first square increases by rate 1 cm./sec. and side length of the second square decreases by rate  $\frac{1}{2}$  cm./sec. , then the area of the shaded region after  $\frac{1}{2}$  second  $\dots\dots\dots$



(a) stop increasing instantly.

(b) stop decreasing instantly.

(c) increases at rate  $\frac{1}{4}$  length unit/sec.(d) decreases at rate  $\frac{1}{4}$  length unit/sec.

16 When Sameh calculated the value of  $\int (\tan x + \tan^3 x) dx$  the result was  $\frac{1}{2} \sec^2 x + a$  while the result when Youssef calculated the same integration was  $\frac{1}{2} \tan^2 x + b$  If both of them got the full mark , then  $a - b = \dots\dots\dots$

(a) -1

(b)  $-\frac{1}{2}$ (c)  $\frac{1}{2}$ 

(d) 1

17 If  $\int \frac{2}{y} dy = \int \frac{1}{x} dx$ , then  $\ln y^2 = \dots\dots\dots + c$  where  $c$  is constant

(a)  $\ln |x|$ (b)  $|x|$ (c)  $\ln |x + y|$ (d)  $\ln |x - y|$ 

18  $\frac{\pi}{4} \int_2^{\frac{\pi}{2}} \frac{e^{\cot x}}{\sin^2 x} dx = \dots\dots\dots$

(a) 1

(b)  $e - 1$ (c)  $e + 1$ (d)  $e$



19 If  $y = a(1 - \cos \theta)$ ,  $x = a(\theta + \sin \theta)$ , then  $\frac{dy}{dx} = \dots\dots\dots$

- (a)  $\tan \theta$  (b)  $\cot \theta$  (c)  $\tan \frac{\theta}{2}$  (d)  $\cot \frac{\theta}{2}$

20  $\int \frac{e^{2x}}{e^{-x}} dx = \dots\dots\dots + c$

- (a)  $\frac{e^{2x}}{e^{-x}}$  (b)  $\frac{1}{2} \frac{e^{2x}}{e^{-x}}$   
(c)  $\frac{1}{3} \frac{e^{2x}}{e^{-x}}$  (d)  $\frac{1}{e^{3x}}$

21 Let  $f(x) = (\cos x)^{\cos x}$ , then  $f'(0) = \dots\dots\dots$

- (a)  $-3$  (b)  $-2$  (c)  $-1$  (d) zero

22 If  $\int (2x+3) \ln x dx = yz - \int z dy$ , then  $yz$  equals  $\dots\dots\dots$

- (a)  $2x \ln x$  (b)  $(2x+3) \ln x$   
(c)  $\frac{1}{2} (2x+3) \ln x$  (d)  $x(x+3) \ln x$

23 If  $f(x)$  is continuous decreasing on the interval  $[0, 10]$  and has a critical point at  $(4, 2)$  which of the following must be false?

- (a)  $f(4)$  has neither local minimum nor local maximum value.  
(b)  $f'(4)$  does not exist.  
(c)  $f'(4) = 0$   
(d)  $f'(4) < 0$

24 If  $f(x) = \frac{2}{x+1}$ ,  $g(x) = 3x$ , then  $\frac{d}{dx} [(f \circ g)(x)] = \dots\dots\dots$  at  $x = -2$

- (a)  $\frac{-3}{25}$  (b)  $6$  (c)  $\frac{1}{25}$  (d)  $\frac{-6}{25}$

25 If the volume of the solid generated by revolving the region bounded by the curve  $y = \frac{1}{x}$  and the two straight lines  $x = k$ ,  $x = 2k$  where  $(k > 0)$  is completely revolved about the  $x$ -axis equals  $\frac{\pi}{3}$  cubic unit, then  $k = \dots\dots\dots$

- (a)  $\frac{1}{2}$  (b)  $\frac{1}{3}$  (c)  $\frac{3}{2}$  (d)  $\frac{2}{3}$



Answer the following questions :

1 If  $\lim_{x \rightarrow 0} \frac{e^{x+a} - e^a}{x} = \frac{1}{e}$ , then the value of  $a = \dots\dots\dots$

- (a) zero (b) 1 (c)  $e$  (d)  $-1$

2 If  $f$  is a polynomial function, where  $f^{(4)}(x)$  is 7<sup>th</sup> degree then  $f^{(7)}(x)$  is  $\dots\dots\dots$  degree.


- (a) 10<sup>th</sup> (b) 7<sup>th</sup> (c) 4<sup>th</sup> (d) 3<sup>rd</sup>

3  $\int e^2 dx = \dots\dots\dots + c$

- (a)  $e^2 x$  (b)  $\frac{1}{3} e^3$  (c)  $e^2$  (d)  $\frac{1}{3} e^3 x$

4 The opposite figure represents the curves of the two functions  $f(x) = ax - 2$ ,  $g(x) = a\sqrt{x}$  if the area of shaded region equals  $\frac{13}{6}$  square units, then the value of the constant  $a = \dots\dots\dots$

- (a)  $\frac{1}{2}$  (b) 1  
(c)  $\frac{1}{3}$  (d) 2




5 If  $y = \frac{1}{(x+1)^2}$ , then the rate of change of  $\frac{1}{y}$  with respect to  $x^2$  at  $x = 1$  is  $\dots\dots\dots$

- (a)  $\frac{1}{2}$  (b) 4 (c) 2 (d)  $\frac{1}{4}$

6 The opposite figure represents the curve of the function  $f$  then the sign of the expression  $[f'(0) - f'(2)]$  is  $\dots\dots\dots$

- (a) negative. (b) positive.  
(c) zero. (d) not sufficient givens.



7 The absolute maximum value of the function  $f : f(x) = x \ln x$  where  $x \in [e^{-2}, e]$  equals  $\dots\dots\dots$

- (a)  $-2e^{-2}$  (b)  $e$  (c)  $2e$  (d)  $-e^{-1}$



8  $\int \sqrt{x} (1 + \sqrt{x}) dx = \dots + c$

(a)  $x^{\frac{1}{2}} + x$

(b)  $\frac{2}{3} x^{\frac{3}{2}} + 2x^2$

(c)  $\frac{3}{2} x^{\frac{2}{3}} + \frac{1}{2} x^2$

(d)  $\frac{2}{3} x^{\frac{3}{2}} + \frac{1}{2} x^2$

9  $\int \frac{2}{x \ln x^2} dx = \dots + c$  (where  $c$  is constant)

(a)  $2 \ln |\ln x|$

(b)  $\frac{1}{2} \ln |\ln |x||$

(c)  $\ln |\ln |x||$

(d)  $\ln |x|$

10 If  $e^{xy} - x^2 + y^3 = 0$ , then  $\frac{dy}{dx}$  (at  $x = 0$ ) equals .....

(a)  $-1$

(b)  $-\frac{1}{3}$

(c)  $1$

(d)  $\frac{1}{3}$

11 The volume of the solid generated by revolving the region bounded by the curve  $y = x^2$  and the straight line passes through the two points  $(0, 6)$ ,  $(1, 7)$  a complete revolution about  $x$ -axis equals ..... cubic unit.

(a)  $\frac{500}{3}$

(b)  $\frac{665}{3} \pi$

(c)  $55 \pi$

(d)  $\frac{500}{3} \pi$

12  $\int \tan^2 x dx = \dots$

(a)  $\tan x - x + c$

(b)  $\tan x + x + c$

(c)  $\sec^4 x + c$

(d)  $\frac{1}{3} \tan^3 x + c$

13 If  $f$  an even continuous function in  $\mathbb{R}$  and  $\int_0^3 f(x) dx = 5$ , then  $\int_{-3}^3 [f(x) - f(x)] dx = \dots$

(a)  $15$

(b)  $-10$

(c)  $-15$

(d)  $10$

14 By discussion the increasing and decreasing intervals of the function  $f : \left[\frac{1}{e}, e\right] \rightarrow \mathbb{R}$ ,  $f(x) = x - \ln x$  we get that :

(a) the function is increasing on the intervals  $\left]\frac{1}{e}, 1\right[$  and  $]1, e[$

(b) the function is decreasing on the interval  $]1, e[$  and increasing on the interval  $\left]\frac{1}{e}, 1\right[$

(c) the function is decreasing on the interval  $\left]\frac{1}{e}, 1\right[$  and increasing on the interval  $]1, e[$

(d) the function is decreasing on the interval  $\left]\frac{1}{e}, e\right[$




- 15 A man of tall 180 cm. moves far from the base of a 3-metre lamp post at a rate of 1.2 m./sec. , then the rate of change of the length of the man's shadow = ..... m./sec.  
(a)  $\frac{5}{9}$  (b) 1.5 (c) 2.25 (d) 1.8
- 
- 16 If  $f : f(x) = \log_x e$ , then  $f'(x) =$  .....  
(a)  $\frac{1}{x}$  (b)  $-\frac{1}{x}$   
(c)  $-\frac{(\log_x e)^2}{x}$  (d)  $\frac{(\ln x)^2}{x}$
- 
- 17 If  $3y^2 \frac{dy}{dx} = 2x + 1$  and  $x = 1$  when  $y = -1$ , then the relation between  $x, y$  is .....  
(a)  $y^3 = x^2 + x - 3$  (b)  $y^3 = x^2 - 2$   
(c)  $y^3 + x^2 = 0$  (d)  $y^3 + 1 = x^2 - x$
- 
- 18 The slope of the normal to the curve  $x = e^t, y = e^{2t+2}$  at the point which lies on the curve and its  $x$ -coordinate equals one is .....  
(a)  $-2e^2$  (b)  $-\frac{1}{2e^2}$  (c)  $2e^2$  (d)  $\frac{e^2}{2}$
- 
- 19 If  $y = \cos x$ , then  $\left(\frac{dy}{dx}\right)^2 - y \frac{d^2y}{dx^2} =$  .....  
(a) zero (b) 1 (c) -1 (d)  $\cos 2x$
- 
- 20 If  $f'(x) = (x-3)(x+4)$ , then the curve of the function  $f$  is convex upward on .....  
(a)  $]-\infty, -4[$  (b)  $]-4, 3[$   
(c)  $]3, \infty[$  (d)  $]-\infty, 3[$
- 
- 21 If  $f(x) = x^{\sin x}$ , then  $f'\left(\frac{\pi}{2}\right) =$  .....  
(a) -2 (b) -1 (c) zero (d) 1
- 
- 22 If  $\int_a^b \frac{d}{dx} (3x^2 - 2x) dx = 0$ ,  $a \neq b$ , then  $a + b =$  .....  
(a) 0 (b)  $\frac{2}{3}$  (c) 1 (d)  $\frac{3}{2}$



23 In the opposite figure :

$\overline{AB}$  is a tangent segment of the circle,  $\overline{AD}$  is secant where  $DC = 5$  cm. If the rate of change of the length of  $\overline{AB}$  equals 1 cm./min. Then the rate of change of the length of  $\overline{AC}$  when  $AC = 4$  cm. equals ..... cm./min.




- (a) 2                      (b)  $\frac{1}{2}$                       (c)  $\frac{12}{13}$                       (d)  $\frac{3}{4}$

24 If  $f$  is differentiable twice function in the interval  $[-1, 1]$  where  $\hat{f}'(x)$  is increasing on  $]-1, 0[$  and  $\hat{f}'(x)$  is decreasing on  $]0, 1[$ , then the statement which is certainly correct from the following is .....

- (a)  $f(0)$  is a local maximum value of the function.  
 (b) the point  $(0, f(0))$  is an inflection point of the function.  
 (c) the function  $f$  is increasing on  $]0, 1[$   
 (d) the function  $f$  is decreasing on  $]0, 1[$

25 In the opposite figure :

If  $AB = 5$  cm. , then the area of the figure ABCD is maximum when .....



- (a)  $x = y$   
 (b)  $x = 2y$   
 (c)  $y = 2x$   
 (d)  $3x = 4y$



Answer the following questions :

1 If  $\lim_{x \rightarrow \infty} \left( \frac{x+a}{x+b} \right)^x = e^c$ , then  $c = \dots\dots\dots$

(a)  $a + b$

(b)  $a - b$

(c)  $b - a$

(d)  $\frac{a+b}{2}$

2 An empty container, its volume  $45 \text{ cm}^3$ , water is poured in it at a rate  $5 \text{ cm}^3/\text{sec.}$ , the container becomes full after  $\dots\dots\dots$  seconds.


(a) 9

(b) 135


(c) 45

(d) 5


3 If  $f(3)$  undefined,  $f'(x) > 0$  when  $x \neq 3$ , then the curve which can represent the continues function  $f$  on  $[1, 5]$  is  $\dots\dots\dots$




(a)



(b)



(c)



(d)

4 The area of the region bounded by the curve  $y = 3x^2 + 4$ ,  $x$ -axis and the two straight lines  $x = -1$ ,  $x = 2$  equals  $\dots\dots\dots$  square units.

(a) 21

(b) 11

(c) -21

(d) 16




- 5 If  $y = \sqrt{2(\sec x + \tan x)}$  where  $x \in ]0, \frac{\pi}{2}[$ , then  $\frac{1}{y} \left( \frac{dy}{dx} \right) = \dots\dots\dots$
- (a)  $2 \sec x$  (b)  $-\frac{1}{2} \sec x$  (c)  $\frac{1}{2} \sec x$  (d)  $-2 \sec x$
- 
- 6 The shortest distance between the straight line :  $x - 2y + 10 = 0$  and the curve  $y^2 = 4x$  equals ..... length unit.
- (a) 4 (b)  $6\sqrt{5}$  (c)  $\frac{6}{5}\sqrt{5}$  (d) 2
- 
- 7 If  $x \frac{dy}{dx} - y = 3$  and  $x = 1$  when  $y = -2$ , then the relation between  $x, y$  is .....
- (a)  $|y| = |x + 3|$  (b)  $|y + 3| = |x|$   
 (c)  $y + x = -1$  (d)  $y + 2x = 0$
- 
- 8  $\int \frac{2e^x + 1}{e^x} dx = \dots\dots\dots + c$  (where  $c$  is constant)
- (a)  $2x + e^{-x}$  (b)  $2x - e^{-x}$  (c)  $2 + \ln e^x$  (d)  $2 - e^x$
- 
- 9 The volume of a solid generated by revolution of the region bounded by the curve  $xy = 3$  and the two straight lines  $y = 1$ ,  $y = 3$  and  $y$ -axis a complete revolution about  $x$ -axis equals ..... cubic unit.
- (a)  $12\pi$  (b)  $8\pi$  (c)  $4\pi$  (d)  $6\pi$
- 
- 10  $f, g$  are polynomial functions,  $f(x) = cx^2 + g(x)$ ,  $g(1) = k$  and  $g'(1) = 6$  where  $c, k$  are constants. If  $(1, 5)$  is an inflection point to the curve of  $f$ , then  $k - c = \dots\dots\dots$
- (a) -11 (b) 5 (c) 11 (d) -5
- 
- 11 The function  $f : f(x) = -|x| + 1$  is decreasing in the interval .....
- (a)  $]0, \infty[$  (b)  $] -\infty, 0[$  (c)  $]1, \infty[$  (d)  $] -\infty, 1[$
- 
- 12 If  $\frac{dy}{dx} = \tan x$ ,  $\frac{dz}{dx} = \cot x$ ,  $\frac{d^2y}{dz^2} - a^2 > 0$  when  $x = \frac{\pi}{4}$ , then  $a \in \dots\dots\dots$
- (a)  $] -2, 2[$  (b)  $\mathbb{R} - [-2, 2]$  (c)  $] -\infty, 2[$  (d)  $\mathbb{R} - ] -2, 2[$



13 In the opposite figure :

If  $m(\angle ADE) = m(\angle C)$ , then  $\int y \, dx = \dots + c$

- (a)  $\tan x + \frac{1}{3} \tan^3 x$   
 (b)  $\tan x + \frac{1}{2} \tan^2 x$   
 (c)  $\tan x + \frac{1}{9} \tan^3 x$   
 (d)  $\frac{1}{3} \sec^3 x + \frac{1}{3} \tan^3 x$



14 If  $\tilde{f}(x)$  is continuous,  $f(4) = -3$ ,  $\int_0^4 f(x) \, dx = 8$ , then  $\int_0^4 x \tilde{f}(x) \, dx = \dots$

- (a) -20 (b) -13 (c) -12 (d) -7


15 Let  $y = \ln(\sin x)$ , then  $\frac{d^2 y}{dx^2} = \dots$

- (a)  $-\csc^2 x$  (b)  $\sec x$  (c)  $-\csc x \cot x$  (d)  $\sec x \tan x$

16 The opposite figure :

ABCDEF regular hexagon, its side length increase by rate  $\sqrt{3}$  cm./sec., then the rate of change of the shaded area =  $\dots$  cm<sup>2</sup>/sec. when the side length of the hexagon equals 6 cm.

- (a) 27 (b)  $27\sqrt{3}$   
 (c) 9 (d)  $9\sqrt{3}$



17 If  $f: \mathbb{R} \rightarrow \mathbb{R}$  where  $f(x) = x^3 + 3x^2 - 9x$  and  $a, b$  are the absolute minimum and maximum values of function  $f$  on the interval  $[-2, 2]$ , then  $b - a = \dots$

- (a) 17 (b) -17 (c) 27 (d) -27

18 If  $x^2 = e^{2y}$ , then  $x \left( \frac{dy}{dx} \right) \dots$

- (a) -1 (b) 1 (c)  $x$  (d)  $e^{2x}$

19 If  $y^2 = 8x$  where  $y > 0$ , then the point lies on this curve at which  $\frac{dy}{dx} = \frac{dx}{dy}$  is  $\dots$

- (a) (0, 0) (b)  $(1, 2\sqrt{2})$  (c)  $(\frac{1}{2}, 2)$  (d) (2, 4)



- 20 The tangent of the curve  $2x + 1 = \sin y$  is parallel to y-axis at the point .....
- (a)  $(\frac{\pi}{2}, 0)$       (b)  $(-\frac{1}{2}, \pi)$       (c)  $(0, \frac{\pi}{2})$       (d)  $(-\frac{1}{2}, 0)$
- 
- 21 If  $y = f(x)$ ,  $f(2) = 5$ ,  $g(x) = f(3 - x)$ , then  $g(1) = \dots\dots\dots$
- (a) 1      (b) -1      (c) 5      (d) -5
- 
- 22 The curve of the function  $f$  is convex downwards on  $\mathbb{R}$  if  $f(x)$  equals .....
- (a)  $3 - x^2$       (b)  $3 - x^3$       (c)  $3 - x^4$       (d)  $3 + x^4$
- 
- 23 If the tangent to the curve of the function  $f : f(x) = ax^3$  at  $x = 1$  is perpendicular to the tangent to the curve of the function  $g : g(x) = b \sin^2 x$  at  $x = \frac{\pi}{4}$ , then  $a \times b = \dots\dots\dots$
- (a)  $\frac{4}{3}$       (b)  $\frac{2}{3}$       (c)  $\frac{1}{3}$       (d)  $-\frac{1}{3}$
- 
- 24  $\int (\sin x + \cot x)^9 (\cos x - \csc^2 x) dx = \dots\dots\dots + c$
- (a)  $\frac{1}{2} (\cot x - \csc^2 x)^2$       (b)  $\frac{1}{10} (\sin x + \cot x)^{10}$   
(c)  $(\sin x + \cot x)^{10}$       (d)  $-\frac{1}{10} (\sin x + \cot x)^{10}$
- 
- 25 If  $f$  is continuous and even function in  $\mathbb{R}$  where  $_{-5} \int^5 f(x) dx = 2a$  and  $_0 \int^3 f(x) dx = b$ , then  $_{-5} \int^{-3} f(x) dx = \dots\dots\dots$
- (a)  $b - a$       (b)  $2a - b$       (c)  $a - b$       (d)  $b - 2a$



Answer the following questions :

1  $\lim_{x \rightarrow \infty} \left(1 + \frac{1}{x}\right)^{2x} = \dots\dots\dots$

(a) 1

(b) 2

(c) e

(d)  $e^2$

2 The curve  $y = x e^x$  at  $\dots\dots\dots$

(a)  $x = -1$  has local minimum value.

(b)  $x = -1$  has local maximum value.

(c)  $x = 0$  has local minimum value.

(d)  $x = 0$  has local maximum value.

3 The tangent to the curve  $y = 3x^2 - 5$  at the point  $(1, -2)$  also passes by the point  $\dots\dots\dots$

(a)  $(5, -2)$

(b)  $(3, 1)$

(c)  $(2, -4)$

(d)  $(0, -8)$

4 If the perimeter of a circular sector is P (where P is constant), then its surface area is maximum at  $r = \dots\dots\dots$

(a)  $\frac{P}{2}$

(b)  $\frac{1}{\sqrt{P}}$

(c)  $\sqrt{P}$

(d)  $\frac{P}{4}$

5 If  $x = e^{2t}$ ,  $y = t^3$ , then  $\frac{d^2 y}{d x^2} = \dots\dots\dots$  at  $t = 1$

(a)  $\frac{3}{2}$

(b)  $\frac{3}{2} e^{-4}$

(c) 0

(d)  $3e^2$

6 If  $f$  is a continuous even function on the interval  $[-4, 4]$ ,

$_{-4} \int^4 f(x) dx = 20$ ,  $_0 \int^2 f(x) dx = 6$ , then  $_{-4} \int^2 f(x) dx = \dots\dots\dots$

(a) 120

(b) 14

(c) 26

(d) 16

7 If  $f(x) = \cot x$ , then  $\int \left(\frac{\pi}{4}\right) = \dots\dots\dots$

(a)  $-\frac{4}{9}$

(b)  $\frac{4}{9}$

(c) 4

(d)  $\frac{9}{2}$

8 If  $f(x) = \begin{cases} 3x^2 & , \quad x \leq 1 \\ 2x+1 & , \quad x > 1 \end{cases}$ , then  $_0 \int^5 f(x) dx = \dots\dots\dots$

(a) 125

(b) 30

(c) 110

(d) 29



9 If  $f(x) = 2x^3 - 3x^2 - 12x + 12$ , then the local maximum value of the function equals .....

- (a) 2                      (b) -8                      (c) -1                      (d) 19

10 The length of each side of an equilateral triangle =  $a$ , and increase at a rate  $k$ , then the rate of increasing of its surface area equals .....

- (a)  $\frac{2}{\sqrt{3}} ak$                       (b)  $\sqrt{3} ak$                       (c)  $\frac{\sqrt{3}}{2} ak$                       (d)  $2 ak$

11 The curve of the function  $f$  is convex downwards in  $\mathbb{R}$  if  $f(x) = \dots\dots\dots$


- (a)  $3 - x^2$                       (b)  $3 - x^3$                       (c)  $3 - x^4$                       (d)  $3 + x^4$

12  $\int \tan \theta \, d\theta = \dots\dots\dots$

- (a)  $-\ln |\cos \theta| + c$                       (b)  $-\ln \cos \theta + c$                       (c)  $\ln \cos \theta + c$                       (d)  $|\ln \cos \theta| + c$

13 In the opposite figure :

The volume of the solid generated by revolving the shaded region a complete revolution about  $x$ -axis equals ..... cube units.



- (a)  $\frac{14}{3} \pi$                       (b)  $\frac{15}{2} \pi$   
(c)  $\frac{15}{2}$                       (d)  $\frac{14}{3}$

14 The area of the region bounded by the curve of the function  $y = x^3$  and the two straight lines  $y = 0$ ,  $x = 2$  equals ..... square unit.


- (a) 4                      (b) -4                      (c)  $\frac{1}{2}$                       (d) 8

15  $\frac{d^2}{dx^2} (\cos^4 x + \sin^4 x) = \dots\dots\dots$


- (a) zero                      (b)  $-2 \sin 2x$                       (c)  $-4 \cos 4x$                       (d) 1




- 16 The surface area of a sphere increases at constant rate  $6 \text{ cm}^2/\text{sec}$ . at the instant at which its radius is  $30 \text{ cm}$ , then the rate of increase of the volume of the sphere = .....  $\text{cm}^3/\text{sec}$ .
- (a) 180 (b) 40 (c) 90 (d)  $90\pi$
- 
- 17 If  $f(x) = 2 \sin \frac{x}{2} \cos \frac{x}{2}$ , then the  $1000^{\text{th}}$  derivative of  $f(x)$  equals .....
- (a)  $\sin x$  (b)  $-\sin x$  (c)  $-\cos x$  (d)  $\cos^{1000} x$
- 
- 18 If  $f''(x) = (x-3)(x+4)$ , then the curve of the function  $f$  is convex upwards on .....
- (a)  $]-\infty, -4[$  (b)  $]-4, 3[$  (c)  $]3, \infty[$  (d)  $]-\infty, 3[$
- 
- 19 The slope of the normal to the curve at any point on it  $(x, y)$  equals  $\frac{-2y}{x}$ , then the equation of the curve, given that it intercepts 3 units from the positive part of y-axis, is .....
- (a)  $2y^2 = x^2 + 18$  (b)  $2y^2 = x^2$   
 (c)  $2y^2 = \frac{1}{2}x^2 + 9$  (d)  $y^2 = \frac{1}{2}x^2 + 3$
- 
- 20 If  $f(x) = |x|$ , then  $f'(-6) = \dots\dots\dots$
- (a) 6 (b) -1 (c) 0 (d) -6
- 
- 21  $\lim_{x \rightarrow 0} (1+x)^{\frac{1}{3x}} = \dots\dots\dots$
- (a)  $\frac{1}{3}$  (b)  $e^3$  (c)  $e^{\frac{1}{3}}$  (d)  $\frac{e}{3}$
- 
- 22 If  $f''(x) < 0$ ,  $f''(x) > 0$ , for each  $x \in [a, b]$ , which of the following represents the curve of the function  $f$  in  $[a, b]$ ?




(a)



(b)



(c)



(d)



23 The greatest area of the rectangle whose perimeter equals 14 cm. equals .....  $\text{cm}^2$

(a) 10

(b) 12

(c) 12.25

(d) 49

24  $\int \frac{1}{1 - \cos^2 x} dx = \dots\dots\dots + c$

(a)  $\csc^2 x$

(b)  $\cot x$

(c)  $-\sin^{-1} x$

(d)  $-\cot x$

25 If  $\int_{-2}^3 f(x) dx = 9$  ,  $\int_5^3 f(x) dx = 4$  , then the value of :

$\int_{-2}^5 [3f(x) - 6x] dx = \dots\dots\dots$

(a) -48

(b) -58

(c) -144

(d) -147



Answer the following questions :

- 1 The function  $f : f(x) = \frac{x}{\ln x}$  is increasing in the interval .....  
 (a)  $]0, \infty[$  (b)  $]0, e[$  (c)  $]e, \infty[$  (d)  $] - \infty, \infty[$

---

- 2  $\int (4 - \csc x \cot x) dx = \dots + c$   
 (a)  $4x - \csc x$  (b)  $4x + \csc x$  (c)  $4x - \cot x$  (d)  $4x + \cot x$

---

- 3  $\lim_{x \rightarrow \infty} \left( \frac{x+6}{x+2} \right)^{x+5} = \dots$   
 (a)  $e^4$  (b)  $e^5$  (c)  $e$  (d)  $e^6$

---

- 4 The area of the greatest rectangle which can be drawn in a circle of radius 4 cm. equals .....  $\text{cm}^2$   
 (a)  $4\sqrt{2}$  (b)  $8\sqrt{2}$  (c) 32 (d) 64

---

- 5 The curve of the function  $f$  where :  $f(x) = 2x^3 + 3x^2 - 12x + 5$  has inflection point at  $x = \dots$   
 (a) -2 (b) 1 (c)  $11\frac{1}{2}$  (d)  $-\frac{1}{2}$

---

- 6 If  $\int_{-2}^2 f(x) dx = 0$ , then  $f(x) = \dots$   
 (a)  $x^2 + 1$  (b)  $x$  (c)  $x + 1$  (d)  $x - 1$

---

- 7 If  $f(x) = \sin 2x \cos 2x$ , then  $f\left(\frac{\pi}{3}\right) = \dots$   
 (a) -4 (b) 0 (c)  $4\sqrt{3}$  (d) 8

---

- 8 If  $y = (\sin x)^{\tan x}$ , then  $\frac{dy}{dx} = \dots$   
 (a)  $(\sin x)^{\tan x} (\sec^2 x \ln \sin x + 1)$  (b)  $(\tan x) (\sin x)^{\tan x - 1}$   
 (c)  $(\cos x)^{\sec^2 x}$  (d)  $\sec^2 x \ln \sin x + 1$



- 9 The height of right circular cone equals its base diameter. The rate of change of its base radius =  $\frac{1}{\pi}$  cm./sec. then the rate of change of its volume = .....  $\text{cm}^3/\text{sec}$ .  
when its base radius length = 5 cm.

(a)  $50\pi$  (b)  $\frac{250}{3}\pi$  (c) 150 (d) 50

- 10  $\int x^2 e^x dx = \dots + c$


(a)  $\frac{1}{3} x^3 e^x$

(b)  $x^2 e^x - 2x e^x + 2e^x$

(c)  $x^2 e^x - 2x e^x$

(d)  $2x e^x$

- 11 The opposite figure represents the curve of the function  $f'$ , then the curve of the function  $f$  has an inflection point at  $x = \dots$



(a) 0

(b) 1

(c) 2

(d) 3

- 12 In the opposite figure :

ABCD is a square of side length 24 cm. ,  $AF = 2AE$


, then the greatest area of the figure FBCE = .....  $\text{cm}^2$

(a) 324

(b) 252

(c) 6

(d) 648



- 13 The area of the region bounded by the curve  $y = x^3$  and the straight lines  $x = -1$  ,  $x = 1$  ,  $y = 0$  equals ..... square unit.

(a) zero

(b)  $\frac{1}{2}$

(c)  $\frac{1}{4}$

(d) 6

- 14 The equation of the curve :  $y = f(x)$  if  $y' = 6x - 4$  and the curve has local minimum at (1 , 5) is .....

(a)  $f(x) = x^3 - 2x^2 + x + 5$

(b)  $f(x) = x^3 - 2x^2 + x$


(c)  $f(x) = 3x^2 - 4x + 1$

(d)  $f(x) = 3x^2 - 4x$



- 15 The curve of the function  $f : f(x) = (x - 2)e^x$  is convex downwards in the interval .....
- (a)  $]-\infty, \infty[$  (b)  $]-1, 2[$  (c)  $]0, 2[$  (d)  $]0, \infty[$
- 
- 16 The rate of increasing of the length of each of two sides in a triangle is  $0.1 \text{ cm./sec.}$  and the rate of increasing of angle including between them is  $\frac{1}{5} \text{ rad/sec.}$ , then the rate of increasing of area of the triangle at the instant when the length of each side of the triangle is  $10 \text{ cm.}$  equals .....  $\text{cm}^2/\text{sec.}$
- (a)  $\frac{\sqrt{3}}{4}$  (b)  $\frac{\sqrt{3}}{2}$  (c)  $5$  (d)  $5.866$
- 
- 17 If  $y = \sin x + \sec x$ ,  $x = 3\pi z$ , then  $\frac{dy}{dz} = \dots\dots\dots$  at  $z = 1$
- (a)  $3\pi$  (b)  $-3\pi$  (c)  $-6\pi$  (d)  $-1$
- 
- 18 If  $f(x) = 3x - 2$ , then  $(f \circ f)(1) = \dots\dots\dots$
- (a)  $6$  (b)  $-6$  (c)  $9$  (d)  $3$
- 
- 19  $\int_{-1}^3 |x - 1| dx = \dots\dots\dots$
- (a)  $2$  (b)  $0$  (c)  $4$  (d)  $\frac{3}{2}$
- 
- 20 The volume of the solid generated by revolving the region bounded by the straight lines  $y = 3 - x$ ,  $x = 0$ ,  $y = 0$  a complete revolution about  $x$ -axis equals ..... cubic unit.
- (a)  $9\pi$  (b)  $9$  (c)  $\frac{9}{2}\pi$  (d)  $\frac{9}{2}$
- 
- 21 If  $y = \ln(\tan x)$ , then  $\frac{dy}{dx} = \dots\dots\dots$
- (a)  $2 \sec 2x$  (b)  $2 \csc 2x$  (c)  $\sec^2 x$  (d)  $\cot x$
- 
- 22 In the opposite figure :  
If  $f(x) = g(x) - 3h(x)$   
, then  $f(5) = \dots\dots\dots$

- (a) zero  
(b) 2  
(c) 3  
(d) 4





23  $\int \cos^{99} X \sec^{100} X dX = \dots\dots\dots + c$

(a)  $\frac{1}{100} \cos^{100} X$

(b)  $\frac{1}{101} \cos^{101} X$

(c)  $\sec X \tan X$

(d)  $\ln |\sec X + \tan X|$

24 If the point (1, 12) is the inflection point to the curve of the function  $f$  where  $f(X) = aX^3 + bX^2$ , then  $2a + b = \dots\dots\dots$

(a) 24

(b) -12

(c) 12

(d) 6

25 If  $y = -\sin X$ , then  $\frac{d^2 y}{dX^2} + y = \dots\dots\dots$

(a) -4

(b) 2

(c) 4

(d) zero



Answer the following questions :

1  $\lim_{x \rightarrow 0} \frac{e^x - \sin x - 1}{x} = \dots\dots\dots$

(a) zero

(b) 1

(c) undefined.

(d) -1

2 If the curve of the function  $f$  represents a polynomial function, has a local maximum at the point  $(a, b)$ , then  $f'(a) = \dots\dots\dots$

(a) b

(b) zero

(c)  $-\frac{b}{a}$

(d) undefined.

3 If the tangent to the curve  $y^2 = 4ax$  is perpendicular to  $X$ -axis, then  $\dots\dots\dots$

(a)  $\frac{dy}{dx} = 0$

(b)  $\frac{dy}{dx} = 1$

(c)  $\frac{dx}{dy} = 1$

(d)  $\frac{dx}{dy} = 0$

4 If  $x = a(\cos \theta + \theta \sin \theta)$ ,  $y = a(\sin \theta - \theta \cos \theta)$ , then  $\frac{dy}{dx} = \dots\dots\dots$

(a)  $\sin \theta$

(b)  $\sin 2\theta$

(c)  $\cos \theta$

(d)  $\tan \theta$

5 The volume of the solid generated by revolving the region bounded by the two curves  $y = \tan x$ ,  $y = \sec x$  and the two straight lines  $x = \frac{\pi}{6}$ ,  $x = \frac{\pi}{3}$  a complete revolution about  $X$ -axis is  $\dots\dots\dots$  cubic unit.

(a)  $\frac{\pi^2}{6}$

(b)  $\frac{\pi^2}{3}$

(c)  $\frac{2\pi^2}{5}$

(d)  $2\pi^2$

6 If  $f: \left[\frac{1}{e}, e\right] \rightarrow \mathbb{R}$  and  $f(x) = x - \ln x$

, then the function  $f$  has absolute maximum value =  $\dots\dots\dots$

(a) e

(b)  $e - 1$

(c) 1

(d)  $\frac{1}{e} + 1$

7 The rate of change for  $\sqrt{x^2 + 16}$  with respect to  $\frac{x}{x-1}$  at  $x = 3$  equals  $\dots\dots\dots$

(a) -60

(b)  $-\frac{5}{12}$

(c)  $-\frac{12}{5}$

(d)  $-\frac{3}{5}$

8  $\int \frac{\ln x^2}{\ln x} dx = \dots\dots\dots + c$  (where :  $x \in \mathbb{R}^+ - \{1\}$ )

(a)  $\frac{x}{2}$

(b)  $\frac{1}{x}$

(c)  $2x$

(d)  $\ln|x|$



9 If  $f(x) = 2x^2 + x - 3$ , then  $\int_{-1}^2 f(x) dx = \dots\dots\dots$

- (a) 8                      (b) 9                      (c) 10                      (d) 11

10 If the rate of change in volume of a sphere equals the rate of change of its radius, then  $r = \dots\dots\dots$  length unit.

- (a) 1                      (b)  $\sqrt{2\pi}$                       (c)  $\frac{1}{\sqrt{2\pi}}$                       (d)  $\frac{1}{2\sqrt{\pi}}$

11  $\int x \cos x dx = \dots\dots\dots + c$

- (a)  $x \sin x - \cos x$                       (b)  $x$   
(c)  $-\frac{1}{2} x^2 \sin x$                       (d)  $x \sin x + \cos x$

12 The curve  $y = (2x - c)^3 + 4$  has an inflection point at  $x = 5$ , then  $c = \dots\dots\dots$

- (a) 2                      (b) 4                      (c) 5                      (d) 10

13 If  $y = e^x$ ,  $z = \sin x$ , then  $\frac{dy}{dz} = \dots\dots\dots$

- (a)  $\frac{e^x}{\sin x}$                       (b)  $e^x \tan x$                       (c)  $e^x \cos x$                       (d)  $\frac{e^x}{\cos x}$

14 If  $f(x) = 2x^3 - 3x^2 - 36x + 14$ , then the curve of the function is convex downwards on the interval  $\dots\dots\dots$

- (a)  $]\frac{1}{2}, \infty[$                       (b)  $]-\infty, \frac{1}{2}[$                       (c)  $]-2, 3[$                       (d)  $\mathbb{R} - [-2, 3]$

15 The slope of the tangent to the curve at any point on it  $(x, y)$  is given by the relation

$\frac{dy}{dx} = \sin x \cos x$ , then the equation of the curve known that it passes through

the point  $(\frac{\pi}{6}, 1)$  is  $\dots\dots\dots$

- (a)  $y = \frac{1}{2} \sin^2 x$                       (b)  $y = \frac{1}{2} \sin^2 x + 7$   
(c)  $y = \frac{1}{2} \sin^2 x + \frac{7}{8}$                       (d)  $2y = \sin^2 x - \frac{7}{4}$




- 16 A trapezium is drawn in a semi-circle, and its base is the diameter of the semi-circle, then the base angle of the trapezium such that its area is as maximum as possible is of measure .....


(a)  $45^\circ$  (b)  $60^\circ$  (c)  $30^\circ$  (d)  $120^\circ$

- 17 The opposite figure represents the curve of function  $f$ , then all the following statements are true except .....


(a)  $f(4) = 6$   
 (b)  $f'(4) = 0$   
 (c)  $f''(x) > 0$  for  $x < 4$   
 (d)  $f''(x) < 0$  for  $x < 4$




- 18 If  $y = ax^n - bx^{n-1}$  is a polynomial function,  $a, b \in \mathbb{R}$ , then  $\frac{d^n y}{dx^n}$  could be represented by the figure .....




(a)



(b)



(c)




(d)

- 19 In the opposite figure :

If the area of the shaded region = 9 square units

, then  $\int_0^4 f(x) dx = \dots$

(a) 24 (b) 4  
 (c) 15 (d) 33



- 20  $\int_{2020}^{2022} (x - 2021)^2 dx = \dots$

(a)  $\frac{1}{3}$  (b)  $\frac{2}{3}$  (c) 1 (d)  $\frac{4}{3}$



21  $\int \frac{\sin^{10} x}{\cos^{12} x} dx = \dots\dots\dots + c$

- (a)  $\tan^{11} x$       (b)  $\frac{1}{11} \tan^{11} x$       (c)  $\frac{1}{11} \tan 11 x$       (d)  $\sec^2 x$

22 ABCD is a square whose side length 10 cm. and  $M \in \overline{BC}$  where  $BM = x$  cm. and  $N \in \overline{CD}$  where  $CN = \frac{3}{2} x$ , then the value of  $x$  which makes the area of  $\Delta AMN$  as minimum as possible =  $\dots\dots\dots$  cm.

- (a)  $\frac{10}{3}$       (b)  $\frac{3}{2}$       (c) 5      (d)  $\frac{15}{2}$

23  $\int \frac{2x^3}{x^4 + 5} dx = \dots\dots\dots + c$

- (a)  $8 \ln |x^4 + 5|$       (b)  $2 \ln |x| + \frac{1}{10} x^4$   
 (c)  $2 \ln |x^4 + 5|$       (d)  $\frac{1}{2} \ln |x^4 + 5|$

24 The tangent to the curve :  $x^2 - xy + y^2 = 27$  drawn at the point (6, 3) makes an angle of measure  $\dots\dots\dots$  with the positive direction of the  $x$ -axis.

- (a)  $90^\circ$       (b) zero      (c)  $45^\circ$       (d)  $180^\circ$

25 The side length of a square is 5 cm. The side length increases at a rate 4 cm./sec. , then the length of the side of the square after  $t$  seconds is given by the relation  $\dots\dots\dots$

- (a)  $4t$       (b)  $4t + 5$       (c)  $4t - 5$       (d) 9

**Answer the following questions :**

- 1 The equation of the normal to the curve  $y = f(x)$  at the point  $(1, 1)$  is  $x + 4y = 5$ , then  $f'(1) = \dots\dots\dots$
- (a)  $-3$                       (b)  $-\frac{1}{4}$                       (c)  $4$                       (d)  $-4$
- 
- 2 If  $y = \ln(\sec x + \tan x)$ , then  $\frac{dy}{dx} = \dots\dots\dots$
- (a)  $\tan x$                       (b)  $\sec x$                       (c)  $\tan^2 x$                       (d)  $\csc x$
- 
- 3 If  $z = x + \frac{1}{x}$ , then  $dz = \dots\dots\dots$
- (a)  $(1 + \frac{1}{x^2}) dx$                       (b)  $(1 + \frac{1}{x^2}) dx + c$                       (c)  $(1 - \frac{1}{x^2}) dx + c$                       (d)  $(1 - \frac{1}{x^2}) dx$
- 
- 4  $\int_{-2}^2 (ax^3 + bx + c) dx$  depends on  $\dots\dots\dots$
- (a) the value of  $b$                       (b) the value of  $c$   
(c) the value of  $a$                       (d) the values of  $a, b$
- 
- 5 If  $\lim_{x \rightarrow 0} \frac{\ln(1+ax)}{bx} = -1$ , then  $a + b = \dots\dots\dots$
- (a) zero                      (b)  $-1$                       (c)  $2$                       (d)  $-2$
- 
- 6 If  $f'(x) = x \cdot f(x)$ ,  $f(3) = -5$ , then  $f''(3) = \dots\dots\dots$
- (a)  $-50$                       (b)  $-40$                       (c)  $15$                       (d)  $27$
- 
- 7 The absolute minimum value of the function  $f : f(x) = xe^{-x}$  in the interval  $[0, 2]$  equals  $\dots\dots\dots$
- (a)  $1$                       (b)  $\frac{1}{e}$                       (c) zero                      (d)  $\frac{2}{e^2}$
- 
- 8 If  $x^2 y^3 = 108$ , and  $\frac{dx}{dt} = 2$  at  $x = 2$ ,  $y = 3$ , then  $\frac{dy}{dt} = \dots\dots\dots$
- (a)  $2$                       (b)  $-2$                       (c)  $-1$                       (d)  $18$



- 9  $\int e^{(x^3 + 2 \ln x)} dx = \dots + c$   
 (a)  $3e^{x^3}$  (b)  $\frac{1}{3}e^{x^3}$  (c)  $e^{x^3}$  (d)  $x^2e^{x^3}$
- 
- 10 The curve of the function  $f : f(x) = x^3 - 12x$  is convex upwards in the interval .....  
 (a)  $]-\infty, 0[$  (b)  $]0, \infty[$  (c)  $\mathbb{R} - \{0\}$  (d)  $\mathbb{R} - ]-2, 2[$
- 
- 11 The dimensions of a rectangle which has the greatest area can be drawn in a triangle , the base length of the triangle equals 16 cm. and its height 12 cm. such that one of the rectangle sides coincides with the base of the triangle and its opposite vertices lie on the other two sides of the triangle are .....  
 (a) 6 cm. , 6 cm. (b) 8 cm. , 8 cm. (c) 6 cm. , 8 cm. (d) 4 cm. , 6 cm.
- 
- 12 The area of the region bounded by the curve  $y = \sqrt{4 - x^2}$  and  $x$ -axis in square unit equals ..... square units.  
 (a) 2 (b) 4 (c)  $2\pi$  (d)  $4\pi$
- 
- 13 The equation of the curve passes through the point A (2 , 3) and the slope of the normal at any point on it is  $3 - x$  .....  
 (a)  $y = \ln |x - 3|$  (b)  $y = (x - 3)^{-2}$   
 (c)  $y = \ln |x - 3| + 3$  (d)  $y = \ln |x - 3| - 3$
- 
- 14 A regular quadrilateral pyramid of metal expands uniformly , the height equals the side length of its base , its volume increases at a rate  $1 \text{ cm}^3/\text{sec}$  , when the rate of increasing of each of its height and its base side equals  $0.01 \text{ cm}/\text{sec}$  , then the base length at this moment = .....  
 (a) 10 cm. (b) 100 cm. (c) 5 cm. (d) 125 cm.
- 
- 15 If  $y \in ]0, \frac{\pi}{4}[$  ,  $x = \frac{2 \tan y}{1 - \tan^2 y}$  , then  $\frac{dy}{dx} = \dots$   
 (a)  $\frac{1}{2} \cos^2 2y$  (b)  $2 \sec^2 2y$  (c)  $\sin^2 2y$  (d)  $\cot 2y$



- 16 If the function  $y = a x^3 + b x^2 + c x + d$  has a critical point at  $(1, 5)$ , then  $2a + b - d = \dots\dots\dots$

(a)  $-6$  (b)  $-5$  (c)  $5$  (d)  $6$

- 17 If  $x = \frac{t}{1+t}$ ,  $y = \frac{t+1}{t}$ , then  $\frac{d^2 y}{d x^2} = \dots\dots\dots$


(a)  $\frac{2}{x}$  (b)  $2x^{-3}$  (c)  $-x^{-2}$  (d) zero

- 18 If  $f: f(x) = 15x + 6x^2 - x^3$ , then the function  $f$  has  $\dots\dots\dots$

(a) an inflection point  $(2, 46)$  (b) two inflection points at  $x = -1, x = 5$   
(c) no inflection point. (d) an inflection point  $(0, 2)$

- 19 The opposite figure represents the curve of  $\ddot{f}(x)$ , then the function  $f$  is convex upwards on the interval  $\dots\dots\dots$

(a)  $]1, 3[$  (b)  $\mathbb{R} - [1, 3]$   
(c)  $] -\infty, 2[$  (d)  $]2, \infty[$




- 20 If  $f(x)$  is a continuous function on  $[-5, 8]$  and  $_{-5} \int^4 f(x) dx = 19$ ,  $\int_8^4 f(x) dx = 7$ , then  $_{-5} \int^8 f(x) dx = \dots\dots\dots$

(a)  $26$  (b)  $12$  (c)  $-12$  (d)  $-26$

- 21 In the opposite figure :

The volume of the solid generated by revolving the shaded area a complete revolution about the  $x$ -axis =  $\dots\dots\dots$  cube units.

(a)  $\frac{\pi}{3}$  (b)  $\frac{\pi}{2}$   
(c)  $\frac{2\pi}{3}$  (d)  $\pi$





22 If the equation of the tangent to the curve of the function  $f : f(x) = ax^3 + 2\sqrt{x}$  is  $y = 4x - 1$  at  $x = 1$ , then  $a = \dots\dots\dots$

(a) 1

(b) 2

(c)  $\sqrt{2}$ 

(d) 4

23 A ladder of length 10 m. rests with its upper end on a vertical wall and its lower end on a horizontal ground. If the lower end slipping away from the wall at speed 2 m./min, then the rate of change in the inclination angle of the ladder to the horizontal at the moment the lower end is 8 m. from the wall equals  $\dots\dots\dots$  rad./min.

(a) 3

(b) -3

(c)  $\frac{1}{3}$ (d)  $-\frac{1}{3}$ 

24  $\int \tan x \, dx = \dots\dots\dots + c$

(a)  $\ln |\cos x|$ (b)  $-\ln |\sec x|$ (c)  $\sec^2 x$ (d)  $\ln |\sec x|$ 

25 In the opposite figure :


The least length of the line segment  $\overline{OA} = \dots\dots\dots$  length unit.

(a)  $\sqrt{2}$ 

(b) 2

(c)  $2\sqrt{2}$ 

(d) 4



**Answer the following questions :**

- 1 The straight line  $y + x - 1 = 0$  touches the curve of the function  $f : f(x) = x^2 - 3x + a$ , then  $a = \dots\dots\dots$
- (a) 1                                      (b) 2                                      (c) 3                                      (d) 4
- 
- 2 If  $\int_{-2}^3 f(x) dx = 12$  ,  $\int_{-2}^5 f(x) dx = 16$  , then  $\int_3^5 f(x) dx = \dots\dots\dots$
- (a) -28                                      (b) -4                                      (c) 4                                      (d) 28
- 
- 3 The local minimum value of the function  $f$  , where :  $f(x) = x^4 - 2x^2$  equals  $\dots\dots\dots$
- (a) 1                                      (b) -1                                      (c) 0                                      (d) -4
- 
- 4 If  $\int (2x - 1) e^{2x+3} dx = yz - \int z dy$  , then  $\int z dy = \dots\dots\dots$
- (a)  $e^{2x+3} + c$                                       (b)  $\frac{1}{2} e^{2x+3} + c$   
(c)  $-e^{2x+3} + c$                                       (d)  $-\frac{1}{2} e^{2x+3} + c$
- 
- 5  $\int \frac{x^3 dx}{x^4 + 3} = \dots\dots\dots + c$
- (a)  $\frac{1}{4} (x^4 + 3)$                                       (b)  $\frac{1}{4} \ln |x^4 + 3|$                                       (c)  $\ln |x^4 + 3|$                                       (d)  $\frac{1}{4} (x^4 + 3)^{-1}$
- 
- 6 In the function  $f$  : where  $f(x) = \frac{x^2 + 9}{x}$  , the absolute minimum value of the function  $f$  where  $x \in [1, 6]$  equals  $\dots\dots\dots$
- (a) 10                                      (b) 6                                      (c) 7.5                                      (d) zero
- 
- 7  $\int \sec^5 x \tan x dx = \dots\dots\dots + c$
- (a)  $\frac{1}{6} \sec^6 x$                                       (b)  $\frac{1}{5} \sec^5 x$                                       (c)  $\sec^7 x$                                       (d)  $\frac{1}{2} \tan^2 x$
- 
- 8  $\frac{d^3}{dx^3} (\sin^2 x) = \dots\dots\dots$
- (a)  $\sin 2x$                                       (b)  $2 \cos 2x$                                       (c)  $4 \cos 2x$                                       (d)  $-4 \sin 2x$




- 9 When the region bounded by the curve  $x = \frac{1}{\sqrt{y}}$ ,  $1 \leq y \leq 4$  and y-axis revolves a complete revolution about y-axis, then the volume of the solid generated in cubic units equals .....
- (a)  $\frac{2}{3} \pi$  (b)  $3\sqrt{2} \pi$  (c)  $2 \pi \ln 2$  (d)  $\frac{2}{3} \pi \log 3$
- 
- 10 On the perpendicular coordinate system a straight line  $\overleftrightarrow{AB}$  passes through the point C (3, 2) and intersects the positive part of x-axis at the point A and the positive part of y-axis at the point B, then the smallest area of the triangle AOB equals ..... square unit (where O is the origin)
- (a) 12 (b) 6 (c) 3 (d) 24
- 
- 11 A ladder of length two metres is leaning against a smooth vertical wall. If the top of the ladder slid down at the same rate as the lower end slid away from the wall, then the distance of the lower end from the wall equals ..... m.
- (a) 2 (b)  $2\sqrt{2}$  (c)  $\sqrt{2}$  (d)  $-\sqrt{2}$
- 
- 12 If  $f: \mathbb{R} \rightarrow \mathbb{R}$  where  $f(x) = \sqrt[3]{x^2}(3x - 7)$  and the function is increasing for  $x \in ]-\infty, a[$ ,  $x \in ]b, \infty[$ , then  $7a + 30b = \dots\dots\dots$
- (a) 14 (b) 28 (c) -28 (d)  $\frac{98}{15}$
- 
- 13 The slope of the normal to the curve:  $x = \cos \theta$ ,  $y = \sqrt{2} + \sin \theta$  at  $\theta = \frac{\pi}{4}$  is .....
- (a) 1 (b) -1 (c) zero (d) undefined.
- 
- 14  $\frac{d}{dx} \left( 2 \sin \frac{\pi}{8} \cos \frac{\pi}{8} \right) = \dots\dots\dots$
- (a)  $\sin \frac{\pi}{4}$  (b)  $\cos \frac{\pi}{4}$  (c)  $\frac{1}{4} \cos \frac{\pi}{4}$  (d) zero
- 
- 15 If  $f(x) = \frac{x^{65}}{65}$ , then  $f^{(65)}(x) = \dots\dots\dots$
- (a) zero (b) 1 (c) x (d)  $\frac{x}{65}$



- 16 The opposite figure represents the function  $f$  and the straight line  $L$  touches the curve of  $f$  at the point

$A(3, 5)$  and  $g(x) = x \cdot f(x)$ , then  $g'(3) = \dots\dots\dots$

- (a) 3 (b) 1  
(c) 5 (d) 8



- 17 The height of a right circular cone equals its base diameter, the rate of change of its base radius  $= \frac{1}{\pi}$  cm./sec., then the rate of change of its volume  $= \dots\dots\dots$  cm<sup>3</sup>/sec. when its base radius  $= 5$  cm.

- (a)  $50\pi$  (b)  $\frac{250}{3}\pi$  (c) 150 (d) 50

- 18  $\lim_{x \rightarrow 0} \frac{6^x - 1}{3x} = \dots\dots\dots$

- (a)  $3 \ln 8$  (b)  $3 \ln 3$  (c)  $\frac{1}{3} \ln 6$  (d)  $\frac{1}{3} e^6$

- 19 If  $y^x = x$ , then  $\frac{dy}{dx} = 0$  at  $x = \dots\dots\dots$

- (a) zero (b) 2 (c)  $e$  (d)  $\frac{1}{e}$

- 20  $\int \frac{dx}{e^x + e^{-x} + 2} = \dots\dots\dots + c$

- (a)  $\frac{1}{e^x + 1}$  (b)  $-\frac{1}{e^x + 1}$  (c)  $\frac{2}{e^x + 1}$  (d)  $-\frac{2}{e^x + 1}$

- 21 The function  $f : f(x) = x^4 - 4x^2$  has  $\dots\dots\dots$

- (a) local minimum value and two local maximum values.  
(b) two different local minimum values and one local maximum value.  
(c) two local minimum values and no local maximum value.  
(d) two equal local minimum values and one local maximum value.




22 In the opposite figure :

$B \in$  the curve  $y = x^2 - 5x + 14$

, then the least perimeter of the rectangle

OABC equals ..... length units.

- (a) 10 (b) 12  
(c) 16 (d) 20



23 If  $f(x) = \begin{cases} 2x-1 & , -1 \leq x \leq 2 \\ 3 & , 2 < x < 5 \end{cases}$ , then  $_{-1} \int^4 f(x) dx = \dots\dots\dots$

- (a) 4 (b) 5 (c) 6 (d) 7


24 In the opposite figure :

$A_1 = 2$  square units ,

$A_2 = 7$  square units

, then  $_{-2} \int^2 f(x) dx = \dots\dots\dots$

- (a) 5 (b) 9  
(c) 15 (d) 19



25  $\int (\sin^2 x + \sin^2 x \tan^2 x) dx = \dots\dots\dots + c$

- (a)  $\sin^2 x + \csc^2 x$  (b)  $\tan x - x$  (c)  $\tan^2 x$  (d)  $\sec x$




## Answer the following questions :

- 1 The rate of change of tangent slope of the function  $f : f(x) = 2x^3$  at  $x = 3$  equals .....  
 (a) 54 (b) 36 (c) 18 (d) 9
- 2 The function  $f : f(x) = x^x$  has a stationary point at  $x =$  .....  
 (a) e (b)  $\frac{1}{e}$  (c) 1 (d)  $\sqrt{e}$
- 3  $\frac{d}{dx} \int_2^3 x\sqrt{x^2+1} dx =$  .....  
 (a) -1 (b) zero (c) 1 (d) 2
- 4 The shortest distance between the point (0, 5) and the curve  $y = \frac{1}{2}x^2 - 4$  equals ..... length units.  
 (a) 4 (b) zero (c) 17 (d)  $\sqrt{17}$
- 5 If  $y = \sin^3 \theta$ ,  $z = \cos^3 \theta$ , then  $\frac{dy}{dz} =$  .....  
 (a)  $-\sin \theta$  (b)  $\cos \theta$  (c)  $-\tan \theta$  (d)  $3 \sin 2\theta$
- 6 The slope of the tangent to the curve at a point  $(x, y)$  which lies on it is  $x\sqrt{x+1}$ , then the equation of the curve given that it passes through  $(0, \frac{11}{15})$  is .....  
 (a)  $15y = 15x + 11$  (b)  $y = \frac{2}{5}(x+1)^{\frac{5}{2}} - \frac{2}{3}(x+1)^{\frac{3}{2}}$   
 (c)  $y = \frac{5}{2}(x+1)^{\frac{5}{2}} - \frac{3}{2}(x+1)^{\frac{3}{2}} + 1$  (d)  $y = \frac{2}{5}(x+1)^{\frac{5}{2}} - \frac{2}{3}(x+1)^{\frac{3}{2}} + 1$

## 7 In the opposite figure :

ABC is isosceles triangle in which  $AB = AC = 13$  cm.,  $BC = 10$  cm.  
 If the straight line L moves from point B parallel to  $\overrightarrow{AC}$  in direction of  $\overrightarrow{BC}$  to intersects  $\overline{AB}$ ,  $\overline{BC}$  at D, E respectively where the rate of change in  $BE = \frac{1}{10}$  cm./sec., then the rate of change of the area of  $\triangle DEB$  at  $BE = 1$  cm. equals .....  $\text{cm}^2/\text{sec}$ .



- (a) 0.48 (b) 0.12 (c) 0.96 (d) 0.24



- 8 The local maximum value of the curve :  $y = \sin x (1 + \cos x)$  where  $x \in ]0, \frac{\pi}{2}[$  equals .....

(a)  $-3\sqrt{3}$  (b)  $\frac{3}{4}\sqrt{3}$  (c)  $\frac{1}{2}$  (d)  $\frac{\pi}{3}$

- 9 If  $y = \pi^{\sin x} + e^{\pi}$ , then  $\frac{dy}{dx} = \dots\dots\dots$

(a)  $\pi^{\cos x}$  (b)  $\sin x \times \pi^{\sin x - 1}$   
(c)  $\pi^{\sin x} \cos x$  (d)  $\pi^{\sin x} \cos x \ln \pi$

- 10  $\int 2 \cos^2 x dx = \dots\dots\dots + c$

(a)  $1 + \frac{1}{2} \sin 2x$  (b)  $x + \frac{1}{2} \sin 2x$   
(c)  $1 - 2 \sin 2x$  (d)  $x + \sin 2x$

- 11 If the function  $f : f(x) = x^2 + \frac{b}{x}$  has a critical point at  $x = 2$ , then  $b = \dots\dots\dots$

(a) 16 (b) 4 (c) -1 (d)  $\frac{1}{16}$

- 12 If  $\int \frac{dx}{1 - \sin x} = a \tan x + b \sec x + c$ , then  $a^2 + b^2 = \dots\dots\dots$


(a) 2 (b) 1 (c) 4 (d) 5

- 13 In the opposite figure :

The area of the region bounded by the two curves

$y = x^2$ ,  $y = |x|$  equals .....

(a)  $2 \int_{-1}^0 (x^2 - x) dx$  (b)  $\int_0^1 (x - x^2) dx$   
(c)  $2 \int_0^1 (x - x^2) dx$  (d)  $\int_{-1}^1 (x - x^2) dx$



- 14 If  $y = \ln \sqrt{\tan x}$ , then  $\frac{dy}{dx} = \dots\dots\dots$  at  $x = \frac{\pi}{4}$

(a) 1 (b) zero (c)  $\frac{1}{2}$  (d)  $\infty$

- 15 If  $f(x) = 8x \sin x \cos x \cos 2x$ , then  $f\left(\frac{\pi}{8}\right) = \dots\dots\dots$

(a) -2 (b) zero (c) 1 (d) 2



- 16 If  $x = (1 - \sqrt[3]{y})(1 + \sqrt[3]{y} + \sqrt[3]{y^2})(1 + y + y^2)$ , then  $\frac{dy}{dx} = \dots\dots\dots$   
 (a)  $3y^2$  (b)  $-3y^2$  (c)  $-\frac{1}{3y^2}$  (d)  $\frac{1}{3y^2}$
- 17 The length of the intercepted part of y-axis by the tangent to the curve  $y = x \sin x$  at  $x = \pi$  equals  $\dots\dots\dots$  length unit.  
 (a)  $-\pi$  (b)  $\pi$  (c)  $-\pi^2$  (d)  $\pi^2$
- 18 A cuboid of dimensions 3, 4 and 12 cm. If the rate of the increase of its first dimension is 2 cm./sec. and the second dimension is 1 cm./sec. but the rate of the decrease of its third dimension is 3 cm./sec., then the rate of change of its volume at the end of two second equals  $\dots\dots\dots$  cm<sup>3</sup>/sec.  
 (a)  $-12$  (b) 468 (c) 144 (d) 252
- 19  $\lim_{x \rightarrow 0} \frac{\sin 4x}{3^x - 1} = \dots\dots\dots$   
 (a)  $\frac{4}{\ln 3}$  (b)  $\frac{4}{e^3}$  (c)  $4e^3$  (d)  $\frac{1}{\ln 3}$
- 20 The absolute maximum value of the function  $f : f(x) = x + \frac{1}{x}$  in the interval  $[\frac{1}{2}, 3]$  equals  $\dots\dots\dots$   
 (a)  $2\frac{1}{2}$  (b)  $4\frac{1}{4}$  (c) 3 (d)  $3\frac{1}{3}$
- 21 ABC is a right-angled triangle at B in which :  $AB + BC = 20$  cm., then the greatest possible area for this triangle equals  $\dots\dots\dots$  cm<sup>2</sup>  
 (a) 50 (b) 10 (c) 150 (d) 100
- 22  $\int x(x-5)^3 dx = \dots\dots\dots + c$   
 (a)  $\frac{1}{5}(x-5)^5 + \frac{5}{4}(x-5)^4$  (b)  $4x(x-5)^4 + 20(x-5)^5$   
 (c)  $\frac{1}{4}(x^2 - 5x)^4$  (d)  $4(x-5)^3 + 15(x-5)^2$
- 23 If  $f(x) = \begin{cases} x^2 & , x < 2 \\ 3x - 2 & , x \geq 2 \end{cases}$ , then  $\int_0^3 f(x) dx = \dots\dots\dots$   
 (a)  $\frac{49}{6}$  (b) 9 (c)  $\frac{15}{2}$  (d)  $\frac{25}{3}$



24 The volume of the solid generated by revolution the region bounded by the curve :  $y^2 = 4 - x$  and the two positive parts of the coordinate axes a complete revolution about  $y$ -axis equals ..... volume units.

(a)  $\frac{256}{15} \pi$

(b)  $\frac{16}{3} \pi$

(c)  $\frac{1472}{15} \pi$

(d)  $8 \pi$

25  $\int \sin^2 x \, dx = \dots\dots\dots + c$

(a)  $\frac{1}{2} x - \frac{1}{2} \sin x$

(b)  $\frac{1}{3} \sin^3 x$

(c)  $-\cos^2 x$

(d)  $\frac{1}{2} x - \frac{1}{4} \sin 2x$



Answer the following questions :

1 The tangent to the curve of the function  $y = \sqrt[3]{x}$  at  $x = 0$  is parallel to .....

- (a)  $x$ -axis. (b)  $y$ -axis.  
(c) the straight line  $y = x$  (d) the straight line  $x + y = 0$

2 If  $\int_1^3 f(x) dx = 5$  ,  $\int_3^4 f(x) dx = 2$  ,  $\int_2^4 f(x) dx = 6$   
 , then  $\int_2^1 f(x) dx = \dots\dots\dots$

- (a) 1 (b) 13 (c) -2 (d) -1

3  $\lim_{x \rightarrow 6} \frac{e^x - e^6}{x - 6} = \dots\dots\dots$

- (a) 1 (b) -1 (c) zero (d)  $e^6$

4 The maximum local value of the function  $y = \frac{\ln x}{x}$  in the interval  $[2, \infty[$  is .....

- (a) 1 (b)  $\frac{2}{e}$  (c)  $e$  (d)  $\frac{1}{e}$

5 The function  $f : f(x) = \frac{x^2 + 1}{x^2 + 3}$  is convex downward on the interval .....

- (a)  $] -1, 1[$  (b)  $] -\infty, -1[ , ] 1, \infty[$   
(c)  $] 0, \infty[$  (d)  $] -\infty, 0[$

6 If  $x^2 y = 2x + 5$  , then :  $x^2 \frac{d^2 y}{dx^2} + 4x \frac{dy}{dx} = \dots\dots\dots$

- (a)  $-2y$  (b)  $-y$  (c)  $2y$  (d)  $y$

7 When the region bounded by the curve  $y = x^2$  and the straight line  $y = 2$  revolves a complete revolution about  $y$ -axis , then the volume of the generated solid equals .....

- (a)  $\int_0^2 y dx$  (b)  $\pi \int_0^2 y dy$  (c)  $\pi \int_0^2 x dx$  (d)  $\pi \int_0^2 x^2 dx$



8 An equilateral triangle, its side length increases at a rate  $\frac{1}{3}$  cm./sec., then the rate of change of its perimeter at this instant equals ..... cm./sec.

- (a) 1 (b) 2 (c) 3 (d) 4

9 If  $x = 2t^3 + 3$ ,  $y = t^4$ , then  $\frac{d^2y}{dx^2} = \dots\dots\dots$  at  $t = 1$

- (a) 9 (b)  $\frac{1}{9}$  (c)  $\frac{2}{3}$  (d)  $\frac{1}{3}$

10  $\int \cot^3 x \, dx = \dots\dots\dots + c$


- (a)  $\frac{1}{4} \cot^4 x$  (b)  $(\ln |\sin x|)^3$   
(c)  $-\frac{1}{2} \cot^2 x + \ln |\csc x|$  (d)  $\frac{1}{2} \cot^3 x + \csc^2 x$

11 The function  $f : f(x) = x^3 + 4x + 2$  is increasing for every  $x \in \dots\dots\dots$

- (a)  $\mathbb{R}^+$  (b)  $\mathbb{R}$  (c)  $\mathbb{R}^-$  (d)  $\mathbb{R} - \{0\}$

12 The opposite figure represents the curve  $f'(x)$ , then the function  $f$  has a local minimum at  $x = \dots\dots\dots$

- (a) 1  
(b) 2  
(c) 3  
(d) 4



13 A metallic circular sector whose area is  $16 \text{ cm}^2$ , then the radius length of the sector's circle which makes its perimeter as minimum as possible equals ..... cm.

- (a) 4 (b) 8 (c) 64 (d) 2

14 The area of the region bounded by the two curves :  $y = 2x^2$ ,  $y = 3 - x^2$  equals ..... square unit.

- (a) 2 (b) 4 (c) 0 (d) 8

15 If the slope of the normal to the curve at any point  $(x, y)$  on it equals  $\frac{5+2y}{2-3x^2}$  and the curve passes through  $(1, 2)$ , then its equation is .....

- (a)  $y^2 + 5y = x^3 - 3x$  (b)  $y^2 + 5y = x^3 - 2x + 15$   
(c)  $2y + 5 = 3x^2 - 12$  (d)  $y^2 + 5y = x^3 - 2x - 15$



- 16 The equation of the tangent to the curve :  $2 + \ln y \cdot \ln x = x^2 + y$  at the point whose  $x$  coordinate equals 1 is .....
- (a)  $2x + y = 0$       (b)  $2x + y = 3$       (c)  $-2x + y = 3$       (d)  $x - 2y = 3$
- 
- 17  $\frac{d}{dx} [(\sec x - 1)(\sec x + 1)] = \dots\dots\dots$
- (a)  $\sec^2 x \tan^2 x$       (b)  $2 \sec^2 x \tan x$   
(c)  $\sec^2 x \tan x$       (d)  $\sec^4 x$
- 
- 18 The straight line :  $13x - y - 7 = 0$  touches the curve  $y = ax^3 + bx^2$  at the point (1, 6), then the value of  $a$   $b = \dots\dots\dots$
- (a) 1      (b) 6      (c) 5      (d) 13
- 
- 19 A man walk across a bridge, 12 m. high above water surface, the speed of the man is 3 m./min., the man observed a boat moving perpendicular to the bridge with a constant speed 6 m./min. exactly under the man, then the rate of diverge between the man and the boat after 6 minutes from the moment that they are on the same vertical line = ..... m./minute
- (a)  $\frac{45}{7}$       (b) 540      (c)  $\frac{2700}{7}$       (d)  $3\sqrt{5}$
- 
- 20 If  $x e^{xy} = y + \sin^2 x$ , then :  $\frac{dy}{dx}$  (at  $x = 0$ ) equals .....
- (a) -1      (b) -2      (c) 2      (d) 1
- 
- 21 The area of the largest rectangle that can be inscribed in a circle of radius 4 cm. equals .....  $\text{cm}^2$
- (a) 32      (b)  $4\sqrt{2}$       (c) 8      (d) 64
- 
- 22  $\int x^3 e^{x^2} dx = \dots\dots\dots + c$
- (a)  $\frac{1}{2} x^2 e^{x^2} - x e^{x^2}$       (b)  $\frac{1}{2} x^2 e^{x^2} + \frac{1}{2} e^{x^2}$   
(c)  $2 x^2 e^{x^2} - 4 e^{x^2}$       (d)  $\frac{1}{2} x^2 e^{x^2} - \frac{1}{2} e^{x^2}$



23 If  $f(x) = \begin{cases} 2 & , \quad x < 2 \\ x & , \quad x \geq 2 \end{cases}$ , then  $\int_0^6 f(x) dx = \dots\dots\dots$

- (a) 18                      (b) 20                      (c) 12                      (d) 24

24 An architect has designed an arc-like entryway of a hotel whose equation  $y = -\frac{1}{2}(x-1)(x-7)$  where  $x$  in metres. How much does the glass cost if this entryway is covered by the glass which costs L.E. 1500 per square metre?

- (a) L.E. 27000                      (b) L.E. 18  
(c) L.E. 13500                      (d) L.E. 54000

25 If  $g$  is an increasing function on  $\mathbb{R}$ ,  $k$  is a decreasing function on  $\mathbb{R}$  and  $f(x) = 4g(x) - 3k(x)$ , then the function  $f$  ..... on  $\mathbb{R}$

- (a) increases                      (b) decreases  
(c) is constant                      (d) is a zero



Answer the following questions :

1 The function  $f : f(x) = x^3 + 4x + 2$  is increasing, then  $x \in \dots\dots\dots$

(a)  $]-4, \infty[$  only.

(b)  $\mathbb{R}$

(c)  $]-\infty, -\frac{4}{3}[$  only.

(d)  $]-\frac{4}{3}, \infty[$  only.

2 The tangent equation of the curve of the function  $f : f(x) = e^{2x+1}$  at the point  $(-\frac{1}{2}, 1)$  is  $\dots\dots\dots$

(a)  $2y = x + 1$

(b)  $y = 2x + 2$

(c)  $y = 2x - 3$

(d)  $2y = 3x + 1$

3 The absolute minimum value of the function  $f$ , where  $f(x) = x + \frac{1}{x}$  in the interval  $[\frac{1}{2}, 3]$  equals  $\dots\dots\dots$

(a) 3

(b) 2

(c) 1

(d)  $2\frac{1}{2}$

4 A regular octagon, its side length is 10 cm., the side length increases at a rate 0.2 cm./sec., then the rate of increasing of its area at this moment =  $\dots\dots\dots$  cm<sup>2</sup>/sec.

(a) 19.3

(b) 4.825

(c) 193

(d) 38.6

5  $\lim_{x \rightarrow 0} \frac{a^x + b^x + c^x - 3}{x} = \dots\dots\dots$

(a)  $\ln(a + b + c)$

(b)  $\ln(a b c)$

(c)  $\ln a - \ln b - \ln c$

(d)  $\log(a b c)$

6 The normal equation to the curve  $y = f(x)$  at the point  $(1, 1)$  is  $x + 4y = 5$ , then  $f'(1) = \dots\dots\dots$

(a) -3

(b)  $-\frac{1}{4}$

(c) 4

(d) -4

7 The radius length of a circle increases at a rate 2 cm./min. and its area at a rate  $20\pi$  cm<sup>2</sup>/min., then the radius length at this instant equals  $\dots\dots\dots$  cm.

(a)  $\frac{5}{2}$

(b) 5

(c) 10

(d) 20



- 8 If  $y = \ln (\sec x + \tan x)$ , then  $\frac{dy}{dx} = \dots\dots\dots$
- (a)  $\tan x$  (b)  $\sec x$  (c)  $\tan^2 x$  (d)  $\csc x$
- 
- 9 If  $f: \mathbb{R} \rightarrow \mathbb{R}$  where  $f(x) = x^3 - kx^2 + 12x + 7$  is one-to-one function, then  $k \in \dots\dots\dots$
- (a)  $\mathbb{R} - \{-6, 6\}$  (b)  $]-\infty, -6]$  (c)  $[-6, 6]$  (d)  $[6, \infty[$
- 
- 10  $\int (1 + 4x^4) e^{x^4} dx = \dots\dots\dots + c$
- (a)  $x + xe^{x^4}$  (b)  $xe^{x^4}$  (c)  $e^{x^4}$  (d)  $\frac{1}{4}e^{x^4}$
- 
- 11 The length of the hypotenuse in a right-angled triangle equals 10 cm., then the length of each side of the right angle when the area is as great as possible equals .....
- (a)  $\sqrt{10}$  cm.,  $\sqrt{10}$  cm. (b)  $5\sqrt{2}$  cm.,  $5\sqrt{2}$  cm.  
 (c)  $10\sqrt{2}$  cm., 5 cm. (d)  $2\sqrt{5}$  cm.,  $2\sqrt{15}$  cm.
- 
- 12 The slope of the tangent at any point  $(x, y)$  on the curve  $y = f(x)$  is  $6x^2 - 30x + 36$ , then given that the curve has a local maximum value equals 28 the equation of the curve is .....
- (a)  $y = 2x^3 - 15x^2$  (b)  $y = 2x^3 - 15x^2 + 36x$   
 (c)  $y = 2x^3 - 15x^2 + 36x + 28$  (d)  $y = 6x^2 - 30x + 8$
- 
- 13 If  $y = 1 + \frac{x}{1} + \frac{x^2}{2} + \frac{x^3}{3} + \dots$  to  $\infty$ , then  $2\ddot{y} + 3\dot{y} - 4y = \dots\dots\dots$
- (a)  $y$  (b) zero (c)  $2y$  (d)  $9y$
- 
- 14 If  $y = e^x \sin x$ , then  $\frac{d^2 y}{dx^2} - 2 \frac{dy}{dx} + 2y = \dots\dots\dots$
- (a) 0 (b)  $e^x$  (c)  $e^x \sin x$  (d)  $e^x \cos x$
- 
- 15  $\int \frac{\sin^6 x}{\cos^8 x} dx = \dots\dots\dots + c$
- (a)  $\tan^7 x$  (b)  $\frac{1}{7} \tan^7 x$  (c)  $\frac{1}{7} \tan^7 x$  (d)  $\sec^7 x$




- 16 The volume of the solid generated by revolving the plane region bounded from the top by the curve  $x^2 + y^2 = 4$  and from the bottom by the two straight lines  $y = x$  ,  $y = -x$  a complete revolution about  $x$ -axis equals ..... cubic unit.

(a)  $\frac{16\sqrt{2}}{3}\pi$       (b)  $\frac{8\sqrt{2}}{3}\pi$       (c)  $\frac{32\sqrt{2}}{3}\pi$       (d)  $\frac{4\sqrt{2}}{3}\pi$

- 17 If the function  $f : f(x) = x^3 - ax^2 + b$  has local minimum value at the point  $(2, 4)$  , then the value of  $a \times b =$  .....

(a) -12      (b) zero      (c) 12      (d) 24

- 18 The opposite figure represents  $f'$  , then the function  $f$  has a local maximum value at  $x =$  .....



(a) 1  
(b) 4  
(c) 0  
(d) 2.5

- 19 A factory is producing electric appliances profits L.E. 30 in every appliance if it produces 50 appliances monthly. When the production increased than that , the profit in the appliance decreases by 50 piasters for every extra appliance produced , then the number of appliances produced monthly to get maximum profit = ..... appliance.

(a) 52      (b) 55      (c) 60      (d) 65

- 20 If  $\int (2x - 1) \ln x \, dx = yz - \int z \, dy$  , then  $yz =$  .....


(a)  $(2x - 1) \ln x$       (b)  $\frac{2x-1}{x}$       (c)  $(x^2 - x) \ln x$       (d)  $x - 1$



21 In the opposite figure :

The area of the shaded region = ..... square units

- (a) 4  
(b) 8  
(c) 12  
(d) 16



22  $\int \frac{1}{\cos X - 1} dX = \dots\dots\dots + c$

- (a)  $\sec X + \tan X$  (b)  $-\csc X - \cot X$  (c)  $\sin X - X$  (d)  $\csc X + \cot X$

23 If  $y = \sin X - \cos X$ , then  $\frac{d^{17}y}{dX^{17}} = \dots\dots\dots$

- (a)  $\sin X + \cos X$  (b)  $\sin X - \cos X$   
(c)  $\cos X - \sin X$  (d)  $-(\sin X + \cos X)$

24 If  $\sin(Xy) + \cos(Xy) = \text{zero}$ , then  $\frac{dy}{dX} = \dots\dots\dots$

- (a)  $Xy$  (b)  $-Xy$  (c)  $\frac{X}{y}$  (d)  $\frac{-y}{X}$

25 The equation of the normal to the curve  $y^2(1 + X^2) = 8$  at the point  $(-1, 2)$  is .....

- (a)  $X + y - 1 = 0$  (b)  $X - y + 3 = 0$  (c)  $y + 2 = -(X - 1)$  (d)  $y + 1 = X - 2$



Answer the following questions :

1 If  $\int_1^4 f(x) dx + \int_b^8 f(x) dx = \int_1^8 f(x) dx$ , then  $b = \dots\dots\dots$

- (a) 2 (b) 4 (c) 1 (d) 8

2  $\lim_{x \rightarrow 0} (1 + 3 \tan^2 x)^{\cot^2 x} = \dots\dots\dots$

- (a)  $e$  (b)  $e^3$  (c)  $3e$  (d)  $e^{\frac{1}{3}}$

3 The slope of the tangent to the curve of the function  $f$  at any point  $(x, y)$  on it is given by the relation  $g(x) = x e^{3x}$ , then the equation of the curve is  $\dots\dots\dots$  given that it passes through the point  $(\frac{1}{3}, 5)$

- (a)  $y = \frac{1}{3} x e^{3x}$  (b)  $y = \frac{1}{9} x e^{3x} + 5 - \frac{e}{9}$   
(c)  $y = \frac{1}{3} x e^{3x} - \frac{1}{9} e^{3x}$  (d)  $y = \frac{1}{3} x e^{3x} - \frac{1}{9} e^{3x} + 5$

4 The ratio between the slope of the tangent to the curve  $y_1 = \ln 3 \sqrt{x+1}$  and the slope of the tangent to the curve  $y_2 = \ln 5 \sqrt{x+1}$  at  $x = a$  is  $\dots\dots\dots$

- (a) 3 : 5 (b) 5 : 3 (c) 1 : 1 (d)  $\ln 3 : \ln 5$

5 The length of each of two equal sides in an isosceles triangle equals 6 cm. and the measure of the angle between them equals  $(x)$ , the rate of change of  $x$  is  $(\frac{\pi}{90})^{\text{rad.}}$  per minute, then the rate of change of its area at  $x = 30^\circ$  is  $\dots\dots\dots$

- (a)  $\frac{\sqrt{3}}{10} \pi$  (b)  $\frac{\pi}{10}$  (c)  $9\sqrt{3}$  (d) 9

6  $\int (\sin^2 x + \cos^2 x + \cot^2 x) dx = \dots\dots\dots + c$

- (a)  $\csc^2 x$  (b)  $\cot x$  (c)  $-\cot x$  (d)  $x + \frac{1}{3} \cot^3 x$

7 If the function  $f : f(x) = x^3 - 3x + 4$ , then the function is decreasing on the interval  $\dots\dots\dots$


- (a)  $]-\infty, 0[$  (b)  $]0, \infty[$   
(c)  $]-\infty, -1[ \cup ]1, \infty[$  (d)  $]-1, 1[$



8 The normal equation to the curve  $y = x|x|$  at the point  $(-2, -4)$  is .....

- (a)  $y + 4x + 12 = 0$  (b)  $4y + x + 18 = 0$   
 (c)  $4y + x + 14 = 0$  (d)  $y + 4x - 4 = 0$

9 The top right corner of a piece of paper whose dimensions are 8 cm. , 12 cm. is folded to the lower edge as shown in the figure , then the value of  $x$  which makes  $y$  as small as possible = .....




- (a) 6 (b) 4 (c) 2 (d) 8

10  $\pi \int_{-2}^2 (4 - x^2) dx$  , is the volume of .....

- (a) a sphere whose radius is 4 units.  
 (b) a right circular cone whose height is 4 units.  
 (c) a sphere whose radius is 2 units.  
 (d) a right circular cylinder whose height is 4 units.

11 The opposite figure represents the curve  $f'(x)$  of the function  $f$  , then the solution set of the inequality  $f'(x) > 0$  is .....



- (a)  $]-1, \infty[$   
 (b)  $]1, \infty[$   
 (c)  $]-\infty, -1[$   
 (d)  $]-\infty, 1[$

12 If  $y = \sin(5x^3) \csc(5x^3)$  , then  $\frac{dy}{dx} = \dots\dots\dots$

- (a) 0  
 (b)  $-25 \cos(5x^3) \times \csc(5x^3) \cot(5x^3)$   
 (c)  $15x^2 \cos(5x^3) - 15x^2 \csc(5x^3) \cot(5x^3)$   
 (d) 1

13 The area of the region bounded by the curve :  $y = x^2 - 9$  , the  $x$ -axis , the straight line  $x = 4$  and above  $x$ -axis = ..... square unit.

- (a)  $\frac{44}{3}$  (b) 18 (c)  $\frac{10}{3}$  (d)  $\frac{98}{3}$



- 14 The gas leaks from a spherical balloon at a rate  $20 \text{ cm}^3/\text{sec}$ . , then the rate of change of the balloon external surface area at the moment in which the radius length is 10 cm. equals .....  $\text{cm}^2/\text{sec}$ .

(a) -4                      (b)  $80\pi$                       (c) -2                      (d) -1

- 15 If  $\sqrt{x} + \sqrt{y} = 1$  , then  $\frac{dy}{dx}$  at the point  $(\frac{1}{4}, \frac{1}{4})$  equals .....

(a)  $\frac{1}{2}$                       (b) 1                      (c) -1                      (d) 2

- 16 If  $f$  is a fifth degree polynomial , then the fifth derivative of the function  $f$  equals .....

(a)  $x$                       (b)  $5x$   
(c) zero                      (d) non zero constant.

- 17  $\int \frac{\ln x^5}{x \ln x^3} dx = \dots + c$

(a)  $\frac{5}{3} \ln |x|$                       (b)  $\frac{3}{5} \ln |x|$                       (c)  $\ln (\ln |x|)$                       (d)  $\ln (\frac{5}{3} |x|)$

- 18 If the two curves of the functions  $f$  and  $g$  are touching at the point  $(2, 4)$  and  $\hat{f}(2) = 3$  , then  $\hat{g}(2) = \dots$

(a) 2                      (b) 3                      (c) 4                      (d) 5

- 19 If  $a^y = b^x$  where  $a, b \in \mathbb{R}^+ - \{1\}$  , then  $\frac{dy}{dx} = \dots$

(a)  $\log \frac{a}{b}$                       (b)  $\log_a b$                       (c)  $\log_b a$                       (d)  $\log \frac{b}{a}$

- 20 If the function  $f : f(x) = kx^2 + (k+5)x + k - 2$  has local maximum value at  $x = 2$  , then  $k = \dots$

(a) -2                      (b) -1                      (c) zero                      (d) 1

- 21 If  $a \neq b$  and  $\int_a^b (3x^2 - 1) dx = \text{zero}$  , then  $a^2 + b^2 = \dots$

(a)  $ab$                       (b)  $1 - ab$                       (c)  $ab + 1$                       (d)  $a + b$

- 22 If the perimeter of a circular sector is constant  $p$  , then the area has maximum value at  $r = \dots$

(a)  $\frac{p}{2}$                       (b)  $\frac{2}{\sqrt{p}}$                       (c)  $\sqrt{p}$                       (d)  $\frac{p}{4}$



23 The absolute maximum value of the function  $f$  where  $f(x) = 10xe^{-x}$ ,  $x \in [0, 4]$  is .....

- (a)  $\frac{10}{e}$       (b) zero      (c) 1      (d)  $e$


24 If  $f(x) = \frac{1 - \cot x}{1 + \cot x}$ , then  $f\left(\frac{\pi}{4}\right) = \dots\dots\dots$

- (a) zero      (b) 1      (c)  $\sqrt{2}$       (d)  $\frac{1}{\sqrt{2}}$

25 In the opposite figure :

The volume of the solid generated by revolving the shaded area a complete revolution about the y-axis = ..... cube units.

- (a)  $\frac{4}{3} \pi$   
 (b)  $\frac{8}{3} \pi$   
 (c)  $6 \pi$   
 (d)  $\frac{16}{3} \pi$





Answer the following questions :

1  $\lim_{x \rightarrow 0} \frac{(10)^{\sin x} - 1}{\tan x} = \dots\dots\dots$

- (a) 10                      (b)  $\ln 10$                       (c)  $\ln \sin x$                       (d) 1

2  $\int_0^6 |3 - x| dx = \dots\dots\dots$

- (a) 9                      (b) -9                      (c)  $\frac{9}{2}$                       (d)  $-\frac{9}{2}$

3 The absolute maximum value of the function  $f : f(x) = \frac{x}{x^2 + 1}$ ,  $x \in [0, 2]$  equals .....

- (a) 0                      (b)  $\frac{1}{2}$                       (c)  $\frac{2}{5}$                       (d) 1


4 The height of a cylinder which has the greatest volume placed inside a sphere whose radius length (r) equals ..... length unit.

- (a)  $\frac{2r}{\sqrt{5}}$                       (b)  $\frac{2r}{\sqrt{3}}$                       (c)  $2r$                       (d)  $2\sqrt{3}r$


5 If  $y = \ln \sqrt{\tan x}$ , then  $\frac{dy}{dx} = \dots\dots\dots$  when  $x = \frac{\pi}{4}$

- (a) 1                      (b) zero                      (c)  $\frac{1}{2}$                       (d)  $\infty$


6 Which of the following figures represents the curve of the continuous function  $f$  in which  $f(0) = 3$ ,  $f(2) = f(-2) = 0$ ,  $f'(x) > 0$  when  $-2 < x < 2$ ?




(a)



(b)



(c)



(d)

7  $\int \frac{\ln x}{x} dx = \dots\dots\dots + c$

- (a)  $(\ln x)^2$                       (b)  $\frac{1}{2}(\ln x)^2$                       (c)  $\frac{1}{x^2}$                       (d)  $\ln x - 1$



8 If  $x = a \sec^2 \theta$ ,  $y = a \tan^3 \theta$ , then  $\frac{d^2 y}{d x^2} = \dots\dots\dots$

- (a)  $\frac{3}{2} \sec^2 \theta$  (b)  $\frac{3 \cot \theta}{4 a}$   
 (c)  $\frac{3 a \cot \theta}{4}$  (d)  $3 a \sec^4 \theta \tan \theta$

9 The current intensity  $I$  (Ampere) in a circuit for alternating current at any moment  $t$  (second) is given by the relation  $I = 2 \cos t + 2 \sin t$ , then the maximum value of the current in this circuit equals  $\dots\dots\dots$

- (a)  $2\sqrt{2}$  (b)  $\sqrt{2}$  (c)  $\frac{\pi}{4}$  (d) 8

10 A ladder of constant length its upper end slides on a vertical wall at a rate  $k$  length unit/sec., then the rate of increasing of the distance between the lower end and the wall when the ladder inclined to the vertical with an angle  $\theta$  where  $\csc \theta = \frac{5}{4}$  equals  $\dots\dots\dots$  unit/sec.


- (a)  $\frac{3}{4}$  (b)  $\frac{12}{25}$  (c)  $\frac{12}{25} k^2$  (d)  $\frac{3 k}{4}$

11 In the opposite figure :

If  $A_1 = 5$  square unit,  $A_2 = 2$  square unit,  $A_3 = 8$  square unit

, then  $-\int_{-2}^5 f(x) dx + \int_{-2}^5 |f(x)| dx = \dots\dots\dots$

- (a) 15 (b) 20  
 (c) 22 (d) 26



12 If  $f(x) = x - x \ln x$ , then the slope of the tangent to the curve at  $x = e$  equals  $\dots\dots\dots$

- (a) zero (b) -1 (c) 1 (d) e

13 The volume of the solid generated by revolving the plane region bounded by the curve  $y = 2\sqrt{x-1}$  (where  $x \geq 1$ ) and the tangent at the point  $(2, 2)$  and the straight line  $y = 0$  a complete revolution about  $x$ -axis equals  $\dots\dots\dots$  cubic unit.

- (a)  $\frac{\pi}{3}$  (b)  $\frac{16\pi}{3}$  (c)  $\frac{2\pi}{3}$  (d)  $\frac{\pi}{2}$

14 If the normal to the curve  $y = x \ln x$  is parallel to the straight line  $2x - 2y + 3 = 0$ , then the normal equation is  $\dots\dots\dots$

- (a)  $x - y = 3e^{-2}$  (b)  $x - y = 6e^{-2}$   
 (c)  $x - y = 3e^2$  (d)  $x - y = 6e^2$



15  $\int \frac{\sec^2 X}{\tan X} dX = \dots\dots\dots + c$

- (a)  $-\frac{1}{2}\tan^{-2} X$       (b)  $\ln |\tan X|$       (c)  $\ln |\sec^2 X|$       (d)  $\frac{1}{3} \sec^3 X$

16 The derivative of  $e^{\sin X}$  with respect to  $\sin X$  equals .....

- (a)  $e^{\sin X}$       (b)  $e^{\frac{1}{\sin X}}$       (c)  $\cos X$       (d)  $\sin X$

17  $n$  is the number of sides of a regular polygon whose side length increases at a constant rate (a) cm./sec. , then the measure of its vertex angle .....

- (a) increases at a constant rate (a)<sup>rad.</sup>/sec.  
 (b) increases at a constant rate (na)<sup>rad.</sup>/sec.  
 (c) increases at a non constant rate and unknown.  
 (d) remains constant.

18 The two curves  $y = X^2 + aX + b$  ,  $y = cX - X^2$  are touching at the point (1, 0) , then  $b + c - a = \dots\dots\dots$

- (a) 0      (b) 2      (c) 3      (d) 6

19 If  $\hat{f}(X) = X f(X)$  ,  $f(3) = -5$  , then  $\hat{\hat{f}}(3) = \dots\dots\dots$

- (a) -50      (b) -40      (c) 15      (d) 27

20 If  $f(X) = \cot\left(\frac{\pi}{3} \sin X\right)$  , then  $\hat{f}\left(\frac{\pi}{6}\right) = \dots\dots\dots$

- (a)  $-\frac{2\sqrt{3}\pi}{3}$       (b)  $-\frac{\sqrt{3}\pi}{2}$       (c)  $\pi$       (d)  $\frac{\sqrt{3}\pi}{6}$


21 If the curve  $y = X^3 + aX^2 + bX$  has an inflection point at (3, -9) , then  $a + b = \dots\dots\dots$

- (a) 15      (b) 6      (c) -9      (d) -12

22 In the opposite figure :

$\int_0^1 \hat{f}(X) \cdot f(X) \cdot dX = \dots\dots\dots$


- (a)  $\frac{7}{2}$       (b)  $\frac{5}{2}$   
 (c)  $-\frac{5}{2}$       (d)  $-\frac{7}{2}$






- 23 The opposite figure represents the curve of first derivative of the function  $y = f(x)$ , then all the following statements are true except .....

- (a) at  $x = -1$ , there is an inflection point of the function  $y = f(x)$   
 (b) at  $x = 1$ , there is a local minimum value of the function  $y = f(x)$   
 (c) at  $x = -2$ , there is a local minimum value of the function  $y = f(x)$   
 (d) at  $x = 3$ , there is a local maximum value of the function  $y = f(x)$




- 24 The opposite figure represents the curve  $\hat{f}$ , then the function  $f$  is increasing on the interval .....

- (a)  $]-\infty, 2[$                       (b)  $]-\infty, 1[$   
 (c)  $]1, 2[$                           (d)  $]1, 3[$



- 25 The opposite figure represents a quadratic function, its vertex is  $(k, 9)$ , then the area of the shaded region = ..... square units.

- (a) 6  
 (b) 9  
 (c) 12  
 (d) 18





Answer the following questions :

1  $\lim_{x \rightarrow \infty} \left( \frac{x+5}{x+3} \right)^x = \dots\dots\dots$

(a)  $e$

(b)  $e^2$

(c)  $\frac{1}{e}$

(d)  $\frac{2}{e}$

2 The rate of change of tangent slope of a curve at any point  $(X, y)$  on it is  $6(1 - 2X)$  and the curve has a critical point at  $X = 1$  and the function has a local minimum value equals 4, then the normal equation to the curve at  $X = -1$  is .....

(a)  $12X + y + 3 = 0$

(b)  $X - 12y + 109 = 0$

(c)  $12X + y - 3 = 0$

(d)  $y - 12X - 3 = 0$

3 The measure of the angle which the tangent to the curve  $\sin 2X = \tan y$  makes with the positive direction of  $X$ -axis at the point  $\left( \frac{3\pi}{4}, \frac{3\pi}{4} \right)$  equals .....

(a) zero

(b)  $135^\circ$

(c)  $45^\circ$

(d)  $26^\circ 34'$

4 If  $y = X^2 + 3X + 2$ ,  $z = 3X^2 - 5X + 4$ , then  $\frac{d^2 y}{dz^2}$  at  $X = 2$  equals .....

(a)  $-\frac{4}{7}$

(b)  $-\frac{4}{49}$

(c) 8

(d) 56

5 In the opposite figure :


$\int_0^2 [f(x)]^2 f'(x) dx = \dots\dots\dots$

(a) -9

(b) 9

(c) 2

(d) 1



6 A rectangle whose perimeter is 40 m., its area is as great as possible when its dimensions are ..... m.

(a) 15, 5

(b) 10, 10

(c) 12.5, 7.5

(d) 9, 11

7 If  $g(9) = 7$ ,  $g(4) = 3$ , then  $\int_2^3 2X g(X^2) g'(X^2) dX = \dots\dots\dots$

(a) 10

(b) 20

(c) 5

(d) 2



- 8 ABC is right-angled triangle at  $\angle C$ , its area is constant and equals  $24 \text{ cm}^2$  the rate of change of (b) equals  $1 \text{ cm./sec.}$ , then the rate of change of (a) at the instant when b equals  $8 \text{ cm.}$  equals .....  $\text{cm./sec.}$

(a)  $\frac{-3}{4}$  (b)  $\frac{-3}{2}$  (c)  $\frac{3}{8}$  (d)  $\frac{3}{4}$

- 9 If  $f: \mathbb{R}^+ \rightarrow \mathbb{R}^+$  where  $f(x) = x^{2x}$ , then  $\dot{f}(e) = \dots\dots\dots$

(a)  $4e^{2e}$  (b)  $4$  (c)  $2e^{2e}$  (d)  $4e$

- 10 If  $f(x) = \begin{cases} 2x^3 + 3 & , x \leq -1 \\ 3x + 4 & , x > -1 \end{cases}$ , then  $_{-2} \int^2 f(x) dx = \dots\dots\dots$

(a)  $6$  (b)  $9$  (c)  $12$  (d)  $15$

- 11 The area of the region bounded by the curve  $y = x^2 - 9$  and  $x$ -axis and the straight line  $x = 4$  and above  $x$ -axis equals ..... square unit.

(a)  $\frac{10}{3}$  (b)  $36$  (c)  $\frac{98}{3}$  (d)  $18$

- 12 If  $f$  is a differentiable odd function in the interval  $]-\infty, \infty[$  and  $\dot{f}(3) = 2$ , then  $\dot{f}(-3) = \dots\dots\dots$

(a) zero (b)  $1$  (c)  $2$  (d)  $4$

- 13  $\int \frac{\cos^2 x}{1 - \sin x} dx = \dots\dots\dots + c$

(a)  $-\cos x$  (b)  $1 - \cos x$  (c)  $x + \cos x$  (d)  $x - \cos x$


- 14 If  $x \in [0, \pi]$ , then the function  $f: f(x) = x \sin x + \cos x$  has an absolute minimum value at  $x = \dots\dots\dots$

(a) zero (b)  $\frac{\pi}{2}$  (c)  $\pi$  (d)  $-1$

- 15 In the opposite figure :

The volume of a solid generated by revolving the shaded region a complete revolution about  $x$ -axis and the straight line  $x = -1$ ,  $x = k$  equals  $\frac{\pi}{2} (e^{10} - e^{-2})$  cube unit, then  $k = \dots\dots\dots$

(a)  $5$  (b)  $10$   
(c)  $2$  (d)  $1$





16 The slope of the tangent to the curve  $y = \sqrt{x + \sec x}$  at  $x = 0$  equals .....

- (a) -1                      (b) zero                      (c)  $\frac{1}{2}$                       (d) 1



17 If  $f(x) = 20x^{n-1}$  and  $f'(x) = c$ ,  $c \in \mathbb{R}$ ,  $n \in \mathbb{Z}^+$ , then  $n + c = \dots\dots\dots$

- (a) 104                      (b) 124                      (c) 123                      (d) 125

18 If  $f: \mathbb{R} \rightarrow \mathbb{R}$  where  $f(x) = \pi^x e^x$ , then  $f'(x) = \dots\dots\dots$

- (a)  $f(x) \ln \pi$                       (b)  $f(x) e^x$   
(c)  $f(x) \ln(\pi e)$                       (d)  $f(x)$

19 Water poured into a right circular cylinder at a constant rate as shown in the opposite figure which of the following figures represents the relation between the water height ( $h$ ) in the cylinder and the time ( $t$ )?



20 If the equation of the normal to the common tangent of the two functions  $f$  and  $g$  at  $x = 1$  is  $y = -\frac{1}{3}x + \frac{3}{2}$ , then  $(f \times g)'(1) = \dots\dots\dots$

- (a) 4                      (b) 7                      (c) 2                      (d) 10

21 If  $f$  is an even function and continuous on  $\mathbb{R}$  and the function has an inflection point at  $x = -a$ , then the sign of  $f'(a^-) \times f'(a^+)$  is the same as .....

- (a)  $f'(a^-) \times f'(a^+)$                       (b)  $f'(a^-) \times f'(a^+)$   
(c)  $f'(-a)$                       (d)  $f'(-a^-) \times f'(a^-)$




22  $\int \frac{\ln X^4}{\ln X} dX = \dots\dots\dots$

- (a)  $4X + c$                       (b)  $\frac{X}{4} + c$                       (c)  $\frac{4}{X} + c$                       (d)  $4X^2 + c$


23 The opposite figure represents the curve of the function  $f$ , then  $\hat{f}$  is negative in the interval .....

- (a)  $]1, 2[$                       (b)  $]0, 3[$   
(c)  $\mathbb{R} - [1, 2]$                       (d)  $]0, 2[$




24 The opposite figure represents the curve of the function  $f$  where  $f(X) = 3X - X^2$ . If the point  $C(a, b)$  lies on the curve, then the greatest value of the expression  $a + b$  is .....

- (a) 6                      (b) 5                      (c) 4                      (d) 3



25 The opposite figure represents the curve of the first derivative of function  $y = f(X)$ , then all the following statements are true except .....

- (a)  $f$  increases on  $]-\infty, 3[$   
(b)  $f$  decreases on  $]3, \infty[$   
(c)  $f(-2) > f(-3)$   
(d)  $f$  decreases on  $]-\infty, -2[$





Answer the following questions :

1 If  $\int_3^5 f(x) dx = 6$ , then  $\int_3^5 [4f(x) - 1] dx = \dots\dots\dots$

- (a) 18 (b) 22 (c) 23 (d) 26

2 If  $f(x) = e^{\tan x}$ , then  $\lim_{x \rightarrow \frac{\pi}{4}} \frac{f(x) - f(\frac{\pi}{4})}{x - \frac{\pi}{4}} = \dots\dots\dots$

- (a)  $e$  (b)  $2e$  (c)  $e^2$  (d)  $2e^2$

3 The minimum value of the function  $f : f(x) = x \ln x$  equals  $\dots\dots\dots$

- (a)  $e$  (b)  $\frac{1}{e}$  (c)  $-\frac{1}{e}$  (d)  $-e$


4 The local maximum value of the function :  $y = \frac{1}{3}x^3 - 9x + 2$  equals  $\dots\dots\dots$

- (a) 20 (b) -16 (c) 3 (d) -3

5 In the opposite figure :

The area of the shaded part =  $\dots\dots\dots$  square unit.

- (a) 2  
(b)  $2\sqrt{2}$   
(c)  $2\sqrt{2} + 2$   
(d)  $2\sqrt{2} - 2$



6 If  $f(x) = x^{2019}$ , then the 2019<sup>th</sup> derivative of this function equals  $\dots\dots\dots$

- (a) 2019 (b) 2018 (c) 2019 (d) zero

7  $\int e^x (\cot x - \csc^2 x) dx = \dots\dots\dots + c$

- (a)  $e^x \cot x - 2e^x \csc^2 x$  (b)  $e^x (-\csc x - \cot^2 x)$   
(c)  $e^x \cot x$  (d)  $e^x \tan x - e^x \cot x$



8 A 5-metre rod is fixed by a hinge to the ground at its base. If its top rises up by a winch at a rate of 1 m./min. , then the rate of decreasing the projection length of the rod on the ground when the height of the top is 3 metres = ..... m./min.

- (a)  $\frac{3}{4}$  (b)  $\frac{3}{2}$  (c)  $\frac{2}{3}$  (d)  $\frac{4}{3}$

9 If  $f(x) = (\cos x)^{\cos x}$ , then  $f'(0) = \dots\dots\dots$

- (a) -3 (b) -2 (c) -1 (d) zero

10  $\int_{-2}^4 |x^2 - 3x| dx = \dots\dots\dots$

- (a) 9 (b) 15 (c) 18 (d) 24

11 The function  $f : f(x) = x^3 + 4x + 8$  increases at  $x \in \dots\dots\dots$

- (a)  $]-4, \infty[$  only. (b)  $]-\infty, -\frac{4}{3}[$  only. (c)  $]-\frac{4}{3}, \infty[$  only. (d)  $\mathbb{R}$

12 If  $y = \sec^n(x)$ , then  $\frac{dy}{dx} = \dots\dots\dots$

- (a)  $n y \sec x$  (b)  $n y \tan x$   
(c)  $n y \sec^2 x$  (d)  $n y \tan^2 x$

13 A rectangle is drawn such that two adjacent vertices of the rectangle lie on the curve  $y = x^2 - 12$  and the other two vertices lie on the curve  $y = 12 - x^2$ , then the maximum area of this rectangle = ..... square unit.

- (a) 96 (b) 64 (c) 8 (d) 112

14 The curve of the function  $f$  where  $f(x) = \sqrt[3]{x-3}$  is convex upward in the interval .....

- (a)  $]3, \infty[$  (b)  $]-\infty, 3[$  (c)  $]-\infty, 0[$  (d)  $]0, \infty[$

15 The volume of the solid generated by revolving the region bounded by the curve  $y = \sqrt{x+5}$  and the straight lines  $y=0$ ,  $x=1$ ,  $x=3$  a complete revolution about  $x$ -axis = ..... cubic unit.

- (a)  $14\pi$  (b)  $5.3\pi$  (c)  $19.5\pi$  (d)  $32\pi$



- 16 ABC is an equilateral triangle of side length  $2l$ , E is the midpoint of  $\overline{BC}$ ,  $D \in \overline{AB}$ ,  $F \in \overline{AC}$  such that  $\overline{DF} \parallel \overline{BC}$ , then greatest area of the triangle DEF = ..... the area  $\Delta ABC$

(a)  $\frac{1}{4}$  (b)  $\frac{1}{2}$  (c) 2 (d) 4

- 17 A train starts its journey at 11 O'clock towards east with velocity 45 km./h., while another train began its journey at 12 O'clock from the same point towards south with velocity 60 km./h., then the rate of increasing of distance between the two trains at 3 O'clock afternoon = ..... km./h.


(a) 180 (b)  $52.5\sqrt{2}$  (c)  $180\sqrt{2}$  (d) 75

- 18  $\lim_{x \rightarrow 0} \left(1 + \frac{x}{a}\right)^{\frac{a}{x}} = \dots\dots\dots$

(a) zero (b)  $\frac{1}{a}$  (c)  $\frac{1}{e}$  (d) e

- 19 The given figure represents the curve  $f''(x)$ , then the curve of  $f(x)$  is convex upwards at  $x \in \dots\dots\dots$

(a)  $]-\infty, 0[$   
 (b)  $]-\infty, 3[$   
 (c)  $]0, \infty[$   
 (d)  $]3, \infty[$



- 20  $-\pi \int_{-\pi}^{\pi} (4 + \pi \cos 2x) dx = \dots\dots\dots$

(a)  $\pi$  (b)  $2\pi$  (c)  $4\pi$  (d)  $8\pi$

- 21 The measure of the positive angle which the tangent to the curve :  $y^2 + 2x^2 = 6$  at the point (1, 2) makes with the positive direction of x-axis = .....°

(a) 45 (b) 135 (c) 120 (d) 150

- 22 If  $y = e^{ax}$ , then  $\frac{d^4 y}{dx^4} = \dots\dots\dots$

(a) ay (b)  $a^4 x$  (c)  $a^4 y$  (d)  $-a^4 y$



23 The slope of the tangent to a curve at any point  $(x, y)$  on it equals  $\frac{\sqrt{2y+1}}{\sqrt{3x-2}}$ , then the equation of the curve given that it passes through  $(1, 4)$  is .....

(a)  $\sqrt{2y+1} = \sqrt{3x-2}$

(b)  $2y+1 = 3x$

(c)  $\sqrt{2y+1} = \frac{2}{3}\sqrt{3x-2} + \frac{7}{3}$

(d)  $\sqrt{2y+1} = 2\sqrt{3x-2} + 1$

24  $\frac{d}{dx}[(\csc x - \cot x)(\csc x + \cot x)] = \dots\dots\dots$

(a) zero

(b)  $\csc^2 x - \cot^2 x$

(c)  $\csc x \cot x + \sec^2 x \tan x$

(d)  $\csc x \cot x - \csc^2 x$

25  $\int \frac{dx}{\sqrt[3]{2x+9}} = \dots\dots\dots + c$

(a)  $\frac{3}{2}(2x+9)^{\frac{2}{3}}$

(b)  $\frac{3}{4}(2x+9)^{\frac{2}{3}}$

(c)  $\frac{-3}{8}(2x+9)^{-\frac{4}{3}}$

(d)  $\frac{1}{3}(2x+9)^{\frac{2}{3}}$



Answer the following questions :

- 1 If  $y = \frac{z+1}{z-1}$ ,  $x = \frac{z-1}{z+1}$ , then  $\frac{dy}{dx} = \dots\dots\dots$  at  $x = 2$   
 (a)  $-9$  (b)  $4$  (c)  $\frac{1}{4}$  (d)  $-\frac{1}{4}$
- 
- 2 The tangent to the curve  $x = e^\theta \cos \theta$ ,  $y = e^\theta \sin \theta$  at the point at which  $\theta = \frac{\pi}{4}$  makes with the positive direction of  $x$ -axis an angle of measure  $\dots\dots\dots$   
 (a) zero (b)  $\frac{\pi}{4}$  (c)  $\frac{\pi}{3}$  (d)  $\frac{\pi}{2}$
- 
- 3  $\int_0^2 \sqrt{4-x^2} dx = \dots\dots\dots$   
 (a) zero (b)  $2$  (c)  $\pi$  (d)  $\frac{\pi}{2}$
- 
- 4 If  $f(2x-1) = 2x^3 - 4x^2 + 6x + 7$ , then  $f(3) = \dots\dots\dots$   
 (a)  $7$  (b)  $33$  (c)  $36$  (d)  $14$
- 
- 5  $\int \frac{\sec^2 x}{\tan x} dx = \dots\dots\dots + c$   
 (a)  $-\frac{1}{2} \tan^{-2} x$  (b)  $\ln |\tan x|$  (c)  $\ln |\sec^2 x|$  (d)  $\frac{1}{3} \sec^3 x$
- 
- 6  $\lim_{x \rightarrow \infty} \left( \frac{x+4}{x-2} \right)^{x+3} = \dots\dots\dots$   
 (a)  $e^2$  (b)  $e^{-6}$  (c)  $-e^2$  (d)  $e^6$
- 
- 7  $\int \frac{x - \frac{1}{2}}{\sqrt{2x-1}} dx = \dots\dots\dots + c$   
 (a)  $\frac{1}{2} (2x-1)^{\frac{1}{2}}$  (b)  $\frac{1}{3} (2x-1)^{\frac{3}{2}}$   
 (c)  $(2x-1)^{\frac{3}{2}}$  (d)  $\frac{1}{6} (2x-1)^{\frac{3}{2}}$
- 
- 8 If  $f$  is a function where  $f(x) = x|x-2|$ , then the function  $f$  is increasing on the interval  $\dots\dots\dots$   
 (a)  $]1, 2[$  only. (b)  $] -\infty, 1[$ ,  $]2, \infty[$  only.  
 (c)  $] -\infty, 1[$  only. (d)  $]1, \infty[$  only.



- 9 The greatest value of the expression  $(\sin x + \sqrt{3} \cos x)$  is at  $x = \dots\dots\dots$  where  $x \in [0, \frac{\pi}{2}]$
- (a)  $\frac{\pi}{3}$  (b)  $\frac{\pi}{4}$  (c)  $\frac{\pi}{6}$  (d) zero

- 10 In a closed electric circuit,  $V$  is the potential difference (Volt),  $I$  is the current intensity (Ampere)  $R$  is the resistance (Ohm). If the potential difference increases at a rate of 1 Volt/sec. and the current intensity decreases at a rate of  $\frac{1}{2}$  Ampere/sec., then the rate of the resistance at the moment which  $V = 12$  Volt and  $I = 2$  Amperes equals  $\dots\dots\dots$  ohm./sec. (where  $V = IR$ )

(a) 2 (b) 4 (c) 6 (d) 1

- 11 If  $y = 2 \sin x - x \cos x$ , then:  $\frac{d^2 y}{d x^2} + y = \dots\dots\dots$

(a)  $x \cos x$  (b)  $\sin x$  (c)  $2 \sin x$  (d)  $2 \cos x$

- 12 If  $\int (2x + 3) \ln x \, dx = yz - \int z \, dy$ , then  $yz = \dots\dots\dots$

(a)  $2x \ln x$  (b)  $(2x + 3) \ln x$   
(c)  $\frac{1}{2} (2x + 3) \ln x$  (d)  $x(x + 3) \ln x$

- 13 If the area of the region bounded by the two curves  $y = 2x^2$ ,  $y^2 = 4a$  and  $x$  equals  $\frac{2}{3}$  square unit, then  $a = \dots\dots\dots$  where  $a > 0$

(a)  $\frac{2}{3}$  (b) 1 (c)  $\frac{4}{9}$  (d)  $\frac{9}{4}$

- 14 If  $f(x) = (a - 2)x^2 + 3x - 5$ ,  $x \in \mathbb{R}$ , then the curve of the function  $f$  is concave downwards when  $\dots\dots\dots$

(a)  $a > 2$  (b)  $a < 2$  (c)  $a = 2$  (d)  $a = 0$

- 15 If  $f(x) = 2 \sin \frac{x}{2} \cos \frac{x}{2}$ , then the thousandth derivative of this function equals  $\dots\dots\dots$

(a)  $\sin x$  (b)  $\cos x$  (c)  $-\sin x$  (d)  $-\cos x$

- 16 The volume of the solid generated by revolving the region bounded by the curve  $y = \frac{4}{x}$  and the straight line  $y + x = 5$  a complete revolution about  $x$ -axis =  $\dots\dots\dots$  cubic unit.

(a)  $3\pi$  (b)  $6\pi$  (c)  $21\pi$  (d)  $9\pi$




17 In the opposite figure :

$CD = 2AF$  ,  $FE = ED$  , the perimeter of the figure  $ABCDEF = 40$  cm.

, then the greatest area of the figure  $ABCDEF$  equals .....  $\text{cm}^2$

- (a) 90 (b) 95  
(c) 89 (d) 91



18 If  $\int_0^{\pi} \frac{\cos x}{1+x^8} dx = k$  , then  $-\pi \int_{-\pi}^{\pi} \frac{2 \cos x}{1+x^8} dx = \dots\dots\dots$

- (a)  $2k$  (b)  $4k$  (c)  $6k$  (d) zero


19 The opposite figure represents the

curve  $f'(x)$  , then the curve has

maximum value at  $x = \dots\dots\dots$

and minimum value at  $x = \dots\dots\dots$

- (a)  $-3, 6$  (b)  $6, -3$   
(c)  $6, 0$  (d)  $0, 3$



20 The sides of right-angled triangle changes , and its perimeter remains constant 40 cm. the rate of change of the hypotenuse is 7 cm./min. when the side lengths 8 , 15 , 17 cm. , then the rate of change of each side of the right-angle at this moment equals ..... cm./min.

- (a)  $-32, 25$  (b)  $-2, 9$  (c)  $2, -9$  (d)  $32, -25$

21 If  $y = \ln x$  , then  $\frac{d^{10} y}{d x^{10}} = \dots\dots\dots$

- (a)  $\frac{9}{-x^{10}}$  (b)  $\frac{10}{-x^9}$  (c)  $\frac{9}{x^{10}}$  (d)  $\frac{10}{x^9}$

22 If the curve of the function  $f$  lies above all tangents drawn from all points on the curve then the curve of the function is .....

- (a) convex upwards. (b) increasing.  
(c) convex downwards. (d) decreasing.



23 If the tangent to the curve  $y = x^2$  passes through the point  $(3, 5)$ , then the equation of this tangent is .....

(a)  $y = 6x - 13$

(b)  $y = 2x - 1$  or  $y = 10x - 25$

(c)  $y = -10x + 25$  or  $y = -2x + 1$

(d)  $y = x + 8$

24  $\int x^5 \left(1 + \frac{3}{x}\right)^5 dx = \dots + c$

(a)  $\frac{1}{6} (x+3)^6$

(b)  $\frac{1}{11} (x+3)^{11}$

(c)  $\frac{1}{18} (x+3)^6$

(d)  $\frac{1}{18} x^6$

25 If  $\frac{dz}{dx} = 2x - 3$ ,  $\frac{dy}{dx} = x^2 + 1$ , then  $\frac{d^2z}{dy^2}$  at  $x = 1$  equals .....

(a) 1

(b)  $\frac{3}{4}$

(c)  $\frac{3}{2}$

(d)  $\frac{4}{3}$




Answer the following questions :

- 1 The slope of the tangent to the curve  $x y^3 = 3$  at the point  $(3, 1)$  equals .....

(a)  $\frac{1}{9}$  (b)  $-\frac{1}{9}$  (c)  $\frac{2}{3}$  (d)  $-\frac{2}{3}$

- 2 The opposite figure represents the first derivative of the function  $f$  and the function  $f$  has a local minimum equals  $-3$ , then :  $\int_0^3 f(x) dx = \dots\dots\dots$

(a)  $\frac{27}{4}$  (b)  $-27$   
(c)  $\frac{9}{4}$  (d)  $-\frac{27}{4}$



- 3 If the curve of a function which passes through the point  $(\frac{\pi}{2}, \frac{\pi^2}{4} + 9)$  given that the slope of its tangent at any point on it  $(x, y)$  is given by the relation  $m = 2x + \frac{1}{2} \sec^2 \frac{x}{2}$ , then the equation of this curve is .....

(a)  $y = x^2 + \tan \frac{x}{2}$  (b)  $y = x^2 + \tan x + 8$   
(c)  $y = x^2 + 2 \tan \frac{x}{2} + 4$  (d)  $y = x^2 + \tan \frac{x}{2} + 8$

- 4  $\int \frac{1 + \sin^2 x}{1 - \sin^2 x} dx = \dots\dots\dots + c$

(a)  $2 \tan x - x$  (b)  $2 \sec^2 - 1$   
(c)  $x$  (d)  $\frac{1}{3} \sec^3 x - x$

- 5  $\lim_{x \rightarrow 0} \frac{e^x - e^{\sin x}}{x - \sin x} = \dots\dots\dots$

(a) 1 (b) e (c) zero (d)  $-e$

- 6  $-\int_a^a \frac{x}{x^4 + \cos x} dx = \dots\dots\dots$

(a)  $-a$  (b)  $2a$  (c) zero (d)  $\frac{a^2}{a^5 + \sin a}$

- 7 The normal equation to the curve  $2y = 3 - x^2$  at the point  $(1, 1)$  is .....

(a)  $x + y = 0$  (b)  $x + y + 1 = 0$   
(c)  $x - y + 1 = 0$  (d)  $x - y = 0$



- 8 The volume of the solid generated by revolving the region bounded by the curve  $f(x) = x^2$  and  $x$ -axis and the two straight lines  $x = -2$  ,  $x = 2$  a complete revolution about  $x$ -axis equals .....

(a)  $\frac{16\pi}{5}$  (b)  $\frac{32\pi}{5}$  (c)  $\frac{64\pi}{5}$  (d)  $4\pi$

- 9 A 3-metre wall is 3 metres away from a house , then the minimum length of the ladder that joined the ground and the house resting on the wall = ..... metre.

(a) 3 (b) 6 (c)  $3\sqrt{2}$  (d)  $6\sqrt{2}$

- 10 If  $\sin x = xy$  , then :  $x^2(y + \dot{y}) + 2 \cos x = \dots\dots\dots$

(a) 0 (b)  $2x$  (c)  $2y$  (d)  $y$


- 11 The local minimum value of the function  $f : f(x) = x + \frac{1}{x}$  is at  $x = \dots\dots\dots$

(a) -1 (b) zero (c) 1 (d) 2


- 12 If  $y = x^x$  ,  $x > 0$  , then  $\frac{dy}{dx} = \dots\dots\dots$

(a)  $\ln x$  (b)  $2 + \ln x$   
(c)  $x^x \ln x$  (d)  $x^x (1 + \ln x)$


- 13 If  $f'(x) < 0$  ,  $f''(x) > 0$  for every  $x \in [a, b]$  , which of the following shown curves represents the curve of the function  $f$  in the interval  $[a, b]$  ?




(a)



(b)



(c)



(d)

- 14 The function  $f : f(x) = 2 \ln x - x^2$  is decreasing on the interval .....


(a)  $]0, 1[$  (b)  $] -\infty, 1[$  (c)  $]1, \infty[$  (d)  $]0, \infty[$




- 15 A cuboid whose dimensions at an instant are 3, 4, 12 cm., the first dimension increases at a rate 2 cm./sec., and the second dimension increases at a rate 1 cm./sec. and third dimension decreases at a rate 3 cm./sec., then the rate of change of volume of the cuboid after 2 seconds = ..... cm<sup>3</sup>/sec.

(a) 528 (b) -12 (c) 96 (d) 252


- 16 From the following figures which one represents the general shape of the curve of the continuous function  $f$  where  $f(1) = f(5) = \text{zero}$ ,  $f(3) = -3$ ,  $f'(x) < 0$  for every  $x < 3$ ,  $f'(x) > 0$  for every  $x > 3$   
 $f''(x) < 0$  for every  $x \neq 3$ ?




(a)



(b)



(c)



(d)

- 17 25 metre rope passes over a pulley which is 12 m. high. One of its end tied to a heavy mass and the other end tied to a car moves on the ground with velocity 6 m./sec. away from the projection of the pulley on the ground, then the rate of change of height of the mass at the moment when the car at a distance 16 m. from the projection of the pulley = ..... m/sec.

(a) 7 (b) 4.8 (c) 9.6 (d) 6

- 18 If  $x = \sin y$  where  $y$  is an acute angle, then  $\frac{dy}{dx} = \dots\dots\dots$

(a)  $\sqrt{1-x^2}$  (b)  $\frac{1}{\sqrt{1-x^2}}$  (c)  $\sqrt{x^2-1}$  (d)  $\frac{1}{\sqrt{x^2-1}}$

- 19 The equation of the curve :  $y = f(x)$ . If the slope of the normal at any point on the curve is  $(2y+1) \csc x$  and the curve passes through the origin is .....

(a)  $y^2 + y = \cos x - 1$  (b)  $y^2 + y = -\cos x - 1$   
 (c)  $y^2 + y = \csc x \tan x$  (d)  $y^2 + 1 = \cos x$




## 20 In the opposite figure :

Area of region located in the first quadrant and including between the curves :

$x + y = 3$  ,  $x^2 = 4y$  ,  $y^2 = 4x$   
equals ..... square unit.


- (a)  $\frac{5}{4}$  (b)  $\frac{10}{3}$   
(c)  $\frac{55}{12}$  (d)  $\frac{13}{6}$



## 21 In the opposite figure :

The straight line L is a tangent to the function  $f$  at the point C and intersects the  $x$ -axis at the point A  $(-4, 0)$ , the coordinates of the point B  $(4, 0)$  and  $f(4) + f'(4) = 9$ , then the area of  $\triangle ABC$  = ..... square unit.

- (a) 36 (b) 72 (c) 32 (d) 18

22 The equation of the tangent to the curve  $y = \ln(e^{2x} + e^x + 1)$  at  $x = 0$  is .....

- (a)  $y = x + \ln 3$  (b)  $y = x - \ln x$   
(c)  $3y = x + 3 \ln 3$  (d)  $y = 3x + \ln 3$


23 If  $\tan(x^2 + y^2) = 0$ , then  $\frac{dy}{dx} =$  .....

- (a)  $\frac{x}{y}$  (b)  $\frac{y}{x}$  (c)  $-\frac{x}{y}$  (d)  $-\frac{y}{x}$

## 24 From the opposite figure :

$${}_2 \int_5 \frac{f'(x)}{f(x)} dx = \dots\dots\dots$$

- (a)  $\ln 2$   
(b)  $\ln 3$   
(c)  $\log_9 5$   
(d)  $\ln 6$

25 If  ${}_2 \int_k \frac{dx}{4x} = \ln 2$ , then  $k =$  ..... where  $k > 2$ 

- (a) 64 (b) 48 (c) 36 (d) 32

**Answer the following questions :**

1 If  $f(\sin X) = \sin^2 X$ , then  $f'(1) = \dots\dots\dots$

(a) 1

(b) 2

(c)  $\pi$

(d)  $\frac{\pi}{2}$

2 The curve  $\left(\frac{x}{a}\right)^t + \left(\frac{y}{b}\right)^t = 2$  touches the straight line  $\frac{x}{a} + \frac{y}{b} = 2$  at the point (a, b) when .....

(a)  $t = 3$  only.

(b)  $t = 2$  only.

(c) for all values of  $t$

(d) not true for any  $t$

3 The normal equation to the curve  $y = e^{2x} \cos x$  at  $x = 0$  is .....

(a)  $x + y = 2$

(b)  $2y + x = 2$

(c)  $2x + y = 2$

(d)  $x - y = 2$

4 A circular segment, the radius length of its circle is 10 cm. and measure of its central angle ( $x$ ) changes in the rate  $3^{\text{rad}}$ . per minute, then the rate of increasing in the area of the circular segment at  $x = \frac{\pi}{3}$  is .....  $\text{cm}^2/\text{min}$ .

(a) 125

(b) 75

(c) 150

(d) 300

5 The function  $f : f(x) = 3 - \ln x^2$  increases in the interval .....

(a)  $]-\infty, \infty[$

(b)  $]-\infty, 0[$

(c)  $]0, \infty[$

(d)  $]3, \infty[$

6  $\lim_{h \rightarrow 0} \frac{\cot(x+h) - \cot(x)}{h} = \dots\dots\dots$

(a)  $-\csc^2 x$

(b)  $\sec^2 x$

(c)  $-\cot^2 x$

(d)  $\cot x \csc x$



7 The curve of tangent slope at any point on it equals a  $\csc^2 X$  where  $a$  is a constant, and the curve passes through the two points  $(\frac{\pi}{4}, 5)$ ,  $(\frac{3\pi}{4}, 1)$ , then the equation of the curve is .....

(a)  $y = 2 \cot X + 3$

(b)  $y = -2 \cot X - 3$

(c)  $y = -\cot X$

(d)  $y = 2 \cot X - 3$

8 A circular sector whose perimeter is 30 cm., and its area is as great as possible, then the radius length of its circle = .....

(a) 15

(b) 30

(c)  $\frac{15}{4}$

(d) 7.5

9 The curve  $y = X^3 - 6X^2$  is convex downwards in the interval .....

(a)  $\mathbb{R} - ]0, 4[$

(b)  $]0, 4[$

(c)  $]2, \infty[$

(d)  $] -\infty, 2[$

10  $\int \sec^{2017} X \tan X dX = \dots + c$

(a)  $\frac{1}{2018} \sec^{2018} X$

(b)  $\frac{1}{2016} \sec^{2016} X$

(c)  $\frac{1}{2017} \sec^{2017} X$

(d)  $\frac{1}{2015} \sec^{2015} X$

11 In the opposite figure :

If  $A_2$  is the region bounded by the two curves

$y^2 = 4X, X^2 = 4y$


, then  $A_1 : A_2 : A_3 = \dots$

(a) 2 : 1 : 2

(b) 1 : 2 : 1

(c) 1 : 1 : 1

(d) 3 : 2 : 3



12 The rate of change of  $(X - \sin X)$  with respect to  $(1 - \cos X)$  at  $X = \frac{\pi}{3}$  equals .....

(a)  $\frac{\sqrt{3}}{3}$

(b)  $\sqrt{3}$


(c)  $2\sqrt{3}$

(d)  $\frac{2}{3}$



- 13 The opposite figure represents the curve  $\hat{f}(x)$ , the inflection is at  $x = \dots\dots\dots$


- (a) zero  
(b) -3  
(c) 4  
(d) all the previous.



- 14 In the opposite figure :

$\overrightarrow{AC}$  is a tangent to the curve  $y^2 = 18x$  at the point  $A(2, 6)$  and  $\overrightarrow{AB} \perp \overrightarrow{AC}$ , then the length of  $\overline{BC} = \dots\dots\dots$  length unit.

- (a) 9  
(b) 11  
(c) 2  
(d) 13



- 15 If  $y = f(x)$  represents a curve of polynomial function of third degree and  $\hat{f}(x) < 0$  at  $x < \frac{-2}{3}$ ,  $\hat{f}(x) > 0$  at  $x > \frac{-2}{3}$  and passes through the point  $(1, 6)$  and there is a critical point at  $(-1, 2)$ , then the equation of the curve is  $\dots\dots\dots$

- (a)  $y = 2x^3 + 4$   
(b)  $y = 2x^3 + 4x^2$   
(c)  $y = x^3 + 2x^2 + x + 2$   
(d)  $y = x^3 + 2x^2 + 3$

- 16 The maximum value of the expression  $(\sin x + \cos x)$  is  $\dots\dots\dots$

- (a) 1  
(b) 2  
(c)  $\sqrt{2}$   
(d)  $\frac{1}{\sqrt{2}}$


- 17 If  $f(x) = x^2$ ,  $g(x) = \cot x$  and  $h(x) = (f \circ g)(x)$ , then  $h\left(\frac{\pi}{4}\right) = \dots\dots\dots$

- (a) -4  
(b) 4  
(c) zero  
(d) -1



- 18 The opposite figure shows a graphical representation to the curves  $f(x)$ ,  $f'(x)$ ,  $f''(x)$  where  $f(x)$  is polynomial, then the curves a, b, c represent in order .....

- (a)  $f(x)$ ,  $f'(x)$ ,  $f''(x)$   
 (b)  $f'(x)$ ,  $f''(x)$ ,  $f(x)$   
 (c)  $f''(x)$ ,  $f(x)$ ,  $f'(x)$   
 (d)  $f''(x)$ ,  $f'(x)$ ,  $f(x)$




- 19 An engineer design a hotel in form of the curve whose equation  $y = 4x - \frac{1}{2}x^2$  where  $x$  is in metres. If it is covered by glass, the cost of one square metre is L.E. 1200, then the cost of the glass = L.E. ....

- (a)  $\frac{128}{3}$  (b) 715200 (c) 51200 (d) 15200

- 20 In the opposite figure :

If the straight line  $L$  is a tangent to the curve  $y = f(x)$  at the point  $A(2, k)$ , then  $\int_{-2}^2 [f''(x) + f'(x)] dx = \dots\dots\dots$

- (a)  $4\sqrt{3}$  (b)  $2\sqrt{3}$   
 (c) 3 (d) 2



- 21  $\int e^x (f''(x) + f'(x)) dx = e^x \times \dots\dots\dots + c$

- (a)  $x$  (b)  $f(x)$  (c)  $f'(x)$  (d)  $f''(x)$

- 22 The function  $f : f(x) = \begin{cases} 5 - x^2 & , -3 \leq x \leq 2 \\ x^2 - 3 & , 2 < x \leq 3 \end{cases}$

then the function has an absolute minimum value = .....

- (a) 5 (b) 1 (c) -5 (d) -4



- 23 If  $f$  is a polynomial function of fifth degree, then the greatest possible number of inflection points is .....

(a) 2

(b) 3


(c) 4

(d) 5

- 24 The opposite figure represents the curve :

$$y = \frac{\ln x}{\sqrt{x}} \text{ and the line } x = e, \text{ then}$$

the volume of the generated solid  
by revolving the shaded region  
a complete revolution about  
the  $x$ -axis = ..... cube units.

(a)  $\frac{1}{3} \pi$ (b)  $\pi$ (c)  $3 \pi$ (d)  $9 \pi$ 

- 25 A man observes a plane flies horizontally at 3 km. high exactly above him and with speed 480 km/h. , then the rate of change of the distance between the man and the plane after 30 sec. later = .....

(a)  $\frac{320}{3}$  km/h.

(b) 384 km/h.

(c) 384 m./sec.

(d)  $\frac{320}{3}$  cm./sec.



# School Book Examinations

in

## Differential & Integral calculus







### First Answer the following question

1 Choose the correct answer :

(1) Which of the following functions satisfies the relation  $\frac{d^3 y}{d x^3} = y$  ?

- (a)  $\frac{1}{12} (x+1)^4$       (b)  $\sin x$       (c)  $e^x$       (d)  $\frac{x}{x-1}$

(2) If the radius length of a circle increases at a rate  $\frac{1}{\pi}$  cm./sec. the circumference of the circle increases at a rate of ..... cm./sec.

- (a)  $\frac{2}{\pi}$       (b) 2      (c)  $\pi$       (d)  $2\pi$

(3) The curve of the function  $f$  where  $f(x) = x^3 - 3x^2 + 2$  is convex upwards when  $x \in$  .....

- (a)  $]-\infty, 0[$       (b)  $]-\infty, 1[$       (c)  $]1, 3[$       (d)  $]1, \infty[$

(4)  $\int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} (\sin x + \cos x) dx$  equals .....

- (a) 4      (b) 2      (c) zero      (d)  $\pi$

(5) If  $f$  is a continuous function on  $\mathbb{R}$ ,  $\int_3^5 2f(x) dx = 8$ ,  $\int_3^4 3f(x) dx = 9$ , then  $\int_4^5 5f(x) dx$  equals .....

- (a) zero      (b) 1      (c) 3      (d) 5

(6) The area of the region bounded by the curve  $y = \sqrt{16 - x^2}$  and  $x$ -axis measured in square units equals .....

- (a)  $16\pi$       (b)  $12\pi$       (c)  $8\pi$       (d)  $4\pi$

### Second Answer three questions only of the following

2 [a] Find :  $\int \sin x \cos^3 x dx$

$$\ll -\frac{1}{4} \cos^4 x + c \gg$$

[b] If  $e^{xy} - x^2 + y^3 = 0$ , find :  $\frac{dy}{dx}$  when  $x = 0$

$$\ll \frac{1}{3} \gg$$

3 [a] Find the equation of the tangent to the curve :  $x^2 - 3xy - y^2 + 3 = 0$  at point  $(-1, 4)$

$$\ll x + 5y - 6 = 0 \gg$$



[b] The lengths of the legs of the right-angle of a right-angled triangle at a moment  
 , are 6 cm. and 30 cm. If the length of the first leg increases at a rate of  $\frac{1}{3}$  cm./min.  
 and the length of the second leg decreases at a rate of 1 cm./min. , **find :**

( 1 ) The rate of increase in the area of the triangle after 3 minutes.

( 2 ) The time at which the increase of the area of the triangle stops. « 1 cm<sup>2</sup>/min. , 6 »

4 [a] Determine the increasing and decreasing intervals to the function  $f$  where :

$$f(x) = x + 2 \sin x \quad , \quad 0 < x < 2\pi$$

[b] A rectangle is drawn such that two adjacent vertices of the rectangle lie on the curve  
 $y = x^2 - 12$  and the other two vertices lie on the curve  $y = 12 - x^2$  , find the maximum  
 area of this rectangle. « 64 »

5 [a] Find the volume of the solid generated by revolving the region bounded by the two  
 curves  $y = \frac{4}{x}$  and  $y = (x - 3)^2$  a complete revolution about  $x$ -axis « 5.4  $\pi$  cubic unit »

[b] Sketch the curve of the function  $f$  which satisfies the following properties :

( 1 )  $f(1) = f(5) = 0$  ,  $f(2) = -3$

( 2 )  $f'(x) < 0$  for each  $x \neq 2$

( 3 )  $f'(x) < 0$  for each  $x < 2$

( 4 )  $f'(x) > 0$  for each  $x > 2$



### First Answer the following question

1 Choose the correct answer :

(1) The equation of the tangent to the curve of the function  $f$  where  $f(x) = e^{2x+1}$  at point  $(-\frac{1}{2}, 1)$  is .....

- (a)  $2y = x + 1$       (b)  $y = 2x + 2$       (c)  $y = 2x - 3$       (d)  $2y = 3x + 1$

(2) If  $y = 4n^3 + 4$  ,  $z = 3n^2 - 2$  , then the rate of change of  $z$  with respect to  $y$  equals .....

- (a)  $2n$       (b)  $2$       (c)  $\frac{1}{2n}$       (d)  $4$

(3) The maximum value of the expression :  $8x - x^2$  where  $x \in \mathbb{R}$  is .....

- (a)  $8$       (b)  $16$       (c)  $32$       (d)  $64$

(4) If the slope of the tangent to the curve of the function  $f$  at any point on it equals  $\frac{1}{x-2}$  and the curve passes through point  $(3, 0)$  , then  $f(e^2 + 2)$  equals .....

- (a)  $2$       (b)  $3$       (c)  $\ln 2$       (d)  $\ln 3$

(5) If  $f$  is a continuous function on  $\mathbb{R}$  ,  $\int_1^2 f(x) dx = 9$  and  $\int_6^2 f(x) dx = -7$  , then  $\int_1^6 f(x) dx$  equals .....

- (a)  $2$       (b)  $8$       (c)  $16$       (d)  $-63$

(6) The volume of the solid generated by revolving the region bounded by the curve  $y = \sqrt{x+1}$  and the straight lines  $y = 0$  ,  $x = -1$  and  $x = 1$  equals .....

- (a)  $\pi$       (b)  $\frac{3\pi}{2}$       (c)  $2\pi$       (d)  $\frac{5\pi}{2}$

### Second Answer three questions only of the following

2 [a] Find : (1)  $\int x(2x-1)^3 dx$

(2)  $\int x e^{-2x} dx$

[b] Find the rate of change for :  $\sqrt{16+x^2}$  with respect to  $\frac{x}{x-2}$  when  $x = -3$

« $\frac{15}{2}$ »



3 [a] If  $x \cos y + y \cos x = 1$ , find :  $\frac{dy}{dx}$

[b] Find the absolute extrema values of the function  $f$  in the interval  $[-1, 1]$

where  $f(x) = 2x^3 + 6x^2 + 5$

« 13, 5 »

4 [a] If  $f(x) = \begin{cases} 2x + x^2 & \text{when } x < 0 \\ 2x - x^2 & \text{when } x \geq 0 \end{cases}$

Find : (1) The local maximum and minimum values of the function  $f$

(2)  $\int_{-1}^3 f(x) dx$

«  $-\frac{2}{3}$  »

[b] The volume of a cube increases regularly such that it keeps its shape at a rate of  $27 \text{ cm}^3/\text{min.}$ , find the increase of the area of its faces at the moment which its edge length is 3 cm.

«  $36 \text{ cm}^2/\text{min.}$  »

5 [a] Find the area of the region bounded by the two curves :

$y = x^2$  and  $y = 6x - x^2$  (in square units).

« 9 square unit »

[b] If the function  $f$  where  $f(x) = x^3 + ax^2 + bx$  has an inflection point at  $(2, 2)$ , find the two values of the two constants  $a$  and  $b$ , then sketch the curve of the function.

« -6, 9 »



### First Answer the following question

1 Choose the correct answer :

(1) The slope of the tangent to the curve of the circle :  $x^2 + y^2 = 25$   
when  $x = 3$  equals .....

- (a)  $-\frac{4}{3}$  (b)  $-\frac{3}{4}$  (c)  $\frac{5}{12}$  (d)  $\frac{4}{3}$

(2) If  $f(x) = \frac{x}{x-2}$ , then  $f'(3) = \dots\dots\dots$

- (a) -36 (b) -12 (c) 6 (d) 4

(3) If  $\frac{dy}{dx} = \csc^2 x$ ,  $y = 2$  and  $x = \frac{\pi}{4}$ , then  $y$  equals .....

- (a)  $-(2 + \cot x)$  (b)  $-(3 + \cot x)$  (c)  $2 - \cot x$  (d)  $3 - \cot x$

(4) If  $\int_2^4 f(x) dx = 7$ ,  $\int_4^2 g(x) dx = 2$ , then  $\int_2^4 [2f(x) - 3g(x) - 5] dx$   
equals .....

- (a) -18 (b) -8 (c) 10 (d) 14

(5) The area of the region bounded by the straight lines :

$y = 2x - 3$ ,  $y = x + 1$ ,  $x = 2$  equals .....

- (a) 2 (b) 3 (c)  $\frac{9}{2}$  (d) 6

(6) The volume of the solid generated by revolving the region bounded by the two curves

$y = \tan x$ , and  $y = \sec x$  and two straight lines  $x = \frac{\pi}{6}$ ,  $x = \frac{\pi}{3}$

a complete revolution about  $x$ -axis approximated in cubic units equals .....

- (a)  $\frac{\pi^2}{6}$  (b)  $\frac{\pi^2}{3}$  (c)  $\frac{2\pi^2}{5}$  (d)  $2\pi^2$

### Second Answer three questions only of the following

2 [a] Find the derivative of  $y$  with respect to  $x$  where  $y = x^2 \ln x$

«  $x(2 \ln x + 1)$  »

[b] If  $f(x) = \sqrt[3]{(x-4)^2}$ , find the convexity intervals upwards and downwards and the inflection points (if existed) to the curve of the function  $f$



3 [a] Find : (1)  $\int x(x-5)^3 dx$

(2)  $\int 4x e^{2x} dx$

[b] Find the absolute maximum values of the function  $f$  where :

$f(x) = x^4 - 4x^3$  on the interval  $[0, 4]$

« 0 »

4 [a] The volume of a solid of revolution generated by revolving the region bounded by the curve  $y = x^3$  and the two straight lines  $x = 0$  and  $y = 1$  a complete revolution about  $x$ -axis is equal to the volume of a cylinder-like wire whose length is 42 units.

What is the radius length of that wire ?

«  $\frac{1}{7}$  length unit »

[b] The two equal legs of the isosceles triangle with a constant base whose length is  $l$  cm. decrease at a rate of 3 cm./min. What is the rate of decrease in the area when the triangle becomes an equilateral triangle ?

«  $\sqrt{3} l \text{ cm}^2/\text{min.}$  »

5 [a] Find the area of the region bounded by the two curves :  $x - y = 0$  ,  $y = 4x - x^2$

«  $\frac{9}{2}$  square unit »

[b] Sketch the curve of the continuous function  $f$  which has the following properties :

(1)  $f(0) = 3$

(2)  $\hat{f}(2) = \hat{f}(-2) = 0$

(3)  $\hat{f}(x) > 0$  when  $-2 < x < 2$

(4)  $\hat{\hat{f}}(x) < 0$  when  $x > 0$  ,  $\hat{\hat{f}}(x) > 0$  when  $x < 0$



### First Answer the following question

1 Choose the correct answer :

(1) If  $y = \frac{3x-5}{x-2}$ , then at  $x = 1$ ,  $\frac{d^3 y}{d x^3}$  equals .....

- (a) -12 (b) -6 (c) 6 (d) 12

(2)  $\int \sec^3 x \tan x \, dx$  equals .....

- (a)  $\frac{1}{4} \sec^4 x + c$  (b)  $\frac{1}{3} \sec^3 x + c$  (c)  $\frac{1}{2} \tan^2 x + c$  (d)  $-\frac{1}{2} \tan^2 x + c$

(3) The normal to the circle  $x^2 + y^2 = 12$  at any point on it passes through the point .....

- (a) (2, 3) (b) (1, 1) (c) (0, 0) (d) (-2, -2)

(4) The curve of the function  $f$  where  $f(x) = (x-2)e^x$  is convex downwards on the interval .....

- (a)  $]-\infty, \infty[$  (b)  $]-1, 2[$  (c)  $]0, 2[$  (d)  $]0, \infty[$

(5)  $\int_{-1}^3 3x|x-4| \, dx$  equals .....

- (a) -27 (b) -20 (c) 20 (d) 27

(6) When the region bounded by the curve  $x = \frac{1}{\sqrt{y}}$ ,  $1 \leq y \leq 4$  and y-axis revolves a complete revolution about y-axis, then the volume of the solid generated measured in cubic units equals .....

- (a)  $\frac{2}{3} \pi$  (b)  $3\sqrt{2} \pi$  (c)  $2 \pi \ln 2$  (d)  $\frac{2}{3} \pi \log 3$

### Second Answer three questions only of the following

2 [a] Find : (1)  $\int (3x^2 - 4e^{2x}) \, dx$  (2)  $\int \frac{x-1}{\sqrt{x+3}} \, dx$

[b] If  $\sin y + \cos 2x = 0$  Prove that :  $\frac{d^2 y}{d x^2} - \left(\frac{d y}{d x}\right)^2 \tan y = 4 \cos 2x \sec y$

3 [a] If  $\int_1^4 f(x) \, dx = 7$ ,  $\int_4^1 g(x) \, dx = 3$

Calculate the value of :  $\int_1^4 [f(x) + 2g(x) - 4] \, dx$

« - 11 »



- [b] If the curve of the function  $f$  where  $f(x) = ax^3 + bx^2 + cx + d$  has a local maximum value at  $(2, 4)$  and an inflection point at  $(1, 2)$ , find the equation of the curve.

$$« f(x) = -x^3 + 3x^2 »$$

- 4 [a] Find the area of the region bounded by the curve :

$$\sqrt{x} + \sqrt{y} = 1 \text{ and the two straight lines } x = 0, y = 0$$

$$« \frac{1}{6} \text{ square unit } »$$

- [b] Graph the curve of the continuous function  $f$  which satisfies the following properties :

$$(1) f(4) = 2, f(3) = 4$$

$$(2) f(2) = 0$$

$$(3) f'(x) < 0 \text{ when } x > 4 \text{ or } x < 2$$

$$(4) f'(x) < 0 \text{ when } x > 3, f'(x) > 0 \text{ when } x < 3$$

- 5 [a] Prove that the volume of the solid generated by revolving the region bounded by

the two curves  $y = \frac{4}{x}$  and  $y = 5 - x$  just one revolution about  $x$ -axis equals  $9\pi$  of the cubic units.

- [b] If  $A$  is the area of the part bounded by two concentric circle whose radii lengths are

$r_1$  and  $r_2$  where  $r_2 > r_1$ , find the rate of change of  $A$  with respect to time at any

moment at which  $r_2 = 10$  cm. ,  $r_1 = 6$  cm. , if known that at this moment  $r_1$

increases at a rate of  $0.3$  cm/s. and  $r_2$  decreases at a rate of  $0.2$  cm/s. «  $-7.6\pi \text{ cm}^2/\text{sec.}$  »




### First Answer the following question

- 1 The opposite figure shows the curve  $\hat{f}(x)$  of the function  $f$  where  $f(x) = ax^3 + bx^2$ ,  $a, b$  are two constants.

Complete :

- (1) The function  $f$  is decreasing for each  $x \in \dots\dots\dots$
- (2) The curve of  $f$  has critical points when  $x \in \dots\dots\dots$
- (3) The curve of  $f$  is convex upwards on the interval  $\dots\dots\dots$
- (4) There is a local minimum value of the function  $f$  when  $x = \dots\dots\dots$
- (5)  $f(1) = \dots\dots\dots$
- (6) The area of the region bounded by the curve of the function  $f$  and the two straight lines  $x = 2$ ,  $x = 0$  and  $y = 0$  in square units equals  $\dots\dots\dots$



### Second Answer three questions only of the following

- 2 [a] Find :

$$(1) \int \csc^2\left(\frac{x+5}{2}\right) dx$$

$$(2) \int \frac{5x}{3x^2-1} dx$$

- [b] The function  $f$  where  $f(x) = x^3 - 6x^2 + 9x - 1$

- (1) Determine the increasing and decreasing intervals of function  $f$
- (2) Find the maximum values of the function  $f$  in the interval  $[0, 2]$

- 3 [a] If  $f(x) = 4 + \cot x - \sec^2 x$ , find the equation of the normal to the curve of the function  $f$  at a point lying on the curve and its  $x$ -coordinate equals  $\frac{\pi}{4}$

$$\ll 4x - 24y - \pi + 72 = 0 \gg$$

- [b] An empty tank whose capacity is 10 cubic metres. If the water is poured gradually in that tank at a rate of  $(2t + 3) \text{ m}^3/\text{min}$ , where  $t$  time in minutes, find the time needed to fill the tank.

$$\ll 2 \text{ min. } \gg$$



4 [a] Find :  $\lim_{x \rightarrow \infty} \left( \frac{2x-1}{2x+1} \right)^{2x}$

«  $\frac{1}{e}$  »

[b] A rectangle - like poster contains  $800 \text{ cm}^2$  of the printed material where the widths of both lower and upper margins are 10 cm. and the two side margins are 5 cm. what are the two dimensions of the posters which make its area as minimum as possible.

« 60 , 30 cm. »

5 [a] Find the volume of the solid generated by revolving the region bounded by the curve  $y = 4 - x^2$  and the two positive parts of the axes of coordinates a complete revolution about X-axis.

«  $\frac{256}{15} \pi$  Cubic unit »

[b] If  $f(x) = x^3 + ax^2 + bx + 4$  where a and b are two constants , find the two values of a and b if the function f has a local minimum value when  $x = 2$  and an inflection point when  $x = 1$  , then sketch the curve of the function f



### First Answer the following question

- 1 In each of the following phrases, choose (a) if the phrase is true and (b) if the phrase is false :
- (1) The local maximum value of the function is greater than its local minimum value. (a) (b)
- (2) The rate of change of  $\sqrt{n^2 + 3}$  with respect to  $\frac{n}{n+1}$  is  $\frac{n(n+1)^2}{\sqrt{n^2 + 3}}$  (a) (b)
- (3) If  $\sqrt{y} - \sqrt{x} = 2$ , then  $\frac{d^2 y}{d x^2} = \frac{-1}{x\sqrt{x}}$  (a) (b)
- (4)  $\int \frac{x-4}{(x-2)^6} dx = \frac{7(x-4)^2}{2(x-2)^7} + c$  (a) (b)
- (5) If  $y = x \ln x - x$ , then  $\frac{dy}{dx} = \ln x$  (a) (b)
- (6) If  $(a, f(a))$  is an inflection point to the curve of the continuous function  $f$ , then  $f''(a) = \text{zero}$  (a) (b)

### Second Answer three questions only of the following

2 [a] Find :

$$(1) \int \frac{7x^3}{2-5x^4} dx$$

$$(2) \int \left( 3e^{-5x} + \frac{\pi}{x} \right) dx$$

[b] If  $y = a e^{x^2+1}$  Prove that :  $\frac{d^3 y}{d x^3} = 4xy(3+2x^2)$

3 [a] Find :  $\int \cot x \csc^3 x dx$

[b] If  $s$  is the distance between point  $(1, 0)$  and point  $(x, y)$  lying on the curve  $y = \sqrt{x}$ , find the coordinates of point  $(x, y)$  at which  $s$  is as minimum as possible.  $\left( \frac{1}{2}, \frac{1}{\sqrt{2}} \right)$

4 [a] Identify the absolute extrema values of the function  $f$  where  $f(x) = |x|(x-4)$  in the interval  $[-1, 3]$



[b] If the slope of the tangent to the curve  $y = f(x)$  at any point on it equals  $6x^2 + b$  and  $f(0) = 5$ ,  $f(2) = -3$ , find the value of the constant  $b$ , then sketch the curve of the function  $f$

«  $b = -12$  »

5 [a] Find the rate of change of  $\ln(9 + x^3)$  with respect to  $x^2 + 3$  and  $x = 1$

«  $\frac{3}{20}$  »

[b] If  $A(0, 3)$ ,  $B(1, 4)$ ,  $C(2, 0)$  Find using integration :

(1) The surface area of  $\Delta ABC$

(2) The volume of the solid generated by revolving  $\Delta AOC$  a complete revolution about y-axis.

«  $\frac{5}{2}$  square unit,  $4\pi$  cubic unit »



### First Answer the following question

1 In each of the following phrases, choose (a) if the phrase is true and (b) if the phrase is false :

(1) If  $y^2 = 3x^2 - 7$ , then  $\frac{dy}{dx} = \frac{y}{3x}$  (a) (b)

(2) The function  $f : f(x) = x^3 - 3x + 1$  has an inflection point which is (0, 1) (a) (b)

(3)  $\frac{d}{dx} [\cot(\cos 3x)] = 3 \sin 3x \csc^2(\cos 3x)$  (a) (b)

(4)  $\int (1 - \cos x)^4 \sin x \, dx = -\frac{1}{5} (1 + \cos x)^5 + c$  (a) (b)

(5)  $\lim_{x \rightarrow \infty} \left(1 + \frac{5}{x}\right)^x = e^5$  (a) (b)

(6)  $\int \left(\frac{2e}{x} + \frac{x}{e}\right) dx = 2e \ln|x| - \frac{x^2}{e} + c$  (a) (b)

### Second Answer three questions only of the following

2 [a] Find :

(1)  $\int x \sin x \, dx$

(2)  $_{-1} \int^1 \sqrt{x^4 + x^2} \, dx$

[b] Find the equation of the tangent to the curve  $y = \ln(2 - \sqrt{2} \cos x)$  at the point lying on it and its  $x$ -coordinate equals  $\frac{\pi}{4}$  «  $x - y - \frac{\pi}{4} = 0$  »

3 [a] Identify the convexity intervals downwards and upwards and the inflection points (if existed) to the curve of the function  $f$  where  $f(x) = (x - 1)^4 + 3$

[b] A cuboid of metal whose base is square. If the side length of the base increases at a rate of 0.4 cm./sec. and the height decreases at a rate of 0.5 cm./sec., find the rate of change of the volume when the side length of the base is 6 cm. and the height is 5 cm.

« 6 cm.<sup>3</sup>/sec. »



4 [a] Find :  $\int_0^3 x\sqrt{x+1} \, dx$

«  $\frac{116}{15}$  »

[b] A rectangle - like playground in which two opposite sides end in a semi-circle outside the rectangle of a diameter length equal to the length of this side.

If the perimeter of the playground is 400 metres , prove that the surface area of the playground is as maximum as possible when the ground is a circle - like , then find its radius length.

«  $\frac{200}{\pi}$  »

5 [a] If  $f(x) = x^3 - 3x + 3$  , find :

( 1 ) The absolute extrema value of the function  $f$  in the interval  $f [0 , 2]$

( 2 ) The area of the region bounded by the curve of the function  $f$  and the straight lines  $x = 0$  ,  $x = 2$  ,  $y = 0$

« 4 square unit »

[b] Find the volume of the solid generated by revolving the region bounded by the curve

$xy = 2$  and the two straight lines  $x = 1$  and  $x = 2$  about  $x$ -axis

«  $2\pi$  cubic unit »



### First Answer the following question

1 Complete the following :

(1) If  $x^3 y^2 = 1$ , then  $\left[\frac{dy}{dx}\right]_{y=1} = \dots\dots\dots$

(2)  $\frac{d}{dx} [7 e^{\sec x}] = \dots\dots\dots$

(3) The function  $f : f(x) = x^3 - 3x - 1$  has an inflection point which is  $\dots\dots\dots$

(4) If  $f$  is a continuous function on the interval  $[2, 7]$

, then  $\int_2^7 f(x) dx + \int_7^4 f(x) dx = \dots\dots\dots$

(5) The area of the region bounded by the two curves  $y = x^2$  and  $y = 4x$  equals  $\dots\dots\dots$  square units.

(6) If  $y = x^2 \ln \frac{x}{a}$ ,  $a \neq 0$ , then  $\left[\frac{d^3 y}{dx^3}\right]_{x=4} = \dots\dots\dots$

### Second Answer three questions only of the following

2 [a] Find :

(1)  $\int \frac{(x+3)^3 - 27}{x} dx$

(2)  $\int x^2 e^{-x} dx$

[b] Find the equation of the tangent to the curve of the function  $f$  where  $f(x) = 2 \tan^3 x$  at the point lying on the curve of the function  $f$  and its  $x$ -coordinate equals  $\frac{\pi}{4}$

«  $y = 12x - 3\pi + 2$  »

3 [a] Find :  $\int_0^5 |x-2| dx$

«  $\frac{13}{2}$  »


[b] The opposite figure shows the two curves of the two functions  $g$  and  $z$  where :

$g(x) = \ddot{f}(x)$ ,  $z(x) = \ddot{\ddot{f}}(x)$  and  $f$  is

a polynomial function at the variable  $x$

Sketch the curve of  $f$  knowing that it passes through the two points

$(-1, 0)$ ,  $(1, 4)$





- 4 [a] Identify the absolute extrema values of the function  $f$  in the interval  $[0, 2]$ , where  $f(x) = 3\sqrt{4-x^2}$

- [b] A five metre length rod is fixed by a hinge to the ground at its base. If its top rises up by a winch at a rate of 1 m./min., find the rate of decreasing the projector length of the rod on the ground when the height of the top is 3 metres. «  $\frac{3}{4}$  m./min. »

- 5 [a] If a trapezoid is drawn in a semi-circle such that its base is the diameter of the semi-circle, determine the measure of the angle of the trapezoid base such that its area is as maximum as possible. «  $60^\circ$  »

- [b] If  $a$  is the region bounded by the curve  $xy = 4 + x^2$  and the straight lines  $x = 1$ ,  $x = 4$  and  $y = 0$ , find :

(1) The area of region  $a$  in square units to the nearest unit. «  $4 \ln 4 + \frac{15}{2}$  square unit »

(2) The volume of the solid generated by revolving the region about  $x$ -axis.

«  $57\pi$  cubic unit »



### First Answer the following question

1 Choose the correct answer :

(1) If  $x = 2n^2 + 7$ ,  $y = \sqrt{n^3}$ ,  $n = 1$ , then  $\frac{dy}{dx}$  equals .....

- (a)  $\frac{3}{8}$  (b)  $\frac{3}{4}$  (c) 2 (d) 6

(2) The curve of the function  $f$  is convex downwards on  $\mathbb{R}$  if  $f(x)$  equals .....

- (a)  $2 - x^2$  (b)  $2 + x^3$  (c)  $2 - x^4$  (d)  $2 + x^4$

(3) If the curve of the function  $f : f(x) = x^3 + kx^2 + 4$ ,  $k \in \mathbb{R}$  has an inflection point when  $x = 2$ , then  $k =$  .....

- (a) -6 (b) -3 (c) 6 (d) 9

(4) If  $f$  is a continuous function on  $\mathbb{R}$ ,  $_{-1} \int^3 f(x) dx = 7$ ,  $_5 \int^3 f(x) dx = -11$ , then  $_{-1} \int^5 f(x) dx$  equals .....

- (a) -4 (b) 18 (c) -18 (d) 77

(5)  $_{-1} \int^3 |x - 1| dx$  equals .....

- (a) -6 (b) 0 (c) 4 (d) 8

(6) The area of the region bounded by the curve  $y = x^3$  and the two straight lines  $y = 0$  and  $x = 2$  equals .....

- (a) 1 (b) 2 (c) 4 (d) 8

### Second Answer three questions only of the following

2 [a] Find : (1)  $\int \frac{3x}{x^2 - 1} dx$  (2)  $\int 9x^2 e^{3x} dx$

[b] Find the measure of the positive angle which the tangent of the curve  $y^3 = x^2$  makes with the positive direction of  $x$ -axis when  $x = 8$  to the nearest minute.

« 18° 26' »



- 3 [a] If  $\sin x = xy$ , prove that :  $x^2(y + \dot{y}) + 2 \cos x = 2y$
- [b] If the curve  $y = 2x^3 + 3x^2 + 4x + 5$  has two parallel tangent, one of them touches the curve at point  $(-1, 2)$ , find the equation of the other tangent. «  $4x - y + 5 = 0$  »

- 4 [a] A balloon rises up vertically at a constant rate of 28 m./min. If the balloon is observed by a ground observer distant 200 m. away from the site of launching the balloon, find the rate of change of the angle of elevation of the observer when the balloon is 200 m. up. «  $0.07 \text{ rad./min.}$  »
- [b] If the slope of the tangent to the curve of the function  $f$  at any point  $(x, y)$  on the curve is  $3(x^2 - 1)$ , find the local maximum and minimum values to the curve of the function  $f$  and the inflection points if existed known that the curve passes through the point  $(-2, -1)$ , then sketch this curve.

- 5 The straight line  $\overleftrightarrow{AB}$  intersects the curve of the function  $f$  at point  $C(x, y)$ , where  $x > 0$ ,  $A(0, 2)$ ,  $B(6, 4)$  and  $f(x) = \frac{9}{x}$ , find :
- (1) The equation of the straight line  $\overleftrightarrow{AB}$  «  $y = \frac{1}{3}x + 2$  »
- (2) The coordinates of point  $C$  «  $(3, 3)$  »
- (3) The equation of the normal on the curve of  $f$  at point  $C$  and prove that it passes through the origin point  $O$  «  $x - y = 0$  »
- (4) The volume of the solid generated by revolving the region bounded by the normal  $\overleftrightarrow{OC}$  and the curve of the function and the straight line  $x = 6$  and  $x$ -axis a complete revolution about  $x$ -axis. «  $\frac{45}{2} \pi \text{ cubic unit}$  »



### First Answer the following question

#### 1 Complete :

(1)  $\lim_{x \rightarrow \infty} \left(1 + \frac{1}{x}\right)^{x+3} = \dots\dots\dots$

(2)  $\frac{d}{dx} (5 - 2 \cot x)^3 = \dots\dots\dots$

(3) If the function  $f : f(x) = kx^3 + 9x^2$  has an inflection point when  $x = -1$ , then  $k = \dots\dots\dots$

(4)  $\int_{-1}^3 (4x^3 - 6x^2 + 5) dx = \dots\dots\dots$

(5) If  $f$  is a continuous function on the interval  $[1, 4]$ , then  $\int_1^4 f(x) dx + 4 \int_1^4 f(x) dx = \dots\dots\dots$

(6) The area of the region bounded by the two curves  $y = x^4 + 1$  and  $y = 2x^2$  equals  $\dots\dots\dots$  square units.

### Second Answer three questions only of the following

2 [a] Find : (1)  $\int \tan(3x + 1) dx$

(2)  $\int (1 - x^2)(3x - x^3)^5 dx$

[b] If the two parametric equations of the function  $f$  where  $y = f(x)$  are :

$x = 2n^3 + 3$  and  $y = n^4$ , find each of the following when  $n = 1$

(1) The equation of the tangent to the curve of the function  $f$

(2)  $\frac{d^2 y}{dx^2}$

«  $2x - 3y - 7 = 0, \frac{1}{9}$  »

3 [a] Investigate the convexity of the curve of the function  $f$  where  $f(x) = |x^3 - 1|$  and show the inflection points if existed.

[b] If  $\int_{-2}^3 f(x) dx = 9$ ,  $\int_5^3 f(x) dx = 4$

, find the value of :  $\int_{-2}^5 [3f(x) - 6x] dx$

«  $-48$  »



- 4 [a] Find the area of the plane region bounded by the two curves :

$$y + x^2 = 6 \quad , \quad y + 2x - 3 = 0$$

«  $\frac{32}{3}$  square unit »

- [b] A right circular cylinder-like container of internal height 9 cm. and the interior radius length of its base is 6 cm. A metal rod of length 16 cm. is placed in the container. If the rate of sliding the rod away from the edge of the cylinder is 2 cm./sec. , find the rate of sliding the rod on the cylinder base when the rod reaches the end of its base.

«  $\frac{5}{2}$  cm./sec. »

- 5 [a] If the rate of change of the slope of the tangent to a curve at any point  $(x, y)$  on it is  $6(1 - 2x)$  and the curve has a critical point when  $x = 1$  and the function has a local minimum value equals 4

(1) Find the equation of the normal to the curve when  $x = -1$

(2) Sketch the curve of the function and show the maximum and minimum values and the inflection points if existed.

«  $x - 12y + 109 = 0$  »

- [b] Find the value of the solid generated by revolving the plane region bounded by the curves :  $y = x^3 + 1$  ,  $y = 0$  and  $x = 0$  ,  $x = 1$  a complete revolution about  $x$ -axis.

«  $\frac{23}{14} \pi$  cubic unit »




# Egypt Exams

(2017 : 2021 first and second sessions)

in

## Differential & Integral calculus







### Answer the following questions :

- 1 If the function  $f : f(x) = x + \frac{a}{x}$  has a critical point at  $x = 2$ , then the value of  $a = \dots\dots\dots$

(a) 4                      (b) 3                      (c) 2                      (d) 1

- 2 If the curve of the function  $f : f(x) = \cos x - a x^2$  has an inflection point at  $x = \frac{\pi}{3}$ , then the value of  $a = \dots\dots\dots$

(a)  $\frac{1}{4}$                       (b)  $-\frac{1}{4}$                       (c)  $\frac{1}{2}$                       (d)  $-1$

- 3 The absolute maximum value of the function  $f$  such that :  $f(x) = \sin x + \cos x$  in the interval  $[0, 2\pi]$  is  $\dots\dots\dots$

(a) zero                      (b)  $\frac{1}{\sqrt{2}}$                       (c) 1                      (d)  $\sqrt{2}$

### 4 Answer one of the following items :

[a] Determine the local maximum values and the local minimum values (if there exist) for the function  $f : f(x) = (2 - x) e^x$

[b] Find the absolute maximum value and the absolute minimum value of the function  $f$  such that :  $f(x) = 3x^4 - 4x^3$  in the interval  $[-1, 2]$

- 5  $\int 2 \cos^2 x \, dx = \dots\dots\dots$

(a)  $x + \frac{1}{2} \sin 2x + c$                       (b)  $x + 2 \sin 2x + c$   
(c)  $x - \frac{1}{2} \sin 2x + c$                       (d)  $x - \sin 2x + c$

- 6 In the orthogonal coordinate plane, the straight line  $\overleftrightarrow{AB}$  is drawn passing through the point  $C(3, 2)$ , cutting the positive part of  $x$ -axis at the point  $A$  and the positive part of  $y$ -axis at the point  $B$ , find the smallest area for  $\Delta AOB$  such that  $O$  is the origin point.

- 7 If  $f(x) = |x|$ , then  $\int_{-2}^2 f(x) \, dx = \dots\dots\dots$

(a) 4                      (b) 2                      (c) 0                      (d)  $-1$



8 Find the area of the region bounded by the two curves :  $y = x^2$  ,  $y = 5x$

9 Find the volume of the solid generated by revolving the region bounded by the two curves :  $y = x^2$  ,  $y = 3x$  a complete revolution about the  $x$ -axis

10 Answer one of the following items :

[a] Find :  $\int \frac{x}{x+1} dx$

[b] Find :  $\int x^2 \ln x dx$

11 If  $f(x) = a e^x$  , then  $f(-2) = \dots\dots\dots$

(a)  $-f(2)$

(b)  $-f(2)$

(c)  $-f(-2)$

(d)  $f(-2)$

12  $\int \frac{\ln x^2}{\ln x} dx = \dots\dots\dots$

(a)  $\frac{x}{2} + c$

(b)  $\frac{1}{x} + c$

(c)  $2x + c$

(d)  $\ln|x| + c$

13  $\int \cot x dx = \dots\dots\dots$

(a)  $\ln|\sin x| + c$

(b)  $\ln|\cos x| + c$

(c)  $-\ln|\sin x| + c$

(d)  $\ln|\csc x| + c$

14 Find the equation of the normal to the curve  $y = 3e^x$  at the point lying on it and its  $x$ -coordinate equals  $-1$

15 If  $y = \cot\left(\frac{\pi}{6}t\right)$  ,  $t = 3\sqrt{x}$  , then  $\left(\frac{dy}{dx}\right)_{x=1} = \dots\dots\dots$

(a)  $-\frac{\pi}{4}$

(b)  $-\frac{\pi}{9}$

(c)  $-\frac{\pi}{6}$

(d)  $\frac{\pi}{4}$

16 The slope of the tangent to the curve  $xy^2 = 3$  at the point  $(3, 1)$  equals  $\dots\dots\dots$

(a)  $-6$

(b)  $-3$

(c)  $-\frac{1}{6}$

(d)  $\frac{1}{3}$

17 If  $x = \frac{z+1}{z-1}$  ,  $y = \frac{z-1}{z+1}$  , find :  $\frac{d^2y}{dx^2}$  at  $z = 0$

18 If a stone fell in a settle water lake , then a circular wave is formed whose radius increases at a rate of 4 cm./sec. Find the rate of increasing of the surface area of the wave at the end of 5 seconds.





Answer the following questions :

1  $\int \sec^4 x \tan x \, dx = \dots\dots\dots$

- (a)  $\frac{1}{5} \sec^5 x + c$  (b)  $\frac{1}{4} \sec^4 x + c$  (c)  $\frac{1}{3} \tan x + c$  (d)  $\frac{-1}{3} \tan^3 x + c$

2 Find the maximum area for the isosceles triangle that could be drawn inscribed in a circle whose radius equals 12 cm.

3 If  $f(x) = \sin^3 x$ , then  $\int_{-\pi/2}^{\pi/2} f(x) \, dx = \dots\dots\dots$

- (a) 4 (b) 2 (c) zero (d) -1

4 Find the area of the region bounded by the two curves :  $y = x^2$  ,  $y = 4x$

5 Find the volume of the solid generated by revolving the region bounded by the two curves :  $y = x^2$  ,  $y = 2x$  a complete revolution about  $x$ -axis.

6 Answer one of the following items :

[a] Find :  $\int \frac{x}{3x^2 + 1} \, dx$

[b] Find :  $\int \frac{x}{e^{2x}} \, dx$

7 If  $y = \sec x$ , then  $y'(\frac{\pi}{3}) = \dots\dots\dots$

- (a)  $2\sqrt{3}$  (b) 6 (c) 8 (d) 14

8 If  $x = 2t^2 + 3$  ,  $y = \sqrt{t^3}$ , then  $(\frac{dy}{dx})_{t=1} = \dots\dots\dots$

- (a)  $\frac{3}{8}$  (b) 5 (c)  $\frac{8}{3}$  (d) 6

9 If  $y = x \sin x$ , prove that :  $x \frac{d^3 y}{dx^3} + x \frac{dy}{dx} + 2y = 0$

10 A rectangle of length 24 cm. and width 10 cm. , if its length shrinks at a rate of 2 cm./sec. while its width increases at a rate of 1.5 cm./sec. Find the rate of change of its area at the end of 4 seconds , after how many seconds does the area stop increasing ?



11  $\lim_{x \rightarrow 0} \frac{2^x - 1}{3x} = \dots\dots\dots$

(a)  $3 \ln 2$

(b)  $\frac{1}{3} \ln 2$

(c)  $\ln \frac{2}{3}$

(d)  $2 \ln 3$

12  $\int 4x e^{x^2+1} dx = \dots\dots\dots$

(a)  $e^{x^2+1} + c$

(b)  $4e^{x^2+1} + c$

(c)  $\frac{1}{2} e^{x^2+1} + c$

(d)  $2e^{x^2+1} + c$

13  $\int \frac{\ln x^2}{x \ln x^3} dx = \dots\dots\dots$

(a)  $x \ln \frac{1}{x} + c$

(b)  $\frac{2}{3 \ln x} + c$

(c)  $\frac{2}{3} \ln |x| + c$

(d)  $\frac{2}{3x \ln x} + c$

14 If  $y = (x^3 + 5)^x$ , find  $\frac{dy}{dx}$

15 If  $f: ]-1, 4[ \rightarrow \mathbb{R}$ ,  $f(x) = x^3 - 3x$ , then the number of the critical points for the function  $f$  equals .....

(a) zero

(b) 1

(c) 2

(d) 3

16 If the curve  $y = x^3 + ax^2 + bx$  has an inflection point at  $(3, -9)$ , then  $a + b = \dots\dots\dots$

(a) 15

(b) 6

(c) -9

(d) -12

17 The maximum value for the expression :  $4x - x^2$ , where  $x \in \mathbb{R}$  is .....

(a) 4

(b) 2

(c) 3

(d) 6

18 Answer one of the following items :

[a] Determine the maximum and the minimum local values for the function  $f$  such that :  $f(x) = x^3 - 3x^2 - 9x$ , then determine the inflection point (if exists) for the function.

[b] Find the absolute extrema values of the function  $f$  such that :

$f(x) = 10x e^{-x}$ ,  $x \in [0, 4]$





**Answer the following questions :**

1 If  $a^y = b^x$  such that  $a, b \in \mathbb{R}^+$ ,  $a \neq b$ , then  $\frac{dy}{dx} = \dots\dots\dots$

- (a)  $\log \frac{a}{b}$       (b)  $\log_a b$       (c)  $\log_b a$       (d)  $\log \frac{a}{b}$

2 If  $\int_{-2}^3 f(x) dx = 12$ ,  $\int_{-2}^5 f(x) dx = 16$ , then  $\int_3^5 f(x) dx = \dots\dots\dots$

- (a) -28      (b) -4      (c) 4      (d) 28

3 **Answer one of the following items :**

[a] Find :  $\int x^3 (x^2 + 1)^6 dx$

[b] Find :  $\int (x-3) e^{2x} dx$

4  $\int \tan \theta d\theta = \dots\dots\dots$

- (a)  $-\ln |\cos \theta| + c$       (b)  $-\ln \cos \theta + c$   
(c)  $\ln \cos \theta + c$       (d)  $|\ln \cos \theta| + c$

5  $-\pi \int \frac{2x - \sin x}{x^2 + \cos x} dx = \dots\dots\dots$

- (a)  $-\pi$       (b) zero      (c)  $\pi$       (d)  $2\pi$

6 **Answer one of the following items :**

[a] Find the local maximum values and the local minimum values of the function  $f : f(x) = x^3 - 3x - 2$ , and the inflection points of the curve of the function (if exists)

[b] Find the absolute extrema values of the function  $f : f(x) = x(x^2 - 12)$  in the interval  $[-1, 4]$

7 If  $f'(x) = x f(x)$  and  $f(3) = -5$ , then  $f''(3) = \dots\dots\dots$

- (a) -50      (b) 4      (c) 15      (d) 27

8 The curve of the function  $f : f(x) = (x-2)e^x$  is convex upwards in the interval  $\dots\dots\dots$

- (a)  $]-1, 2[$       (b)  $]-\infty, 0[$       (c)  $]0, \infty[$       (d)  $]0, 2[$



- 9 Find the equations of the tangent and the normal to the curve :

$$x = \sec \theta, \quad y = \tan \theta \text{ at } \theta = \frac{\pi}{6}$$

- 10 If  $\sin y + \cos 2x = 0$ , prove that :  $\frac{d^2 y}{dx^2} - \left(\frac{dy}{dx}\right)^2 \tan y = 4 \cos 2x \sec y$

- 11 If  $x = 2t^3 - 15t^2 + 36t + 1$ ,  $y = t^2 - 8t + 11$ , then this curve has a vertical tangent at  $t = \dots\dots\dots$

- (a) 4                      (b) 3 or 2                      (c) 6                      (d) 8

- 12 For the function  $f$  such that  $f'(x) = -2x + 6$ , then all of the following statements are correct except .....

- (a) the curve of the function  $f$  convex upwards in the interval  $]-\infty, \infty[$   
 (b) the function  $f$  has a local minimum value at  $x = 3$   
 (c) the curve of the function  $f$  has no inflection points.  
 (d)  $f(x)$  is decreasing in the interval  $]3, \infty[$

- 13 If  $y = ax^b$  such that  $a$  and  $b$  are constants, prove that :  $\frac{1}{y} \times \frac{dy}{dt} = \frac{b}{x} \times \frac{dx}{dt}$

- 14 Find the volume of the solid generated by revolving the region bounded by the curve  $y = x^2 + 2$ , the  $x$ -axis and the two straight lines  $x = -2$ ,  $x = 2$  a complete revolution about the  $x$ -axis.

- 15  $\lim_{x \rightarrow 0} \left( \frac{2^x - 1}{3x} \right) = \dots\dots\dots$

- (a)  $3 \ln 2$                       (b)  $\frac{1}{3} \ln 2$                       (c)  $\ln \frac{2}{3}$                       (d)  $2 \ln 3$

- 16 If  $f(x) = x(a - \ln x)$  such that  $a$  is constant, the curve of the function has a critical point at  $x = e$ , then  $a = \dots\dots\dots$

- (a) 1                      (b) 0                      (c)  $e$                       (d) 2

- 17 A metallic circular sector whose area is  $4 \text{ cm}^2$ . Find the radius length of the sector's circle which makes its perimeter as minimum as possible.  
What is the measure of its angle then ?

- 18 Find the area of the region bounded by the curve  $y = 4 - x^2$  and the straight line  $y = x + 2$





### Answer the following questions :

1 If  $f(x) = \sqrt{\sin 2x} - \csc x$ , then  $f'(\frac{\pi}{4}) = \dots\dots\dots$

- (a)  $\sqrt{2}$                       (b) 1                      (c) zero                      (d) -1

2 If the curve :  $y = (2x - a)^3 + 4$  has an inflection point at  $x = 5$ , then  $a = \dots\dots\dots$

- (a) 2                      (b) 4                      (c) 5                      (d) 10

3 A lake infected by bacteria has been treated by an antibacterial. If the number of bacteria  $z$  in  $1 \text{ cm}^3$  after  $n$  day is given by the relation  $z(n) = 20 \left( \frac{n}{12} - \ln \left( \frac{n}{12} \right) \right) + 30$  such that  $1 \leq n \leq 15$

- (1) When the number of bacteria be minimum during this interval ?  
(2) What is the least number of bacteria during this interval ?

4 Find the volume of the solid generated by revolving the region bounded by the two curves  $y = x^2$  and  $y = 3x - 2$  a complete revolution about the  $x$ -axis.

5 If  $y = e^{(1 + \ln x)}$ , then  $\frac{dy}{dx} = \dots\dots\dots$

- (a)  $x$                       (b)  $e^x$                       (c)  $e$                       (d) 1

6  $\int_1^1 \frac{x^3}{x^4 + \cos x} dx = \dots\dots\dots$

- (a) -1                      (b) zero                      (c) 1                      (d) 4

7 Answer one of the following items :

[a] Find :  $\int x(x+2)^6 dx$

[b] Find :  $\int (x+5)e^x dx$

8  $\int \frac{x+2}{x+1} dx = \dots\dots\dots$

(a)  $1 + \ln(x+1) + c$

(b)  $x - \ln|x+1| + c$

(c)  $x + \ln(x+1) + c$

(d)  $x + \ln|x+1| + c$



9  $\int_0^{\frac{\pi}{4}} \sec^2 x \tan x \, dx = \dots\dots\dots$

(a) zero

(b)  $\frac{1}{2}$ 

(c) 1

(d) 2

10 Answer one of the following items :

[a] Find the local maximum and minimum values (if found) of the function  $f$  :

$$f(x) = x^4 - 2x^2$$

[b] Find the absolute extrema values of the function  $f : f(x) = \frac{4x}{x^2 + 1}$  in the interval  $[-1, 3]$

11  $\lim_{x \rightarrow \infty} \left(1 + \frac{1}{x}\right)^{3x} = \dots\dots\dots$

(a) 1

(b) 3

(c) e

(d)  $e^3$ 

12 If the curve of the function  $f(x) = ax^2 + 12x + 1$  has a critical point at  $x = 2$ , then  $a = \dots\dots\dots$

(a) 12

(b) -3

(c) -1

(d) 3

13 Find the equations of the tangent and the normal to the curve :  $y = 3 + \sec x$  at the point which lies on the curve and its  $x$ -coordinate equals  $\frac{2\pi}{3}$

14 Find the area of the region bounded by the curve  $y = \sqrt{2x}$  and the straight line  $y = x$

15 If  $y = 2t^3 + 7$ ,  $z = t^2 - 4$ , then the rate of change for  $y$  with respect to  $z$  equals  $\dots\dots\dots$

(a)  $2t$ (b)  $3t$ 

(c) 6

(d) 12

16 The curve of the function  $f : f(x) = (x - 2)e^x$  is convex downwards in the interval  $\dots\dots\dots$

(a)  $]-\infty, \infty[$ (b)  $]-1, 2[$ (c)  $]0, 2[$ (d)  $]0, \infty[$ 

17 If  $\sin x = xy$ , prove that :  $x^2(y + \dot{y}) + 2 \cos x = 2y$

18 If  $xe^y = 2 - \ln 2 + \ln x$  and  $\frac{dx}{dt} = 6$  at  $x = 2$ ,  $y = 0$ , find  $\frac{dy}{dt}$





### Answer the following questions :

- 1 The volume of the solid generated by revolving the region enclosed by the curve  $y = 2x^2$  and the line  $y = 8x$  a complete revolution about the  $x$ -axis is equal to .....

- (a)  $\pi \int_0^8 (8x - 2x^2)^2 dx$       (b)  $\pi \int_0^4 (8x - 2x^2)^2 dx$   
 (c)  $\pi \int_0^4 (64x^2 - 4x^4) dx$       (d)  $\pi \int_0^4 (4x^4 - 64x^2)^2 dx$

- 2 The area of the region bounded by the curve  $y = x^3$  and the straight lines :  $y = 0$  and  $x = 2$  equals ..... unit of area.

- (a) 8      (b) 4      (c) 2      (d) 1

### 3 Answer only one of the following two questions :

[a] Use integration by parts to find :  $\int x^3 \sqrt{4-x^2} dx$

[b] Find :  $\int \sin^3 x dx$

- 4 The function  $f : f(x) = x^4 - 4x^2$  has .....

- (a) one local minimum value and two local maximum values.  
 (b) two different local minimum values and one local maximum value.  
 (c) two local minimum values and no local maximum values.  
 (d) two equal local minimum values and one local maximum value.

- 5 Let  $f$  be the function , defined by :  $f(x) = \frac{x}{\ln x}$  , then the local minimum value of  $f$  is .....

- (a)  $e$       (b)  $\frac{1}{e}$       (c)  $\ln e$       (d)  $-e$

### 6 Answer only one of the following two questions :

[a] Find the values of  $a$  and  $b$  such that the curve of the function  $y = x^3 + ax^2 + bx$  has an inflection point at  $(3, -9)$  , then determine the local maximum and local minimum values of the function.



[b] Find the absolute extrema values of the function  $f$ , where  $f(x) = 2x^2 e^x$ ,  $x \in [-3, 1]$

7  $\lim_{x \rightarrow 0} \frac{a^{2x} - 1}{x} = \dots\dots\dots$

(a)  $a^2$

(b)  $2a$

(c)  $2 \ln a$

(d)  $2 \ln a^2$

8 If  $y = (e^{-x} \ln x)$ , then  $\frac{dy}{dx} = \dots\dots\dots$

(a)  $e^{-x} \left( \frac{1}{x} - \ln x \right)$

(b)  $e^x \left( \frac{1}{x} - \ln x \right)$


(c)  $\frac{e^{-x}}{x} - \ln x$

(d)  $e^{-x} \left( \frac{1}{x} + \ln x \right)$

9 Find the equation of the tangent to the curve :

$y = 3x^2 - \ln x$  at the point  $(1, 3)$  which lies on it.

10 Determine the dimensions of the rectangle of largest area that can be inscribed in the right-angled triangle shown in the figure.



11 If  $y = \sec^n x$ , then  $\frac{dy}{dx} = \dots\dots\dots$

(a)  $n \sec^{n-1} x \tan x$

(b)  $ny \tan x$

(c)  $ny \cot x$

(d)  $ny$

12 The slope of the tangent to the curve :  $\cos(\sqrt{\pi y}) = 3x + 1$  at the point  $\left(\frac{-1}{3}, \frac{\pi}{4}\right)$ , equals .....

(a)  $\frac{-3x}{4}$

(b)  $0$

(c)  $3$

(d)  $-3$

13 As a spherical raindrop falls, it reaches a layer of dry air and begins to evaporate at a rate that is proportional to its surface area. Show that the radius of the raindrop decreases at a constant rate.

Given that : the area  $(A) = 4\pi r^2$ , the volume  $(v) = \frac{4}{3}\pi r^3$



14 If  $y = \frac{10 - \cos x}{x}$ , prove that :  $x \frac{d^2 y}{dx^2} + 2 \frac{dy}{dx} = \cos x$

15  $\int \frac{\ln x^3}{\ln x} dx = \dots\dots\dots$

- (a)  $3x + c$       (b)  $\frac{x}{3} + c$       (c)  $\frac{3}{x} + c$       (d)  $3x^2 + c$

16 Let  $f$  be the function given by :  $f(x) = \frac{x^4 + 1}{x^2}$ , then the function  $f$  is decreasing in .....

- (a)  $]-\infty, -1[$  only      (b)  $]-1, 0[$  and  $]1, \infty[$   
 (c)  $]0, 1[$  only      (d)  $]-\infty, -1[$  and  $]0, 1[$

17 If the slope of the tangent to a curve at any point  $(x, y)$  on it is  $(a \csc^2 x)$ , where  $a$  is constant, find the equation of this curve given that the curve passes through the two points  $(\frac{\pi}{4}, 5)$ ,  $(\frac{3\pi}{4}, 1)$

18 Find :  $\int_0^6 |x - 4| dx$  (write your steps)



Answer the following questions :

1 If  $y = \sec \frac{x}{4} + \sec \frac{\pi}{4}$ , then  $\frac{dy}{dx} = \dots\dots\dots$

(a)  $\frac{1}{4} \sec \frac{x}{4} \tan \frac{x}{4}$

(b)  $4 \sec \frac{x}{4} \tan \frac{x}{4}$

(c)  $\frac{1}{4} \sec \frac{x}{4} \tan \frac{x}{4} + \sqrt{2}$

(d)  $\frac{1}{4} \sec \frac{x}{4} \tan \frac{x}{4} + \frac{1}{4} \sec \frac{\pi}{4} \tan \frac{\pi}{4}$

2 The derivative of  $(x - \sin x)$  with respect to  $(1 - \cos x)$  at  $x = \frac{\pi}{3}$  equals  $\dots\dots\dots$

(a)  $\frac{1}{\sqrt{3}}$

(b)  $\frac{1}{2}$

(c)  $\sqrt{3}$

(d)  $\frac{\sqrt{3}}{2}$

3 ABC is a triangle, in which  $AC = 7$  cm.,  $BC = 3$  cm.,  $AB = x$  cm. and  $m(\angle ABC) = \theta$   
If  $\frac{d\theta}{dt} = 1.3$  rad.min. when  $\theta = \frac{\pi}{3}$ , find  $\frac{dx}{dt}$  at this instant.

4 If  $y = \sec x$ , **prove that** :  $y \frac{d^2 y}{dx^2} + \left(\frac{dy}{dx}\right)^2 = y^2 (3y^2 - 2)$

5  $\lim_{x \rightarrow 0} \frac{2^x - 1}{3x} = \dots\dots\dots$

(a)  $3 \ln 2$

(b)  $\frac{1}{3} \ln 2$

(c)  $\log \frac{2}{3}$

(d)  $2 \ln 3$

6 If  $y = \ln(1 + e^{2x})$ , then  $\frac{dy}{dx} = \dots\dots\dots$

(a)  $\frac{1}{1 + e^{2x}}$

(b)  $\frac{e^{2x}}{1 + e^{2x}}$

(c)  $\frac{2}{1 + e^{2x}}$

(d)  $\frac{2e^{2x}}{1 + e^{2x}}$

7 Find the equation of the tangent to the curve  $y = \ln[2 - \sqrt{2} \cos x]$  at the point which lies on it and its  $x$ -coordinate is  $\frac{\pi}{4}$

8 Find the positive number for which the sum of its multiplicative inverse and four times its square is the smallest possible.



9  $\int \tan X \, dX = \dots\dots\dots$

(a)  $\ln |\cos X| + c$

(b)  $-\ln |\sec X| + c$

(c)  $\sec^2 X + c$

(d)  $\ln |\sec X| + c$

10 Let  $f$  be the function given by :  $f(X) = (X^2 - 4)^{\frac{2}{3}}$ , then the function  $f$  is decreasing in .....

(a)  $]-\infty, -2[$  and  $]0, 2[$

(b)  $]-2, 0[$  and  $]2, \infty[$

(c)  $]-\infty, -2[$  only.

(d)  $]0, 2[$  only.

11 If the slope of the tangent to a curve at any point  $(X, y)$  on it is  $(X\sqrt{X+1})$ , find the equation of this curve given that the curve passes through the point  $(0, \frac{11}{15})$

12 If  $f : f(X) = \begin{cases} 2X + X^2 & , \text{ at } X < 0 \\ 2X - X^2 & , \text{ at } X \geq 0 \end{cases}$ , find  $_{-1}^3 \int f(X) \, dX$  (write your steps)

13 If the function  $f : f(X) = 2aX^2 + bX + 3$  has a local extrema at  $(1, 2)$ , then  $a + b = \dots\dots\dots$

(a)  $-1$

(b)  $\frac{5}{2}$

(c)  $-\frac{3}{2}$

(d)  $\frac{3}{2}$

14 If the graph of  $f''(X)$  (the second derivative of  $f$ )

is shown in the given figure for  $-2 \leq X \leq 4$

, then the graph of the function  $f$  is


convex upwards in .....

(a)  $-1 < X < 1$

(b)  $0 < X < 2$

(c)  $-2 < X < -1$  only.

(d)  $-2 < X < -1$  and  $1 < X < 3$





15 Answer only one of the following two questions :

[a] If the curve of the function :

$y = a x^3 + b x^2$  has an inflection point at  $(1, 4)$ , determine the values of  $a$  and  $b$ , then determine the local maximum and local minimum values of the function.

[b] Find the values of the absolute extrema of the function  $f(x) = 2x e^x$ ,  $x \in [-3, 1]$

16 The volume of the solid generated by revolving the region enclosed by the curve  $y = x^2$  and the line  $y = 3x$  a complete revolution about the  $x$ -axis is equal to .....

(a)  $\pi_0 \int^3 (3x - x^2)^2 dx$

(b)  $\pi_0 \int^3 (9x^2 - x^4) dx$

(c)  $\pi_0 \int^3 (x^4 - 9x^2) dx$

(d)  $\pi_0 \int^3 (x^2 - 3x)^2 dx$

17 The area of the region bounded by the straight lines :  $y = x$ ,  $x = 2$  and  $y = 0$  equals ..... unit of area.

(a)  $\frac{1}{2}$

(b) 1

(c) 2

(d) 4

18 Answer only one of the following two questions :

[a] Use integration by parts to find  $\int x^2 \ln x dx$

[b] Find  $\int x \sin(2x^2) dx$





### Answer the following questions :

1  $\int \frac{6x+9}{x^2+3x} dx = \dots\dots\dots$

(a)  $\ln |x^2 + 3x| + c$

(b)  $3 \ln |x^2 + 3x| + c$

(c)  $\frac{1}{3} \ln |x^2 + 3x| + c$

(d)  $3 \log |x^2 + 3x| + c$

2 The curve of the function  $f : f(x) = x^3 - 9x^2 - 120x + 6$  is convex downwards at  $x \in \dots\dots\dots$

(a)  $]10, \infty[ \cup ]-\infty, 4[$

(b)  $] -4, 10[$

(c)  $]3, \infty[$

(d)  $] -\infty, 3[$

3 Find the equation of the curve passes through the point  $(1, 0)$  and the slope of its tangent at any point on it equals  $x e^y$

4 Find  $-\frac{\pi}{2} \int^{\frac{\pi}{2}} (\sin x + \cos x)^2 dx$  (write your steps)

5 If the curve of the function  $f : f(x) = x^3 + kx^2 + 4$ ,  $k \in \mathbb{R}$  has an inflection point at  $x = 2$ , then  $k = \dots\dots\dots$

(a)  $-6$

(b)  $-3$

(c)  $6$

(d)  $9$

6 The function  $f$  such that  $f(x) = \frac{x}{x-1}$ ,  $x \in [2, 4]$  has  $\dots\dots\dots$

(a) an absolute maximum value at  $x = 4$

(b) an absolute minimum value at  $x = 2$

(c) an absolute minimum value at  $x = 1$

(d) an absolute maximum value at  $x = 2$

### 7 Answer only one of the following two questions :

[a] Identify the increasing and decreasing intervals and the local maximum and local minimum values of the function  $f : f(x) = 2x^3 - 9x^2 + 12x$

[b] Identify the convexity intervals upwards and downwards and the inflection points (if exists) of the function  $f : f(x) = x^4 - 6x^2 + 16$

8 If  $x = \tan \theta$ ,  $y = \sec \theta$ , then  $\frac{dy}{dx} = \dots\dots\dots$

(a)  $xy$

(b)  $\frac{x}{y}$

(c)  $\frac{y}{x}$

(d)  $\frac{x^2}{y^3}$



9  $\frac{dy}{dx} = \sec^2 x$ ,  $y = 3$  at  $x = \frac{\pi}{4}$ , then  $y = \dots\dots\dots$

(a)  $2 - \tan x$

(b)  $1 + \tan x$

(c)  $3 + \tan x$

(d)  $2 + \tan x$

10 Answer only one of the following two questions :

[a] Find :  $\int (x^2 + 1)\sqrt{x+2} dx$

[b] Find :  $\int x \sin x dx$

11  $\frac{d}{dx} [(\sec x - 1)(\sec x + 1)] = \dots\dots\dots$

(a)  $\sec^2 x \tan^2 x$

(b)  $2 \sec^2 x \tan x$

(c)  $2 \sec^2 x$

(d)  $\sec^2 x \tan x$

12 If  $xy^2 + 2xy = 8$ , then the value of  $y$  at the point  $(1, 2)$  equals  $\dots\dots\dots$

(a)  $-\frac{5}{2}$

(b)  $-\frac{1}{2}$

(c)  $-1$

(d)  $-\frac{4}{3}$

13 A plane fly horizontally at a height of 3000 m. from the surface of the ground and with velocity 480 km/h. to pass directly above an observer on the ground.

Find the rate of change of the distance between the plane and the observer after 30 sec.

14 If  $y = a e^{x^2+1}$  such that  $a$  is constant, prove that :  $\frac{d^3 y}{dx^3} = 4xy(3 + 2x^2)$

15  $\lim_{x \rightarrow \infty} \left(1 + \frac{1}{x}\right)^{x+3} = \dots\dots\dots$

(a)  $e$

(b)  $e^3$

(c)  $\frac{1}{3} e$

(d)  $3e$

16 If  $y = x^{\sin x}$ , then  $\frac{dy}{dx} = \dots\dots\dots$

(a)  $y \sin x \cos x$

(b)  $(\sin x)(x)^{\sin x - 1}$

(c)  $y \left( \frac{\sin x}{x} + \ln x \cos x \right)$

(d)  $\frac{y}{x} \sin x \cos x$

17 Find the equations of the two tangents of the curve :  $y = x^3 + 3x - 2$  which are perpendicular to the straight line :  $x + 6y = 1$

18 Find the greatest volume of the cuboid whose base is a square and its total surface area equals  $150 \text{ cm}^2$





Answer the following questions :

1  $\lim_{x \rightarrow \infty} \left(1 + \frac{5}{x}\right)^x = \dots\dots\dots$

- (a)  $5e$                       (b)  $e^5$                       (c)  $e^{-5}$                       (d)  $e^{\frac{1}{5}}$


2 If  $y = \ln |\sin x|$ , then  $\frac{dy}{dx} = \dots\dots\dots$

- (a)  $\tan x \ln 10$                       (b)  $\tan x$                       (c)  $\cot x \log e$                       (d)  $\cot x$

3 Find the equation of the two tangents to the curve :  $x^2 + y^2 = 8$  in which the tangents are perpendicular to the straight line  $y = 4 - x$

4 In the given figure :

ABCD is a rectangle in which :  $AB = 3 \text{ cm.}$  ,  $BC = 2 \text{ cm.}$   
 , a straight line is drawn passes through the point C and intersects  $\overline{AB}$  in E and  $\overline{AD}$  in F.



Find the smallest area of  $\triangle AEF$

5  $\int (\cos x e^{\sin x} + 3x^2) dx = \dots\dots\dots$

- (a)  $e^{\cos x} + x^3 + c$                       (b)  $-e^{\cos x} + x^3 + c$   
 (c)  $e^{\sin x} + x^3 + c$                       (d)  $-e^{\sin x} + x^3 + c$

6 The function  $f : f(x) = x^3 + 3x^2 - 9x$  has .....

- (a) a local a minimum value at the point  $(0, 0)$   
 (b) an inflection point at  $(1, -5)$   
 (c) an inflection point at  $(-1, 11)$   
 (d) a local a minimum value at the point  $(-3, 27)$

7 Find the equation of the curve passes through the point  $(1, 2)$  and the slope of its tangent at any point on it equals  $\frac{1+x}{xy}$ ,  $x \neq 0$ ,  $y \neq 0$



8 Find :  $\int_{-\pi}^{\pi} (4 + \pi \cos 2x) dx$  (write your steps)

9 If the curve of the function  $f : f(x) = x^3 + kx^2 + 4$ , has an inflection point at  $x = 1$ , then  $k$  equals .....

(a) 3

(b) 6

(c) -3

(d) -6

10 The function  $f : f(x) = x + \frac{1}{x}$ ,  $x \in [\frac{1}{2}, 3]$ , has .....

(a) an absolute minimum value at  $x = 1$ (b) an absolute minimum value at  $x = \frac{1}{2}$ (c) an absolute maximum value at  $x = -1$ (d) an absolute minimum value at  $x = 3$ 

11 Answer only one of the following two questions :

[a] Identify the increasing and decreasing intervals and the local maximum and local minimum values of the function  $f : f(x) = x^3 + 3x^2 - 9x - 7$

[b] Identify the convexity intervals upwards and downwards and the inflection points (if exists) of the function  $f : f(x) = x^3 - x^2$

12 If  $x = a \sec^2 \theta$ ,  $y = a \tan^3 \theta$ ,  $a$  is constant, then  $\frac{dy}{dx} = \dots\dots\dots$

(a)  $\tan \theta$ (b)  $\frac{3}{2} \tan \theta$ (c)  $\frac{3}{2} \tan \theta \sec \theta$ (d)  $\frac{3}{2} a \tan \theta$ 

13 If  $\int_2^5 f(x) dx = 4$ , then  $\int_2^5 (3f(x) - 1) dx = \dots\dots\dots$

(a) 12

(b) 11

(c) 9

(d) -8

14 Answer only one of the following two questions :

[a] Find :  $\int x(x+2)^6 dx$

[b] Find :  $\int x \ln |x| dx$

15 If  $y = \sec x + \tan x$ , then : .....

(a)  $\dot{y} + y \sec x = 0$ (b)  $\dot{y} - y \sec x = 0$ (c)  $\dot{y} + y \csc x = 0$ (d)  $\dot{y} - y \csc x = 0$



16 If  $x^2 + xy + y^3 = 0$ , then  $\frac{dy}{dx} = \dots\dots\dots$

(a)  $-\frac{2x+y}{x+3y^2}$

(b)  $-\frac{x+3y^2}{2x+y}$

(c)  $-\frac{2x+y}{x+3y^2-1}$

(d)  $\frac{-2x}{x+3y^2}$

17 A car moves from a fixed point in the north direction with velocity 30 km./h. and after one hour another car moves from the same point in the west direction with velocity 80 km./h.  
Find the rate of change of the distance between the two cars after one hour from the movement of the second car.

18 If  $y = e^{3x} + x^2$  Prove that :  $\frac{d^2y}{dx^2} = 9(y - x^2) + 2$



### Answer the following questions :

- 1 If  $x \in \mathbb{R}^+$ , then the smallest value of  $\sqrt{x} + \frac{1}{x}$  equals .....  
 (a)  $2\sqrt[3]{4}$                       (b)  $\frac{\sqrt[3]{4}}{2}$                       (c)  $\frac{3}{\sqrt[3]{4}}$                       (d)  $\frac{\sqrt[3]{4}}{3}$

---

- 2  $\int \frac{x}{x^2 - 2x + 1} dx = \dots + c$  (where c is constant)  
 (a)  $\ln|x-1| - \frac{1}{x-1}$                       (b)  $\ln|x-1| + \frac{1}{x-1}$   
 (c)  $\ln|x-1| + (x-1)^2$                       (d)  $\ln|x^2 - 2x + 1| + \frac{1}{x-1}$

---

- 3 If the function  $f : f(x) = \sin x + \frac{1}{2} \sin 2x, x \in ]0, \frac{\pi}{2}[$ , then the function f .....  
 (a) has a local minimum value at  $x = \frac{\pi}{4}$   
 (b) has a local maximum value at  $x = \frac{\pi}{6}$   
 (c) has not a local maximum value at this interval.  
 (d) has a local maximum value at  $x = \frac{\pi}{3}$

---

- 4 The absolute minimum value of the function :  $f(x) = \sqrt{x^2 + 9}$  where  $x \in [-3, 3]$  equals .....  
 (a)  $3\sqrt{2}$                       (b)  $-3\sqrt{2}$                       (c) 3                      (d) -3

---

- 5 f, g are polynomial functions,  $f(x) = cx^2 + g(x)$ ,  $g(1) = k$  and  $g'(1) = 6$  where c, k are constants. If (1, 5) is an inflection point to the curve of f, then  $k - c = \dots$   
 (a) -11                      (b) 5                      (c) 11                      (d) -5

---

- 6 The equation of the normal to the curve  $y = \ln(\tan x)$  at the point which lies on the curve and its x-coordinates equals  $\frac{\pi}{4}$  is .....  
 (a)  $4x - 8y = \pi$                       (b)  $8y + 4x = \pi$   
 (c)  $4x + 2y = \pi$                       (d)  $4x - 2y = \pi$

---

- 7 If f is continuous and even function in  $\mathbb{R}$  where  $_{-5}^5 f(x) dx = 2a$  and  $_0^3 f(x) dx = b$ , then  $_{-5}^{-3} f(x) dx = \dots$   
 (a)  $b - a$                       (b)  $2a - b$                       (c)  $a - b$                       (d)  $b - 2a$



8 If  $f(x) = x^3 - 3x - 4$ , then the function  $f$  is decreasing when .....

- (a)  $|x| < 1$  (b)  $|x| > 1$  (c)  $x > 1$  (d)  $x < -1$

9  $\int \frac{2e^x + 1}{e^x} dx = \dots + c$  (where  $c$  is constant)

- (a)  $2x + e^{-x}$  (b)  $2x - e^{-x}$  (c)  $2 + \ln e^x$  (d)  $2 - e^x$

10 The tangent to the curve :

$x = \cos 2\theta$  and  $y = \sin 3\theta$  at  $\theta = 0$  is .....

- (a) parallel to  $y$ -axis. (b) parallel to  $x$ -axis.  
(c) parallel to the straight line  $y = x$  (d) parallel to the straight line  $y = -x$

11 If  $P = f(t)$  where  $P, t \in \mathbb{R}^+$  and the rate of change of  $P$  with respect to  $t$  varies inversely with  $t$ ,  $f(1) = 200$  and  $f(\sqrt[3]{e}) = \frac{700}{3}$ , then  $f(3) = \dots$

- (a)  $100 \ln(3e^2)$  (b)  $200 \ln(9e)$  (c)  $100 \ln(e^2 + 3)$  (d)  $200 \ln(9 + e)$

12 If  $y = \sqrt{2(\sec x + \tan x)}$  where  $x \in ]0, \frac{\pi}{2}[$ , then  $\frac{1}{y} \left( \frac{dy}{dx} \right) = \dots$


- (a)  $2 \sec x$  (b)  $-\frac{1}{2} \sec x$  (c)  $\frac{1}{2} \sec x$  (d)  $-2 \sec x$

13  $\int \frac{2}{x \ln x^2} dx = \dots + c$  (where  $c$  is constant)

- (a)  $2 \ln |\ln x|$  (b)  $\frac{1}{2} \ln |\ln |x||$   
(c)  $\ln |\ln |x||$  (d)  $\ln |x|$

14 ABCDHN is a regular hexagon, the length of its side increases at rate  $\sqrt{3}$  cm/s, when the side length is 4 cm, then the rate of change of the shaded region with respect to time = ..... cm<sup>2</sup>/s.

- (a) 12 (b) 8  
(c)  $12\sqrt{3}$  (d)  $8\sqrt{3}$




15 If  $y = \frac{1}{(x+1)^2}$ , then the rate of change of  $\frac{1}{y}$  with respect to  $x^2$  at  $x = 1$  is .....

- (a)  $\frac{1}{2}$  (b) 4 (c) 2 (d)  $\frac{1}{4}$




- 16 If  $f(3)$  undefined,  $f(x) > 0$  when  $x \neq 3$ , then the curve which can represent the continues function  $f$  on  $[1, 5]$  is .....


(a)




(b)



(c)



(d)



- 17 If  $y = e^{x^3-4}$  and  $2\dot{y} = m x^2 y$  where  $x \neq 0$ , then  $m = \dots\dots\dots$

(a) 6

(b) 4

(c) 3

(d) 2

- 18 If the area of the region pounded by the two curves  $y = x^2$ ,  $y = kx$  equals  $\frac{9}{2}$  square units, where  $k > 0$ , then  $k = \dots\dots\dots$

(a) 9

(b) 3

(c) 6

(d) 12


- 19 The opposite figure represents the curves of the two functions  $y = ax - 2$ ,  $y = a\sqrt{x}$  if the area of shaded region equals  $\frac{13}{6}$  square units, then the value of the constant  $a = \dots\dots\dots$

(a)  $\frac{1}{2}$ 

(b) 1

(c)  $\frac{1}{3}$ 

(d) 2



- 20 The equation of the tangent to the curve :  $xy = x^y$  at the point  $(1, 1)$  which lies on it is .....

(a)  $x - 1 = 0$ (b)  $y + x = 2$ (c)  $y - 1 = 0$ (d)  $x - y = 0$




21 If  $x \frac{dy}{dx} - y = 3$  and  $x = 1$  when  $y = -2$ , then the relation between  $x$ ,  $y$  is .....

- (a)  $|y| = |x + 3|$  (b)  $|y + 3| = |x|$   
 (c)  $y + x = -1$  (d)  $y + 2x = 0$

22 If the volume of the solid generated by revolving the shaded region included between the curve  $xy = 3$ , the two straight lines  $y = 1$ ,  $y = k$  and  $y$ -axis complete revolution about  $y$ -axis equals  $6\pi$  cubic units, then  $k = \dots\dots\dots$

- (a) 4 (b) 2  
 (c) 3 (d) 1.5




23 If  $\lim_{x \rightarrow 0} \frac{e^{x+a} - e^a}{x} = \frac{1}{e}$ , then the value of  $a = \dots\dots\dots$

- (a) zero (b) 1 (c)  $e$  (d)  $-1$


24 The opposite figure represents the curve of the function  $f: f(x) = x^3 + a$ , if the area of the shaded region equals 68 square unit, then  $a = \dots\dots\dots$

- (a) 1 (b) 2  
 (c) 3 (d) 4



25 If the opposite figure represents the curve of the first derivative of a continuous function  $f$  whose domain is  $\mathbb{R}$ , then the wrong statement from the following is .....

- (a) The function has inflection point at  $x = 4$   
 (b) The function has local maximum value at  $x = 2$   
 (c) The curve of the function convex upward in  $]-\infty, 4[$  and convex downward in  $]4, \infty[$   
 (d)  $f(-3) < f(-2)$





### Answer the following questions :

- 1 If  $y = \cot 5x$ , then  $\frac{dy}{dx} + 5y^2 = \dots\dots\dots$   
 (a) 5 (b)  $-5y$  (c)  $5y$  (d)  $-5$

---

- 2 If  $\lim_{x \rightarrow 0} \frac{\ln(1+x)^{4a}}{x} = a^2 + 4$ , then the value of constant  $a = \dots\dots\dots$   
 (a)  $-2$  (b)  $-4$  (c)  $2$  (d)  $4$

---

- 3  $\int 4 \sin 2x e^{\cos 2x} dx = \dots\dots\dots + c$  where  $c$  is constant.  
 (a)  $-2e^{\cos 2x}$  (b)  $2e^{\cos 2x}$  (c)  $4e^{\cos 2x}$  (d)  $-4e^{\cos 2x}$

---

- 4 If  $\int \frac{2}{y} dy = \int \frac{1}{x} dx$ , then  $\ln y^2 = \dots\dots\dots + c$  where  $c$  is constant.  
 (a)  $\ln |x|$  (b)  $|x|$  (c)  $\ln |x+y|$  (d)  $\ln |x-y|$

---

- 5 If the function  $f : f(x) = x^2 + \frac{2a^3}{x}$  where  $a \in \mathbb{R}^-$ , then the function is decreasing in the interval  $\dots\dots\dots$   
 (a)  $]-\infty, 0[ , ]0, \infty[$  (b)  $]a, \infty[$   
 (c)  $]-\infty, a[$  (d)  $]-\infty, 0[ , ]a, 0[$

---

- 6 If  $z = x + 2$ , then  $\int \frac{x-2}{x^2+4x+4} dx = \dots\dots\dots + c$  where  $c$  is constant.  
 (a)  $\ln |z| - \frac{4}{z}$  (b)  $\ln |z| + \frac{4}{z}$  (c)  $\frac{-1}{2z^2} - \frac{4}{3z^3}$  (d)  $\frac{-1}{2z^2} + \frac{2}{3z^3}$

---

- 7 If  $3y^2 \frac{dy}{dx} = 2x + 1$  and  $x = 1$  when  $y = -1$ , then the relation between  $x, y$  is  $\dots\dots\dots$   
 (a)  $y^3 = x^2 + x - 3$  (b)  $y^3 = x^2 - 2$   
 (c)  $y^3 + x^2 = 0$  (d)  $y^3 + 1 = x^2 - x$

---


- 8 If  $x = \tan(1+y)$ , then  $\frac{dy}{dx} = \dots\dots\dots$  at  $x = 2$   
 (a)  $0.5$  (b)  $0.2$  (c)  $1$  (d)  $5$



- 9 The tangent of the curve  $2x + 1 = \sin y$  is parallel to y-axis at the point .....
- (a)  $(\frac{\pi}{2}, 0)$       (b)  $(-\frac{1}{2}, \pi)$       (c)  $(0, \frac{\pi}{2})$       (d)  $(-\frac{1}{2}, 0)$
- 
- 10 The slope of the normal to the curve  $x = e^t$ ,  $y = e^{2t+2}$  at the point which lies on the curve and its x-coordinate equals one is .....
- (a)  $-2e^2$       (b)  $\frac{-1}{2e^2}$       (c)  $2e^2$       (d)  $\frac{e^2}{2}$
- 
- 11 If  $x^2 = e^{2y}$ , then  $x \left( \frac{dy}{dx} \right) = \dots\dots\dots$
- (a)  $-1$       (b)  $1$       (c)  $x$       (d)  $e^{2x}$
- 
- 12 In the opposite figure :
- If  $AB = 5$  cm. , then the area of the figure ABCD is maximum when .....
- (a)  $x = y$   
 (b)  $x = 2y$   
 (c)  $y = 2x$   
 (d)  $3x = 4y$
- 
- 
- 13 The absolute maximum value of the function  $f : f(x) = x \ln x$  where  $x \in [e^{-2}, e]$  equals .....
- (a)  $-2e^{-2}$       (b)  $e$       (c)  $2e$       (d)  $-e^{-1}$
- 
- 14 If  $f$  is an even continuous function in  $\mathbb{R}$  and  $\int_0^3 f(x) dx = 5$ , then  $\int_{-3}^3 [f(x) - f(x)] dx = \dots\dots\dots$
- (a)  $15$       (b)  $-10$       (c)  $-15$       (d)  $10$
- 
- 15 The area of the region bounded by the curve  $y - 1 = e^x$  and x-axis and the straight lines  $x = 0$ ,  $x = k$  where  $k > 0$  equals ..... square unit.
- (a)  $e^k - k + 1$       (b)  $e^k + k + 1$       (c)  $e^k + k - 1$       (d)  $e^k - k - 1$

**16 In the opposite figure :**


Two squares having the same centre and their side lengths are 1 cm. and 4 cm. , if side length of the first square increases by rate 1 cm./sec. and side length of the second square decreases by rate  $\frac{1}{2}$  cm./sec. , then the area of the shaded region after  $\frac{1}{2}$  second .....



- (a) stop increasing instantly. (b) stop decreasing instantly.  
 (c) increases at rate  $\frac{1}{4}$  length unit/sec. (d) decreases at rate  $\frac{1}{4}$  length unit/sec.

**17 In the opposite figure :**


If  $f(x) = 3(x-1)^2$  ,  $_{-1} \int^4 g(x) dx = 150$   
 , then the area of shaded part equals ..... square units.



- (a) 131 (b) 123  
 (c) 119 (d) 115

**18 In the opposite figure :**

If the volume of the solid generated by revolving the shaded region complete revolution about y-axis equals  $8\pi$  cubic units , then  $a = \dots\dots\dots$



- (a) 8 (b) 16  
 (c) 24 (d) 4

**19** If the equation of the tangent to the curve :  $\log_y (4x) = 2x$  at the point (1 , 2) which lies on it is  $y = mx + c$  , then  $m = \dots\dots\dots$

- (a)  $1 - 2 \ln 2$  (b)  $2 \ln 2 - 1$  (c)  $2 \ln 2 + 1$  (d)  $2 + \ln 2$




20 If the function  $f : f(x) = 3ax^3 - bx - 5$  has local maximum value at  $x = 1$ , then  $\frac{b}{a} = \dots\dots\dots$

- (a) -9                      (b) 9                      (c) 20                      (d) -20


21 If  $f(x) = (x + 3)e^x$ , then the curve of the function  $f$  convex downward in the interval  $\dots\dots\dots$

- (a)  $]-\infty, -5[$                       (b)  $]-6, \infty[$                       (c)  $]-5, \infty[$                       (d)  $]-\infty, -3[$


22 If  $f$  is a continuous function has the following properties :  $f(3) = 1$ ,  $f'(x) < 0$  when  $x < 3$ ,  $f'(x) > 0$  when  $x > 3$  and  $f''(x) < 0$  for every  $x \neq 3$ , then the curve which can represent this function is  $\dots\dots\dots$




(a)



(b)



(c)



(d)

23 If  $y = f(x)$ ,  $\frac{d^2 x}{dy^2} = e^{y+1}$  and the tangent to the curve of the function  $f$  at the point  $(x, -1)$  which lies on it is parallel to the straight line :  $y = x - 3$ , then the slope of the tangent to the curve of the function  $f$  at any point  $(x, y)$  lies on it equals  $\dots\dots\dots$

- (a)  $e^{y+1}$                       (b)  $e^{2y+2}$                       (c)  $e^{-y-1}$                       (d)  $e^{-2y-2}$



- 24 The area bounded by the curve  $y = f(x)$  where  $y = n - \frac{n^2}{4}$ ,  $x = \frac{n}{2}$  and  $x$ -axis equal ..... square unit.


(a)  $\frac{7}{3}$

(b)  $\frac{3}{4}$

(c)  $\frac{4}{3}$

(d)  $\frac{3}{7}$

- 25 The opposite figure represents the curve of the continuous function  $g(x)$  on  $\mathbb{R}$  that has a horizontal tangent at  $x = 1$ , if  $f$  is differentiable function on  $\mathbb{R}$  where  $f'(x) = |g(x)|$ , then the true statement from the following is .....



- (a) the function  $f$  has an inflection point at  $x = -1$
- (b) the function  $f$  has minimum value at  $x = -1$
- (c) the function  $f$  has an inflection point at  $x = 1$
- (d) the curve of the function  $f$  convex upward in  $]-1, 1[$



# Al-Azhar Exams

(2019 : 2021 first and second sessions)

in

## Differential & Integral calculus







**Answer the following question :**

**1 Choose the correct answer from the given ones :**

(1) If  $f(x) = x \ln x$ , then  $\int_1^e f(x) dx = \dots\dots\dots$

- (a) 1                      (b) 2                      (c)  $\frac{e^2 - e - 1}{2}$                       (d)  $\frac{1 - e}{e}$

(2) The curve of the function  $f : f(x) = (x - 2)e^x$  is convex downwards on the interval  $\dots\dots\dots$

- (a)  $]-\infty, \infty[$                       (b)  $]-1, 2[$                       (c)  $]0, 2[$                       (d)  $]0, \infty[$

(3) A point moves on a curve whose equation is  $y^2 = 16x$ . If the rate of change of its  $x$ -coordinate with respect to time at  $y = 2$  equals  $\frac{5}{4}$  cm./sec., then the rate of change of its  $y$ -coordinate with respect to time at the same point equals  $\dots\dots\dots$  cm./sec.

- (a) 5                      (b) 10                      (c)  $\frac{4}{5}$                       (d)  $\frac{5}{16}$

(4) The volume of the solid generated by revolving the region bounded by the curve  $y = x^2$  and the straight line  $y = 2x$  complete revolution about  $x$ -axis =  $\dots\dots\dots \pi$  cube unit.

- (a)  $\frac{32}{5}$                       (b)  $\frac{32}{15}$                       (c)  $\frac{64}{15}$                       (d)  $\frac{64}{5}$

(5) If  $\frac{2}{\sqrt{x} + \sqrt{y}} = 9$ , then  $\left(\frac{dy}{dx}\right)^2 = \dots\dots\dots$

- (a)  $\frac{x}{y}$                       (b)  $\frac{y}{x}$                       (c)  $\frac{2y}{x}$                       (d)  $\frac{1}{2\sqrt{xy}}$

(6) If  $x = 2t^3 + 3$ ,  $y = t^4$ , then  $\frac{d^2y}{dx^2} = \dots\dots\dots$  at  $t = 1$

- (a)  $\frac{2}{3}$                       (b)  $\frac{1}{3}$                       (c) 4                      (d)  $\frac{1}{9}$

**Answer only three questions from the following :**

**2 [a] Find the two equations of the tangent and the normal to the curve  $x \sin 2y = y \cos 2x$  at the point  $\left(\frac{\pi}{4}, \frac{\pi}{2}\right)$**



[b] Calculate the area of the region bounded by the curve of the function

$$f : f(x) = 3x^2 + 1, x\text{-axis and the two straight lines } x = -1 \text{ and } x = 2$$

3 [a] If  $x^2 y = a \ln x$ , prove that :  $x^2 \ddot{y} + 5x \dot{y} + 4y = 0$  where  $a$  is constant.

[b] If the slope of the perpendicular to a curve at any point  $(x, y)$  lies on it equals  $\frac{-1}{a \csc^2 x}$  where  $a$  is constant, find the equation of the curve given that it passes

$$\text{through the two points } \left(\frac{\pi}{4}, 3\right) \text{ and } \left(\frac{3\pi}{4}, -1\right)$$

4 [a] In a perpendicular coordinate plane,  $\overleftrightarrow{AB}$  is drawn to pass through the point  $C(3, 2)$  and intersects the positive coordinate axes at point  $A$  and point  $B$ . Prove that the minimum area of triangle  $AOB$  equals 12 squared units where  $O$  is the origin point  $(0, 0)$ .

[b] Find :

$$(1) \int_0^{\ln 2} (e^{2x} + e^x) dx$$

$$(2) \int x^2 \ln x dx$$

5 [a] If the point  $(1, 12)$  is the inflection point to the curve of the function  $f$  where  $f(x) = ax^3 + bx^2$ , find the values of  $a$  and  $b$ , then determine the absolute extrema values of the function  $f$  on the interval  $[-1, 3]$ .

[b] Find :

$$(1) \int \frac{x+1}{\sqrt{x-1}} dx$$

$$(2) \int_0^2 \sqrt{4-x^2} dx$$



**Answer the following question :**

**1 Choose the correct answer :**

(1) The volume of the solid generated by revolving the region bounded by the two curves :  
 $y = x^2$  ,  $y = 1$  one complete revolution about y-axis is .....

- (a)  $\pi$                       (b)  $\frac{1}{2} \pi$                       (c)  $\frac{1}{4} \pi$                       (d)  $-\pi$

(2)  $\int_e^{e^3} \frac{1}{x-1} dx = \dots\dots\dots$

- (a)  $\ln(e-1)$                       (b)  $\ln(e^2 + e + 1)$                       (c)  $\ln(e^2 + e)$                       (d)  $\ln(e^3 - 1)$

(3)  $\lim_{x \rightarrow 1} \frac{\ln x}{x-1} = \dots\dots\dots$

- (a) 0                      (b) e                      (c) 1                      (d)  $e^{-1}$

(4) If  $x = (1-y)(1+y)(1+y^2)(1+y^4)$  , then  $\frac{dy}{dx} = \dots\dots\dots$

- (a)  $\frac{1}{8} y^7$                       (b)  $-8 y^7$                       (c)  $-\frac{1}{8} y^{-7}$                       (d)  $-\frac{1}{4} y^4$

(5) If  $y = \cot\left(\frac{\pi}{6} z\right)$  ,  $z = 3\sqrt{x}$  , then  $\frac{dy}{dx} = \dots\dots\dots$  at  $x = 1$

- (a)  $-\frac{\pi}{3}$                       (b)  $\frac{\pi}{36}$                       (c)  $-\frac{\pi}{4}$                       (d)  $-\frac{\pi}{36}$

(6) If the surface area of a sphere increases at a constant rate  $6 \text{ cm}^2/\text{sec}$ . when the radius length of the sphere equals 30 cm. , then the rate of increasing of the volume of sphere at this moment = .....  $\text{cm}^3/\text{sec}$ .

- (a) -18                      (b) 140                      (c) 90                      (d)  $90 \pi$

**Answer only three questions from the following :**

**2 [a]** Find the two equations of the tangent and the normal to the curve :  $x = \sec^2 \theta - 1$   
 $y = \tan \theta$  at  $\theta = \frac{\pi}{4}$

**[b]** Find the area of the region bounded by the curve of the function  $f : f(x) = \sqrt[3]{2x+2}$  ,  
the straight line  $x = 3$  and lies above x-axis.



- 3 [a] If the perimeter of a circular sector is 30 cm. and its area is as maximum as possible, find the radius length of its circle.

[b] If  $\int_{-2}^3 f(x) dx = 9$ ,  $\int_5^3 f(x) dx = 4$ , find the value of  $\int_{-2}^5 [3f(x) + 6x] dx$

- 4 [a] Find the local maximum, local minimum and the inflection points (if exists) of the function  $f : f(x) = x \ln x$

- [b] If the slope of the tangent to a curve at any point  $(x, y)$  lying on it is equal  $(5 - 2 \sec^2 2x)$ , find the equation of the curve known that the curve passes through the point  $(\frac{\pi}{8}, \frac{5\pi}{8})$

- 5 [a] If  $y = a e^{\frac{b}{x}}$ , **prove that** :  $xy \ddot{y} + 2y \dot{y} - x \dot{y}^2 = 0$   
 , where a and b are two non-zero constants.

[b] **Find :**

(1)  $\int 3x \sqrt{2x+3} dx$

(2)  $\int \frac{\ln 5x}{x} dx$



Answer the following question :

1 Choose the correct answer :

(1) If  $f(x) = 4x + \int 6 \cos^3 x \, dx$ , then  $f'(0) = \dots\dots\dots$

- (a) 10                      (b) 4                      (c) -2                      (d) 2

(2) If the slope of the tangent to the curve at any point  $(x, y)$  equals  $4e^{2x}$ ,  $f(0) = 2$ , then  $f(-2) = \dots\dots\dots$

- (a) 4                      (b)  $4e^{-4}$                       (c)  $2e^{-4}$                       (d)  $2e$

(3) If  $y = x^n$  where  $n$  is a natural number,  $\frac{d^3 y}{dx^3} = 120x^{n-3}$ , then  $n = \dots\dots\dots$

- (a) 7                      (b) 10                      (c) 6                      (d) 5

(4) The curve of the function  $f$  is convex downwards in  $\mathbb{R}$  if  $f(x) = \dots\dots\dots$

- (a)  $3 + x^4$                       (b)  $3 - x^2$                       (c)  $3 - x^3$                       (d)  $3 - x^4$

(5) A cuboid whose base dimensions are 9 cm. and 12 cm., if the rate of increase of its volume is  $27 \text{ cm}^3/\text{min.}$ , then the rate of change of its height =  $\dots\dots\dots \text{ cm./min.}$

- (a) 4                      (b)  $\frac{1}{4}$                       (c) 2                      (d)  $\frac{1}{2}$

(6)  $\int \frac{2}{x \ln 3 x} \, dx = \dots\dots\dots + c$

- (a)  $2 \ln |\ln 3 x|$                       (b)  $\frac{2}{3} \ln |\ln 3 x|$   
(c)  $6 \ln |\ln 3 x|$                       (d)  $\frac{2}{3} \ln |3 x|$

Answer only three questions from the following :

2 [a] Find the two equations of the tangent and the normal to the curve

$$x = \sec \theta, y = \tan \theta \text{ at } \theta = \frac{\pi}{3}$$

[b] Find :  $\int_0^5 |x - 3| \, dx$

3 [a] Find the equation of the curve  $y = f(x)$ , slope of the normal at any point on it is  $(4y + 3) \sec x$  known that the curve passes through the origin point.

[b] If  $y = a e^{\frac{b}{x}}$ , prove that :  $xy\ddot{y} + 2y\dot{y} - x(\dot{y})^2 = 0$



- 4 [a] An open field is bounded by a straight river from one of its sides. Determine how to place a fence around the other sides of the rectangle-like piece of land to surrounded as maximum area as possible by a 800 meter fence. What is the area of the land then ?

[b] If  $\int_1^4 f(x) dx = 7$  ,  $\int_4^1 g(x) dx = 3$   
 , calculate the value of  $\int_1^4 [f(x) + 2g(x) - 4] dx$

- 5 [a] If  $f(x) = (x-1)^4 + 3$  , determine the increasing and decreasing intervals of the function , then find the local maximum values , local minimum values and the inflection points if it exist.

[b] Find :

(1)  $\int x \sec^2 x dx$

(2)  $\int x^3 \sqrt{x-1} dx$



**Answer the following question :**

**1 Choose the correct answer :**

(1) If  $y^2 - 2\sqrt{x} = \text{zero}$ , then  $\frac{dy}{dx} = \dots\dots\dots$

(a)  $\frac{2y}{\sqrt{x}}$

(b)  $\sqrt{x}$

(c)  $\frac{x}{y^2}$

(d)  $\frac{1}{y^3}$

(2) If the curve of the function  $f$  has an inflection point when  $x = 2$  where  $f(x) = x^3 + kx^2 + 4$ , then the value of  $k = \dots\dots\dots$

(a)  $-6$

(b)  $-3$

(c)  $3$

(d)  $6$

(3) If  $\int_2^5 f(x) dx = 4$ , then  $\int_2^5 [3f(x) - 1] dx = \dots\dots\dots$

(a)  $9$

(b)  $11$

(c)  $12$

(d)  $-8$

(4) The sum of two positive integers is 5 and the sum of cubic of smaller number and double of square of the other is as minimum as possible, then the two numbers are represented by the set of elements  $\dots\dots\dots$

(a)  $\{2, 1\}$

(b)  $\{2, 3\}$

(c)  $\{4, 1\}$

(d)  $\left\{\frac{7}{2}, \frac{3}{2}\right\}$

(5) If the radius length of a circle increases at rate  $\frac{1}{\pi}$  cm./sec., the circumference of the circle increases at a rate of  $\dots\dots\dots$  cm./sec.

(a)  $\frac{2}{\pi}$

(b)  $2$

(c)  $\pi$

(d)  $2\pi$

(6) If  $x = \sin y$ , then  $\frac{d^2 y}{dx^2}$  at  $y = \frac{\pi}{4}$  equals  $\dots\dots\dots$

(a)  $1$

(b) undefined.

(c)  $\frac{1}{2}$

(d)  $2$

**Answer only three questions from the following :**

**2 [a] Find the two equations of the tangent and the normal to the curve of the function :**  
 $y = 3 - \cot^2 x$  at  $x = \frac{\pi}{4}$

**[b] Find each of the following :**

(1)  $\int_0^1 \frac{3e^x - 2e^{2x}}{2e^x} dx$

(2)  $\int \frac{(3x-1)^2}{3x} dx$



3 [a] If  $y^2 + 2xy = 8$ , **prove that** :  $(x + y) \frac{d^2 y}{dx^2} + 2 \frac{dy}{dx} + \left( \frac{dy}{dx} \right)^2 = \text{zero}$

[b] The slope of tangent to a curve at any point  $(x, y)$  on it equals  $\frac{2x+3}{x}$   
Find the equation of the curve if it passes through the point  $(e, 2e + 5)$

4 [a] A balloon rises vertically from point A on the ground surface. An apparatus is placed to follow up the motion of the balloon at point B at the same horizontal plane of point A and distance 200 meters from it. At a moment the apparatus observed the elevation angle of the balloon to find it  $\frac{\pi}{4}$ , and it increases at a rate of 0.12 rad./min. Find the rate of the balloon elevation at this moment.

[b] **Find each of the following :**

(1)  $\int x^2 \ln x \, dx$

(2)  $\int_0^5 |x - 2| \, dx$

5 [a] If  $f(x) = \frac{1}{3}x^3 - 9x + 3$  find the intervals of increasing and decreasing, then find the local maximum and minimum values.

[b] **Find the following integrations :**

(1)  $\int \frac{\cos^3 x - 5}{1 - \sin^2 x} \, dx$

(2)  $\int \frac{\ln 5x}{x} \, dx$



Answer the following questions :

1 Choose the correct answer :

(1) If the curve  $y = (2x + a)^3 + 5$  has an inflection point when  $x = 3$ , then  $a = \dots\dots\dots$

- (a) -3                      (b) 6                      (c) 9                      (d) -6

(2) If  $f(x) = e^{\ln(x^2 + 3x + 1)}$ , then  $\ddot{f}(1) = \dots\dots\dots$

- (a) 5                      (b) 2                      (c) -1                      (d) -4

(3) If  $y = \sec^2 \theta - 1$ ,  $x = \tan \theta$ , then  $\frac{d^2 y}{d x^2} = \dots\dots\dots$

- (a)  $2 \sec^2 \theta$                       (b) 2                      (c)  $\sec \theta \tan \theta$                       (d)  $\frac{1}{2}$

(4)  $-\frac{\pi}{2} \int^{\frac{\pi}{2}} (\sin x + \cos x)^2 dx = \dots\dots\dots$

- (a)  $\frac{\pi}{2}$                       (b)  $\frac{\pi}{2} + 1$                       (c)  $\frac{\pi}{2} - 1$                       (d)  $\pi$

(5) The volume of the solid generated by revolving a region bounded by the curve of the function  $y = \sqrt{x+1}$  and the straight lines  $y = 0$ ,  $x = 1$ ,  $x = -1$  a complete revolution about  $x$ -axis =  $\dots\dots\dots$

- (a)  $\pi$                       (b)  $\frac{3}{2} \pi$                       (c)  $2 \pi$                       (d)  $\frac{5}{2} \pi$

(6) If the volume of a cube increases regularly such that it keeps its shape at a rate  $27 \text{ cm}^3/\text{min.}$ , then the increase of the area of one of its faces at the moment which its edge length is  $3 \text{ cm.} = \dots\dots\dots \text{ cm./min.}$

- (a) 6                      (b) 3                      (c) 2                      (d) 1

Answer only three questions from the following :

2 [a] Find the two equations of the tangent and the normal for the curve of the function :  
 $y = 3 - \cot^2 x$  at  $x = \frac{\pi}{4}$

[b] Find :

(1)  $\int x^4 \ln x dx$

(2)  $\int x^2 \sqrt{x-3} dx$



3 [a] If  $\sin y + \cos 2x = 0$ , prove that :  $\frac{d^2 y}{dx^2} - \left(\frac{dy}{dx}\right)^2 \tan y = 4 \cos 2x \sec y$

[b] If the slope of the tangent to the curve of the function  $f$  at any point  $(x, y)$  lying on it is given by the relation  $\frac{3x+2}{x}$

Find the equation of the curve if it passes through the point  $(e, 3e + 5)$

4 [a] A cuboid-like box whose base is in the form of square. If the sum of lengths of all its edges equals 120 cm., find the dimensions of the box that will maximize its volume.

[b] If  $\int_{-2}^3 f(x) dx = 9$ ,  $\int_5^3 f(x) dx = 4$

, find the value of  $\int_{-2}^5 [3f(x) - 6] dx$

5 [a] If  $f(x) = e^x(3 - x)$  Find :

(1) The local maximum and minimum values of the function (if exist).

(2) The intervals of convexity upward and downward for the function and the inflection points (if exist).

[b] Find the area of the region bounded by the two curves  $y = x^2$  and  $y = 4x$  in square units.



Answer the following questions :

1 Choose the correct answer :

- (1) If  $f(x) = x^2 - 3 \ln 5x$ , then  $f'(2) = \dots\dots\dots$   
 (a) -1                      (b) 1                      (c)  $\frac{5}{2}$                       (d) 6
- (2) The curve of the function  $f: f(x) = 3x - x^3$  has a local maximum value at  $x = \dots\dots\dots$   
 (a) 1                      (b) -1                      (c) zero                      (d) 4
- (3) If  $x = \sin y$ , then  $\frac{d^2 y}{d x^2} = \dots\dots\dots$  when  $y = \frac{\pi}{4}$   
 (a) 2                      (b) 1                      (c)  $\frac{1}{2}$                       (d) undefined
- (4) The area of the region bounded by the curve  $y = x^3$ , and the two straight lines  $y = 0$ ,  $x = 2$  equals  $\dots\dots\dots$  square units.  
 (a) 1                      (b) 2                      (c) 4                      (d) 8
- (5) If  $y = x^x$  where  $x > 0$ , then  $\frac{dy}{dx} = \dots\dots\dots$  when  $x = e$   
 (a)  $e^e$                       (b)  $2e^e$                       (c)  $2e^2$                       (d)  $e$
- (6) If  $\frac{dy}{dx} = x + \frac{1}{x}$ ,  $y = \frac{1}{2}$  at  $x = 1$ , then  $y = \dots\dots\dots$  at  $x = e$   
 (a)  $e^2 - e$                       (b)  $\frac{e^2 - 1}{-2}$                       (c)  $\frac{e^2 + 1}{4}$                       (d)  $\frac{e^2}{2} + 1$

Answer only three questions from the following :

- 2 [a] If  $y = \sec x$ , prove that :  $y \frac{d^2 y}{d x^2} + \left( \frac{dy}{dx} \right)^2 = y^2 (3y^2 - 2)$   
 [b] Find the volume of the solid generated by revolving the curve  $y = \frac{4}{x}$  and the straight line  $x + y = 5$  a complete revolution about a-axis
- 3 [a] When inflating a spherical balloon with gas, the rate of increase in its volume was  $8\pi \text{ cm}^3/\text{sec}$ . when its radius length 4 cm., find at this moment :  
 (1) The rate of increasing the radius length.  
 (2) The rate of increasing the surface area.  
 [b] If  $\int_1^4 f(x) dx = 7$ ,  $\int_4^1 g(x) dx = 3$   
 , calculate the value of  $\int_1^4 [f(x) + 2g(x) - 4] dx$



4 [a] Sketch the curve  $f(x) = 5 + 3x - x^3$  and show the local maximum, minimum values and the inflection points (if exist).

[b] If the slope of the tangent to the curve of the function  $f$  at any point  $(x, y)$  on it equals  $7 - 2e^x$  and  $f(\ln 2) = 3$ , find  $f(x)$

5 [a] Find the two equations of tangent and normal for the curve :

$$x = t^2 + 4t, \quad y = 2t^2 \text{ when } t = 1$$

[b] Find :

(1)  $\int 2x \cos(2x - 1) dx$

(2)  $\int x(x - 2)^6 dx$

# Notes

Lined area for writing notes.



يُصرف مجاناً مع هذا الكتاب الجزء الخاص بالإجابات

## Question Bank & Practice Exams

3<sup>rd</sup>  
Sec  
2022

**NOW** At All Bookstores



### Question Bank & Practice Exams in :

- Algebra & Analytic Solid Geometry
- Statics
- Dynamics
- Physics
- Biology
- Chemistry
- Hello English
- French



**AL TALABA BOOKSTORE**

For printing, publication & distribution

El Faggala - Cairo - Egypt

Tel.: 02/ 259 340 12 - 259 377 91

e-mail: info@elmoasserbooks.com

www.elmoasserbooks.com

15014



/ElMoasserEG